

Taxa	Rainy Season															
	1 m				250 m				800 m				1,200 m			
	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF
<i>Haloptilus longicornis</i>	–	–	–	–	0.56	1.51	0.43	10.71	0.02	0.06	0.01	2.38	0.02	0.04	0.20	2.38
<i>Haloptilus</i> spp.	–	–	–	–	0.31	0.76	0.24	7.14	–	–	–	–	–	–	–	–
<i>Pseudhaloptilus</i> sp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Calanidae	11.93	27.78	1.08	11.90	0.31	0.81	0.24	8.33	0.13	0.37	0.06	3.57	0.08	0.17	0.94	3.57
<i>Calanoides carinatus</i>	0.27	1.30	0.02	1.19	1.70	7.00	1.32	8.33	16.17	30.56	7.87	16.67	0.13	0.36	1.57	5.95
<i>Mesocalanus tenuicornis</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Nannocalanus minor</i>	29.99	76.58	2.72	21.43	0.98	2.18	0.76	15.48	0.14	0.31	0.07	11.90	0.16	0.23	1.86	8.33
<i>Neocalanus gracilis</i>	–	–	–	–	0.01	0.06	0.01	1.19	–	–	–	–	–	–	–	–
<i>Neocalanus robustior</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Undinula vulgaris</i>	69.34	134.22	6.29	26.19	1.51	3.68	1.17	14.29	0.23	0.62	0.11	10.71	0.17	0.30	1.95	11.90
<i>Candacia bipinnata</i>	–	–	–	–	0.75	3.51	0.58	3.57	0.98	2.23	0.48	8.33	0.01	0.02	0.07	1.19
<i>Candacia bispinosa</i>	–	–	–	–	–	–	–	–	0.01	0.06	0.01	1.19	0.02	0.07	0.25	1.19
<i>Candacia ethiopica</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Candacia longimana</i>	–	–	–	–	–	–	–	–	0.003	0.01	0.001	1.19	0.02	0.07	0.25	1.19
<i>Candacia pachydactyla</i>	3.68	6.79	0.33	9.52	0.13	0.30	0.10	5.95	0.22	0.85	0.11	4.76	0.01	0.03	0.13	3.57
<i>Candacia simplex</i>	0.46	2.27	0.04	1.19	–	–	–	–	–	–	–	–	0.001	0.002	0.01	1.19
<i>Candacia tenuimana</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Candacia</i> spp.	0.66	2.23	0.06	3.57	0.06	0.25	0.05	3.57	0.08	0.25	0.04	4.76	0.04	0.09	0.50	4.76
Clausocalanidae	71.05	76.55	6.44	19.05	6.55	18.47	5.08	11.90	8.08	33.48	3.93	5.95	0.12	0.18	1.46	7.14
<i>Clausocalanus arcuicornis</i>	–	–	–	–	–	–	–	–	0.003	0.01	0.00	1.19	–	–	–	–
<i>Clausocalanus brevipes</i>	–	–	–	–	0.08	0.36	0.07	2.38	–	–	–	–	0.01	0.02	0.07	1.19
<i>Clausocalanus furcatus</i>	132.82	99.76	12.05	27.38	4.10	7.18	3.18	16.67	0.78	1.81	0.38	13.10	0.72	1.53	8.48	13.10
<i>Clausocalanus mastigophorus</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Ctenocalanus citer</i>	–	–	–	–	0.37	1.83	0.29	1.19	42.07	172.15	20.48	2.38	–	–	–	–
<i>Ctenocalanus vanus</i>	–	–	–	–	0.34	1.64	0.26	1.19	46.76	123.43	22.76	9.52	0.12	0.22	1.37	3.57
Centropagidae	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Centropages furcatus</i>	0.13	0.47	0.01	2.38	–	–	–	–	–	–	–	–	–	–	–	–
<i>Centropages violaceus</i>	1.16	4.63	0.11	2.38	–	–	–	–	–	–	–	–	–	–	–	–
Eucalanidae	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Eucalanus hyalinus</i>	–	–	–	–	–	–	–	–	0.02	0.06	0.01	2.38	–	–	–	–
<i>Paraeucalanus sewelli</i>	–	–	–	–	–	–	–	–	0.27	0.63	0.13	11.90	0.001	0.003	0.01	1.19
Euchaetidae	0.46	2.27	0.04	1.19	0.17	0.68	0.13	4.76	0.01	0.02	0.003	3.57	0.02	0.06	0.28	4.76
<i>Euchaeta marina</i>	0.83	3.20	0.08	2.38	–	–	–	–	–	–	–	–	0.02	0.06	0.28	2.38
<i>Euchaeta media</i>	–	–	–	–	0.02	0.08	0.02	2.38	–	–	–	–	0.001	0.002	0.01	1.19

Taxa	Rainy Season															
	1 m				250 m				800 m				1,200 m			
	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF
<i>Paraeuchaeta sarsi</i>	–	–	–	–	0.13	0.47	0.10	3.57	1.13	4.79	0.55	1.19	0.02	0.07	0.25	1.19
<i>Paraeuchaeta</i> sp.	–	–	–	–	–	–	–	–	–	–	–	–	0.002	0.01	0.03	1.19
Heterorhabdidae	–	–	–	–	0.03	0.11	0.02	2.38	0.01	0.04	0.01	2.38	0.01	0.03	0.14	2.38
<i>Heterorhabdus austrinus</i>	–	–	–	–	–	–	–	–	0.002	0.01	0.001	1.19	–	–	–	–
<i>Heterorhabdus papilliger</i>	–	–	–	–	0.03	0.11	0.02	2.38	0.01	0.02	0.004	2.38	–	–	–	–
<i>Heterorhabdus spinifrons</i>	1.12	4.95	0.10	3.57	0.42	0.78	0.33	14.29	0.06	0.19	0.03	7.14	0.04	0.07	0.42	4.76
<i>Heterorhabdus</i> spp.	–	–	–	–	0.02	0.10	0.02	1.19	–	–	–	–	–	–	–	–
<i>Paraheterorhabdus vipera</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Lucicutia clausii</i>	–	–	–	–	0.58	1.63	0.45	10.71	0.002	0.01	0.001	1.19	0.07	0.14	0.84	4.76
<i>Lucicutia flavicornis</i>	2.19	5.99	0.20	7.14	0.53	0.81	0.41	16.67	0.04	0.10	0.02	5.95	0.23	0.43	2.68	10.71
<i>Lucicutia gaussee</i>	–	–	–	–	0.03	0.10	0.02	2.38	0.001	0.01	0.001	1.19	0.002	0.01	0.02	1.19
<i>Lucicutia longicornis</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.001	0.003	0.02	2.38
<i>Lucicutia magna</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Lucicutia ovalis</i>	–	–	–	–	0.01	0.05	0.01	1.19	0.00	0.01	0.002	2.38	0.003	0.01	0.04	1.19
<i>Lucicutia wolfendeni</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.003	0.01	0.04	1.19
<i>Lucicutia</i> spp.	0.23	1.14	0.02	1.19	2.06	3.84	1.59	14.29	0.10	0.31	0.05	5.95	0.04	0.06	0.48	7.14
<i>Metridia brevicauda</i>	–	–	–	–	–	–	–	–	0.03	0.10	0.02	5.95	–	–	–	–
<i>Metridia princeps</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Metridia</i> spp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Pleuromamma abdominalis</i>	–	–	–	–	0.91	1.51	0.71	16.67	0.08	0.18	0.04	9.52	0.12	0.18	1.44	9.52
<i>Pleuromamma gracilis</i>	–	–	–	–	2.14	5.90	1.66	15.48	0.10	0.26	0.05	7.14	0.03	0.05	0.32	5.95
<i>Pleuromamma piseki</i>	0.50	1.49	0.05	3.57	0.08	0.20	0.06	4.76	0.02	0.07	0.01	2.38	–	–	–	–
<i>Pleuromamma xiphias</i>	–	–	–	–	0.01	0.05	0.01	1.19	0.003	0.01	0.002	2.38	0.001	0.002	0.01	1.19
<i>Pleuromamma</i> spp.	0.77	3.77	0.07	1.19	2.90	5.82	2.25	20.24	0.03	0.13	0.02	3.57	0.09	0.15	1.06	8.33
Paracalanidae	26.95	55.52	2.44	10.71	28.88	133.27	22.37	7.14	50.70	200.49	24.68	7.14	0.001	0.004	0.01	1.19
<i>Acrocalanus gracilis</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Acrocalanus longicornis</i>	50.28	44.69	4.56	27.38	0.59	1.76	0.46	9.52	0.27	0.63	0.13	8.33	0.05	0.13	0.63	4.76
<i>Acrocalanus</i> spp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Calocalanus contractus</i>	–	–	–	–	0.02	0.10	0.02	1.19	–	–	–	–	0.02	0.07	0.28	2.38
<i>Calocalanus pavo</i>	8.55	19.58	0.78	8.33	–	–	–	–	0.01	0.03	0.004	1.19	0.01	0.03	0.11	1.19
<i>Calocalanus pavoninus</i>	21.77	34.57	1.97	16.67	0.38	1.53	0.30	3.57	0.10	0.25	0.05	4.76	0.08	0.14	0.97	5.95
<i>Calocalanus</i> spp.	20.39	37.74	1.85	15.48	–	–	–	–	0.16	0.61	0.08	3.57	0.01	0.04	0.14	2.38
<i>Mecynocera clausi</i>	0.20	0.72	0.02	2.38	0.12	0.38	0.09	7.14	0.02	0.04	0.01	2.38	0.04	0.06	0.42	7.14
<i>Paracalanus aculeatus</i>	5.32	10.64	0.48	14.29	0.08	0.32	0.07	2.38	0.17	0.50	0.08	5.95	0.01	0.02	0.11	2.38

Taxa	Rainy Season															
	1 m				250 m				800 m				1,200 m			
	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF
<i>Paracalanus nanus</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Paracalanus parvus</i>	7.39	24.76	0.67	5.95	0.31	1.18	0.24	2.38	19.51	56.39	9.50	3.57	0.04	0.13	0.45	1.19
<i>Paracalanus quasimodo</i>	68.62	185.07	6.22	16.67	3.70	17.54	2.87	3.57	0.002	0.01	0.001	1.19	0.02	0.06	0.21	1.19
Phaennidae	–	–	–	–	0.17	0.70	0.13	4.76	0.02	0.04	0.01	4.76	0.002	0.01	0.03	1.19
<i>Xanthocalanus marlyae</i>	–	–	–	–	–	–	–	–	0.01	0.03	0.00	1.19	–	–	–	–
<i>Nullosetigera helgae</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Nullosetigera</i> sp.	–	–	–	–	0.02	0.05	0.01	2.38	–	–	–	–	–	–	–	–
Pontellidae	0.68	2.32	0.06	2.38	0.03	0.11	0.02	2.38	0.001	0.01	0.00	1.19	–	–	–	–
<i>Calanopia americana</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Calanopia</i> sp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Labidocera acutifrons</i>	0.96	2.43	0.09	4.76	0.04	0.19	0.03	1.19	–	–	–	–	–	–	–	–
<i>Labidocera fluviatilis</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Labidocera</i> spp.	0.87	2.94	0.08	2.38	–	–	–	–	–	–	–	–	–	–	–	–
<i>Pontellina plumata</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Pontellina</i> sp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Pontellopsis villosa</i>	0.01	0.03	0.00	1.19	–	–	–	–	–	–	–	–	–	–	–	–
<i>Rhincalanus cornutus</i>	1.36	5.06	0.12	2.38	0.14	0.38	0.11	5.95	0.03	0.06	0.01	4.76	0.01	0.02	0.09	3.57
<i>Rhincalanus nasutus</i>	–	–	–	–	0.15	0.44	0.12	5.95	0.05	0.23	0.03	1.19	–	–	–	–
<i>Rhincalanus</i> spp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Scolecitrichidae	1.83	5.30	0.17	4.76	0.21	0.72	0.17	7.14	0.08	0.20	0.04	5.95	0.01	0.05	0.18	2.38
<i>Amallothrix dentipes</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.01	0.04	0.14	1.19
<i>Amallothrix tenuiserrata</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Amallothrix</i> spp.	–	–	–	–	–	–	–	–	0.10	0.41	0.05	1.19	0.001	0.002	0.01	1.19
<i>Lophothrix frontalis</i>	–	–	–	–	0.03	0.10	0.02	2.38	0.01	0.02	0.003	2.38	0.01	0.02	0.10	2.38
<i>Lophothrix latipes</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Lophothrix quadrispinosa</i>	–	–	–	–	0.01	0.04	0.01	1.19	–	–	–	–	–	–	–	–
<i>Lophothrix</i> spp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Pseudoamallothrix ovata</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Pseudoamallothrix profunda</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Racovitzanus levis</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Scaphocalanus brevicornis</i>	–	–	–	–	–	–	–	–	0.002	0.01	0.001	1.19	–	–	–	–
<i>Scaphocalanus echinatus</i>	–	–	–	–	0.25	1.25	0.20	1.19	0.004	0.01	0.002	2.38	–	–	–	–
<i>Scaphocalanus elongatus</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Scaphocalanus magnus</i>	–	–	–	–	–	–	–	–	0.004	0.02	0.00	1.19	0.01	0.03	0.09	2.38

Taxa	Rainy Season															
	1 m				250 m				800 m				1,200 m			
	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF
<i>Scaphocalanus subbrevicornis</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Scaphocalanus</i> spp.	–	–	–	–	–	–	–	–	0.001	0.00	0.00	1.19	0.04	0.12	0.42	1.19
<i>Scolecithricella dentata</i>	–	–	–	–	0.20	0.81	0.16	3.57	0.01	0.02	0.00	1.19	–	–	–	–
<i>Scolecithricella minor</i>	–	–	–	–	1.27	2.71	0.98	8.33	0.04	0.13	0.02	3.57	–	–	–	–
<i>Scolecithricella</i> spp.	–	–	–	–	0.13	0.39	0.10	5.95	0.02	0.06	0.01	4.76	0.01	0.04	0.16	3.57
<i>Scolecithrix bradyi</i>	–	–	–	–	0.01	0.03	0.01	1.19	–	–	–	–	0.001	0.002	0.01	1.19
<i>Scolecithrix danae</i>	4.54	15.52	0.41	4.76	0.31	1.37	0.24	3.57	0.08	0.31	0.04	3.57	0.05	0.10	0.64	5.95
<i>Scolecithrix</i> spp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Scottocalanus securifrons</i>	–	–	–	–	0.03	0.09	0.02	3.57	–	–	–	–	–	–	–	–
Spinocalanidae	–	–	–	–	1.04	3.10	0.81	9.52	0.000	0.002	0.000	1.19	0.005	0.02	0.05	1.19
<i>Spinocalanus</i> sp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Subeucalanidae	–	–	–	–	0.09	0.46	0.07	1.19	0.01	0.02	0.00	2.38	0.003	0.01	0.04	1.19
<i>Subeucalanus crassus</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Subeucalanus longiceps</i>	–	–	–	–	0.01	0.04	0.01	1.19	0.001	0.01	0.00	1.19	0.004	0.01	0.05	1.19
<i>Subeucalanus pileatus</i>	–	–	–	–	–	–	–	–	0.10	0.41	0.05	1.19	–	–	–	–
<i>Subeucalanus subtenuis</i>	–	–	–	–	–	–	–	–	0.29	1.22	0.14	1.19	–	–	–	–
<i>Temora turbinata</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.02	0.07	0.22	1.19
<i>Temora stylifera</i>	80.97	109.87	7.34	26.19	3.38	8.58	2.62	16.67	0.43	0.80	0.21	13.10	0.10	0.19	1.21	9.52
<i>Temoropia mayumbaensis</i>	–	–	–	–	0.25	0.77	0.19	4.76	0.001	0.01	0.00	1.19	0.02	0.06	0.24	2.38
<i>Tharybis asymmetrica</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Oithona plumifera</i>	2.78	6.87	0.25	7.14	0.18	0.57	0.14	3.57	–	–	–	–	0.05	0.09	0.54	4.76
<i>Oithona setigera</i>	–	–	–	–	1.12	3.66	0.87	5.95	0.18	0.61	0.09	3.57	0.11	0.27	1.29	3.57
<i>Oithona similis</i>	3.25	10.18	0.29	4.76	6.71	10.61	5.20	25.00	0.25	0.45	0.12	5.95	0.33	0.49	3.84	10.71
<i>Oithona tenuis</i>	–	–	–	–	0.18	0.69	0.14	3.57	–	–	–	–	–	–	–	–
<i>Oithona</i> spp.	–	–	–	–	3.28	9.18	2.54	14.29	–	–	–	–	0.02	0.07	0.25	1.19
<i>Lubbockia squillimana</i>	–	–	–	–	1.07	2.22	0.83	20.24	0.04	0.08	0.02	7.14	0.06	0.15	0.69	7.14
<i>Lubbockia aculeata</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Lubbockia</i> spp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Oncaeidae	6.37	24.90	0.58	3.57	–	–	–	–	0.12	0.35	0.06	4.76	0.20	0.35	2.33	8.33
<i>Conaea rapax</i>	–	–	–	–	0.02	0.05	0.01	2.38	0.23	0.59	0.11	9.52	0.33	0.39	3.85	11.90
<i>Conaea</i> sp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Oncaea atlantica</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Oncaea media</i>	9.00	16.25	0.82	9.52	2.40	3.52	1.86	20.24	0.04	0.11	0.02	4.76	0.14	0.16	1.69	10.71
<i>Oncaea venusta</i>	80.64	75.70	7.32	28.57	12.90	20.78	9.99	27.38	2.34	3.57	1.14	16.67	1.15	1.98	13.55	14.29

Taxa	Rainy Season															
	1 m				250 m				800 m				1,200 m			
	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF	Mean	SD	RA	OF
<i>Oncaea</i> spp.	2.92	9.99	0.26	2.38	0.01	0.04	0.01	1.19	0.01	0.04	0.00	1.19	0.02	0.06	0.22	2.38
<i>Pachos</i> sp.	–	–	–	–	0.01	0.05	0.01	1.19	–	–	–	–	–	–	–	–
<i>Triconia</i> cf. <i>conifera</i>	5.26	22.76	0.48	2.38	2.59	3.41	2.00	19.05	2.11	4.89	1.03	15.48	0.30	0.53	3.48	11.90
Corycaeidae	9.02	20.05	0.82	9.52	0.31	0.96	0.24	7.14	0.14	0.61	0.07	1.19	0.10	0.25	1.23	7.14
<i>Agetus flaccus</i>	–	–	–	–	0.58	2.83	0.45	1.19	–	–	–	–	–	–	–	–
<i>Agetus limbatus</i>	5.51	17.15	0.50	3.57	0.51	1.06	0.40	10.71	0.20	0.68	0.10	5.95	0.12	0.30	1.46	7.14
<i>Agetus typicus</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Corycaeus speciosus</i>	4.91	10.10	0.45	13.10	0.15	0.68	0.12	3.57	0.06	0.23	0.03	3.57	0.19	0.43	2.18	3.57
<i>Onchocorycaeus giesbrechti</i>	24.60	36.49	2.23	19.05	0.11	0.27	0.08	5.95	0.005	0.02	0.002	1.19	0.07	0.20	0.81	3.57
<i>Onchocorycaeus latus</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Urocorycaeus lautus</i>	1.14	3.42	0.10	5.95	0.80	1.93	0.62	13.10	–	–	–	–	0.04	0.08	0.45	3.57
<i>Farranula gracilis</i>	204.93	222.99	18.59	28.57	3.88	9.51	3.01	16.67	0.60	0.82	0.29	13.10	0.59	0.94	6.89	10.71
<i>Farranula rostrata</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.02	0.07	0.25	1.19
<i>Copilia mirabilis</i>	0.24	1.00	0.02	2.38	–	–	–	–	0.00	0.02	0.00	1.19	0.01	0.03	0.11	1.19
<i>Sapphirina nigromaculata</i>	1.36	4.83	0.12	5.95	–	–	–	–	–	–	–	–	0.01	0.02	0.07	1.19
Harpacticoida	–	–	–	–	–	–	–	–	0.01	0.03	0.00	1.19	–	–	–	–
<i>Euterpina acutifrons</i>	–	–	–	–	–	–	–	–	–	–	–	–	0.001	0.002	0.01	1.19
<i>Macrosetella gracilis</i>	37.31	99.12	3.38	21.43	3.60	6.73	2.79	20.24	0.15	0.25	0.07	7.14	0.32	0.44	3.77	9.52
<i>Microsetella rosea</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Aegisthus mucronatus</i>	–	–	–	–	0.04	0.12	0.03	3.57	0.05	0.14	0.03	3.57	0.01	0.02	0.08	2.38
<i>Clytemnestra scutellata</i>	0.06	0.31	0.01	1.19	0.01	0.05	0.01	1.19	0.05	0.23	0.03	1.19	–	–	–	–
<i>Clytemnestra</i> sp.	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>Miracia efferata</i>	0.10	0.36	0.01	2.38	0.05	0.23	0.04	1.19	0.005	0.02	0.00	1.19	0.01	0.02	0.07	1.19
Damage Copepod	27.98	33.56	2.54	16.67	2.02	4.91	1.57	14.29	0.33	0.89	0.16	7.14	0.07	0.14	0.87	7.14
Nauplius Copepod	–	–	–	–	–	–	–	–	–	–	–	–	0.03	0.11	0.36	1.19