

Table S1. List of references including geographic records of *Trissolcus japonicus* and *T. mitsukurii* distribution.

Species name	List of references including geographic records of species distribution
<i>Trissolcus japonicus</i>	Matsuo et al. 2014, 2016, Talamas et al. 2015a, 2017, Herlihy et al. 2016, Jentsch 2016, Hedstrom et al. 2017, Nystrom Santacruz et al. 2017, Zhang et al. 2017, 2020, Zhong et al. 2017, Kaser et al. 2018, Sabbatini Peverieri et al. 2018, Abram et al. 2019, Jarrett et al. 2019, Kuki et al. 2019, Quinn et al. 2019, 2021, Stahl et al. 2019b, Boyle et al. 2020, Chen et al. 2020, 2021, Formella et al. 2020, Holthouse et al. 2020, 2021, McIntosh et al. 2020, Moraglio et al. 2020, Tillman et al. 2020, Zapponi et al. 2020, 2021, Avila et al. 2021, Dieckhoff et al. 2021, Ludwick et al. 2021, Mi et al. 2021, Nardi et al. 2021, Peterson et al. 2021
<i>Trissolcus mitsukurii</i>	Arakawa et al. 2004, Talamas et al. 2017, Sabbatini Peverieri et al. 2018, Chen et al. 2020, Moraglio et al. 2020, Scaccini et al. 2020, Zapponi et al. 2020, 2021, Bout et al. 2021, Rot et al. 2021

References

- Abram, P. K., Talamas, E. J., Acheampong, S., Mason, P. G., & Garipey, T. D. (2019). First detection of the samurai wasp, *Trissolcus japonicus* (Ashmead) (Hymenoptera, Scelionidae), in Canada. *Journal of Hymenoptera Research*, 68, 29. <https://doi.org/10.3897/JHR.68.32203>
- Arakawa, R., Miura, M., & Fujita, M. (2004). Effects of host species on the body size, fecundity, and longevity of *Trissolcus mitsukurii* (Hymenoptera: Scelionidae), a solitary egg parasitoid of stink bugs. *Applied Entomology and Zoology*, 39(1), 177–181. <https://doi.org/10.1303/AEZ.2004.177>
- Avila, G. A., Chen, J., Li, W., Alavi, M., Mi, Q., Sandanayaka, M., Zhang, F., & Zhang, J. (2021). Seasonal abundance and diversity of egg parasitoids of *Halyomorpha halys* in kiwifruit orchards in China. *Insects*, 12(5), 428. <https://doi.org/10.3390/insects12050428>
- Bout, A., Tortorici, F., Hamidi, R., Warot, S., Tavella, L., & Thomas, M. (2021). First detection of the adventive egg parasitoid of *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae) *Trissolcus mitsukurii* (Ashmead) (Hymenoptera: Scelionidae) in France. *Insects*, 12(9), 761. <https://doi.org/10.3390/INSECTS12090761>
- Boyle, S. M., Weber, D. C., Hough-Goldstein, J., & Hoelmer, K. A. (2020). Parental host species affects behavior and parasitism by the pentatomid egg parasitoid, *Trissolcus japonicus*

(Hymenoptera: Scelionidae). *Biological Control*, 149.

<https://doi.org/10.1016/J.BIOCONTROL.2020.104324>

Chen, H., Talamas, E. J., & Pang, H. (2020). Notes on the hosts of *Trissolcus* Ashmead (Hymenoptera: Scelionidae) from China. *Biodiversity Data Journal*, 8, e53786.

<https://doi.org/10.3897/BDJ.8.E53786>

Chen, M. N., Santander, R. D., Talamas, E. J., Jentsch, P. J., Bon, M. C., & Aćimović, S. G. (2021). Molecular identification of *Trissolcus japonicus*, parasitoid of the brown marmorated stink bug, by species-specific PCR. *Insects*, 12(5), 467. <https://doi.org/10.3390/INSECTS12050467>

Dieckhoff, C., Wenz, S., Renninger, M., Reißig, A., Rauleder, H., Zebitz, C. P. W., Reetz, J., & Zimmermann, O. (2021). Add Germany to the list — adventive population of *Trissolcus japonicus* (Ashmead) (Hymenoptera: Scelionidae) emerges in Germany. *Insects*, 12(5), 414.

<https://doi.org/10.3390/INSECTS12050414>

Formella, A., Dorman, S. J., Taylor, S. v., & Kuhar, T. P. (2020). Effects of aggregation lure and tree species on *Halyomorpha halys* (Hemiptera: Pentatomidae) seasonal oviposition. *Journal of Economic Entomology*, 113(1), 203–210. <https://doi.org/10.1093/JEE/TOZ281>

Hedstrom, C., Lowenstein, D. M., Andrews, H., Bai, B., & Wiman, N. G. (2017). Pentatomid host suitability and the discovery of introduced populations of *Trissolcus japonicus* in Oregon. *Journal of Pest Science*, 90(4), 1169–1179. <https://doi.org/10.1007/s10340-017-0892-6>

Herlihy, M. v., Talamas, E. J., & Weber, D. C. (2016). Attack and success of native and exotic parasitoids on eggs of *Halyomorpha halys* in three Maryland habitats. *PLOS ONE*, 11(3), e0150275. <https://doi.org/10.1371/JOURNAL.PONE.0150275>

Holthouse, M. C., Schumm, Z. R., Talamas, E. J., Spears, L. R., & Alston, D. G. (2020). Surveys in northern Utah for egg parasitoids of *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae) detect *Trissolcus japonicus* (Ashmead) (Hymenoptera: Scelionidae). *Biodiversity Data Journal*, 8, e53363. <https://doi.org/10.3897/BDJ.8.E53363>

- Holthouse, M. C., Spears, L. R., & Alston, D. G. (2021). Comparison of yellow and blue sticky cards for detection and monitoring parasitoid wasps of the invasive *Halyomorpha halys* (Hemiptera: Pentatomidae). *Journal of Insect Science*, *21*(5). <https://doi.org/10.1093/JISESA/IEAB062>
- Jarrett, B. J. M., Pote, J. M., Talamas, E. J., Gut, L. J., & Szqcs, M. (2019). The discovery of *Trissolcus japonicus* (Hymenoptera: Scelionidae) in Michigan. *Great Lakes Entomologist*, *52*, 5.
- Jentsch, P. J. (2016, September 19). *BMSB News: the invasive parasitic wasp, Trissolcus japonicus, recently found in New York State.* The Jentsch Lab. <https://blogs.cornell.edu/jentsch/2016/09/19/bmsb-news-the-invasive-parasitic-wasp-trissolcus-japonicus-recently-found-in-new-york-state/>
- Kaser, J. M., Akotsen-Mensah, C., Talamas, E. J., & Nielsen, A. L. (2018). First report of *Trissolcus japonicus* parasitizing *Halyomorpha halys* in North American agriculture. *Florida Entomologist*, *101*(4), 680–683. <https://doi.org/10.1653/024.101.0406>
- Kuki, D., Wada, Y., Tsunashima, A., & Itoyama, K. (2019). Developmental properties and parasitism capacity of the egg parasitoid *Trissolcus japonicus* (Hymenoptera: Scelionidae), reared on eggs of *Glaucias subpunctatus* (Hemiptera: Pentatomidae). *Applied Entomology and Zoology*, *54*(3), 307–312. <https://doi.org/10.1007/s10340-021-01415-x>
- Ludwick, D. C., Leake, L. B., Morrison, W. R., Lara, J. R., Hoddle, M. S., Talamas, E. J., & Leskey, T. C. (2021). Influence of holding conditions and storage duration of *Halyomorpha halys* (Hemiptera: Pentatomidae) eggs on adventive and quarantine populations of *Trissolcus japonicus* (Hymenoptera: Scelionidae) behavior and parasitism success. *Environmental Entomology*, *50*(3), 550–560. <https://doi.org/10.1093/EE/NVAA183>
- Matsuo, K., Hirose, Y., & Johnson, N. F. (2014). A taxonomic issue of two species of *Trissolcus* (Hymenoptera: Platygasteridae) parasitic on eggs of the brown-winged green bug, *Plautia stali* (Hemiptera: Pentatomidae): resurrection of *T. plautiae*, a cryptic species of *T. japonicus*. *Applied Entomology and Zoology*, *49*(3), 385–394. <https://doi.org/10.1007/s13355-014-0260-4>

- Matsuo, K., Honda, T., Itoyama, K., Toyama, M., & Hirose, Y. (2016). Discovery of three egg parasitoid species attacking the shield bug *Glaucias subpunctatus* (Hemiptera: Pentatomidae). *Japanese Journal of Applied Entomology and Zoology*, 60(1), 43–45. <https://doi.org/10.1303/jjaez.2016.43>
- McIntosh, H. R., Skillman, V. P., Galindo, G., & Lee, J. C. (2020). Floral resources for *Trissolcus japonicus*, a parasitoid of *Halyomorpha halys*. *Insects*, 11(7), 413. <https://doi.org/10.3390/INSECTS11070413>
- Mi, Q., Zhang, J. P., Haye, T., Zhang, B., Zhao, C., Lei, Y., Li, D., & Zhang, F. (2021). Fitness and interspecific competition of *Trissolcus japonicus* and *Anastatus japonicus*, egg parasitoids of *Halyomorpha halys*. *Biological Control*, 152. <https://doi.org/10.1016/J.BIOCONTROL.2020.104461>
- Moraglio, S. T., Tortorici, F., Pansa, M. G., Castelli, G., Pontini, M., Scovero, S., Visentin, S., & Tavella, L. (2020). A 3-year survey on parasitism of *Halyomorpha halys* by egg parasitoids in northern Italy. *Journal of Pest Science*, 93(1), 183–194. <https://doi.org/10.1007/s10340-019-01136-2>
- Nardi, F., Cucini, C., Cardaioli, E., Paoli, F., Sabbatini Peverieri, G., Roversi, P. F., Frati, F., & Carapelli, A. (2021). The complete mitochondrial genome of *Trissolcus japonicus* (Hymenoptera: Scelionidae), the candidate for the biological control of *Halyomorpha halys* (Hemiptera: Pentatomidae). *Mitochondrial DNA Part B*, 6(8), 2307–2309. <https://doi.org/10.1080/23802359.2021.1948370>
- Nystrom Santacruz, E., Venette, R., Dieckhoff, C., Hoelmer, K. A., & Koch, R. L. (2017). Cold tolerance of *Trissolcus japonicus* and *T. cultratus*, potential biological control agents of *Halyomorpha halys*, the brown marmorated stink bug. *Biological Control*, 107, 11–20. <https://doi.org/10.1016/J.BIOCONTROL.2017.01.004>
- Peterson, H. M., Talamas, E. J., Krawczyk, G., Sabbatini Peverieri, G., & Hoelmer, K. A. (2021). Survey for adventive populations of the samurai wasp, *Trissolcus japonicus* (Hymenoptera:

- Scelionidae) in Pennsylvania at commercial fruit orchards and the surrounding forest. *Insects*, 12(3), 258. <https://doi.org/10.3390/INSECTS12030258>
- Quinn, N. F., Talamas, E. J., Acebes-Doria, A. L., Leskey, T. C., & Bergh, J. C. (2019). Vertical sampling in tree canopies for *Halyomorpha halys* (Hemiptera: Pentatomidae) life stages and its egg parasitoid, *Trissolcus japonicus* (Hymenoptera: Scelionidae). *Environmental Entomology*, 48(1), 173–180. <https://doi.org/10.1093/EE/NVY180>
- Quinn, N. F., Talamas, E. J., Leskey, T. C., & Bergh, J. C. (2021). Seasonal captures of *Trissolcus japonicus* (Ashmead) (Hymenoptera: Scelionidae) and the effects of habitat type and tree species on detection frequency. *Insects*, 12(2), 118. <https://doi.org/10.3390/INSECTS12020118>
- Rot, M., Maistrello, L., Costi, E., Bernardinelli, I., Malossini, G., Benvenuto, L., & Trdan, S. (2021). Native and non-native egg parasitoids associated with brown marmorated stink bug (*Halyomorpha halys* [Stål, 1855]; Hemiptera: Pentatomidae) in western Slovenia. *Insects*, 12(6), 505. <https://doi.org/10.3390/INSECTS12060505>
- Sabbatini Peverieri, G., Talamas, E. J., Bon, M. C., Marianelli, L., Bernardinelli, I., Malossini, G., Benvenuto, L., Roversi, P. F., & Hoelmer, K. A. (2018). Two Asian egg parasitoids of *Halyomorpha halys* (Stål) (Hemiptera, Pentatomidae) emerge in northern Italy: *Trissolcus mitsukurii* (Ashmead) and *Trissolcus japonicus* (Ashmead) (Hymenoptera, Scelionidae). *Journal of Hymenoptera Research*, 67(67), 37–53. <https://doi.org/10.3897/JHR.67.30883>
- Scaccini, D., Falagiarda, M., Tortorici, F., Martinez-Sañudo, I., Tirello, P., Reyes-Domínguez, Y., Gallmetzer, A., Tavella, L., Zandigiaco, P., Duso, C., & Pozzebon, A. (2020). An Insight into the role of *Trissolcus mitsukurii* as biological control agent of *Halyomorpha halys* in northeastern Italy. *Insects*, 11(5), 306. <https://doi.org/10.3390/INSECTS11050306>
- Stahl, J., Tortorici, F., Pontini, M., Bon, M. C., Hoelmer, K. A., Marazzi, C., Tavella, L., & Haye, T. (2019). First discovery of adventive populations of *Trissolcus japonicus* in Europe. *Journal of Pest Science*, 92(2), 371–379. <https://doi.org/10.1007/s10340-018-1061-2>

- Talamas, E. J., Buffington, M. L., & Hoelmer, K. A. (2017). Revision of Palearctic *Trissolcus* Ashmead (Hymenoptera, Scelionidae). *Journal of Hymenoptera Research*, 56, 3. <https://doi.org/10.3897/JHR.56.10158>
- Talamas, E. J., Johnson, N. F., & Buffington, M. L. (2015). Key to Nearctic species of *Trissolcus* Ashmead (Hymenoptera, Scelionidae), natural enemies of native and invasive stink bugs (Hemiptera, Pentatomidae). *Journal of Hymenoptera Research*, 43, 45–110. <https://doi.org/10.3897/JHR.43.8560>
- Tillman, G., Toews, M. D., Blaauw, B. R., Sial, A. A., Cottrell, T. E., Talamas, E. J., Buntin, D. G., Joseph, S. v., Balusu, R. R., Fadamiro, H., Lahiri, S., & Patel, D. (2020). Parasitism and predation of sentinel eggs of the invasive brown marmorated stink bug, *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae), in the southeastern US. *Biological Control*, 145, 104247. <https://doi.org/10.1016/J.BIOCONTROL.2020.104247>
- Zapponi, L., Bon, M. C., Fouani, J. M., Anfora, G., Schmidt, S., & Falagiarda, M. (2020). Assemblage of the egg parasitoids of the invasive stink bug *Halyomorpha halys*: insights on plant host associations. *Insects*, 11(9), 588. <https://doi.org/10.3390/INSECTS11090588>
- Zapponi, L., Tortorici, F., Anfora, G., Bardella, S., Bariselli, M., Benvenuto, L., Bernardinelli, I., Butturini, A., Caruso, S., Colla, R., Costi, E., Culatti, P., di Bella, E., Falagiarda, M., Giovannini, L., Haye, T., Maistrello, L., Malossini, G., Marazzi, C., ... Sabbatini Peverieri, G. (2021). Assessing the distribution of exotic egg parasitoids of *Halyomorpha halys* in Europe with a large-scale monitoring program. *Insects*, 12(4), 316. <https://doi.org/10.3390/INSECTS12040316>
- Zhang, J. P., Zhang, F., Garipey, T. D., Mason, P. G., Gillespie, D., Talamas, E. J., & Haye, T. (2017). Seasonal parasitism and host specificity of *Trissolcus japonicus* in northern China. *Journal of Pest Science*, 90(4), 1127–1141. <https://doi.org/10.1007/s10340-017-0863-y>
- Zhang, J. P., Zhong, Y. Z., Tang, R., Rebijith, K. B., Li, F., Chen, G. H., & Zhang, F. (2020). Olfactory reception of host alarm pheromone component by the odorant-binding proteins in the

samurai wasp, *Trissolcus japonicus* (Hymenoptera: Scelionidae). *Frontiers in Physiology*, *11*, 1058. <https://doi.org/10.3389/fphys.2020.01058>

Zhong, Y. Z., Zhang, J. P., Ren, L. L., Tang, R., Zhan, H. X., Chen, G. H., & Zhang, F. (2017). Behavioral responses of the egg parasitoid *Trissolcus japonicus* to volatiles from adults of its stink bug host, *Halyomorpha halys*. *Frontiers in Ecology and Evolution*, *9*, 548. <https://doi.org/10.1007/S10340-017-0884-6>