

Floristic composition in ecotone forests in northern Brazilian Amazonia: preliminary data

Williamar Rodrigues Silva[‡], Carlos Darwin Angulo Villacorta[§], Ricardo Oliveira Perdiz[‡], Hugo Leonardo S. Farias[‡], Andressa Sampaio Oliveira[‡], Arthur Camurça Citó[¶], Lidiany Camila Silva Carvalho[#], Reinaldo Imbrozio Barbosa[¶]

[‡] UFRR/PRONAT, Boa Vista-RR, Brazil

[§] Centro de Investigación, Enseñanza y Producción Agroforestal - CEPIAGRY, Yurimaguas, Peru

[|] INPA/PPGBOT, Manaus-AM, Brazil

[¶] INPA/NAPRR, Boa Vista-RR, Brazil

[#] University of Exeter, Exeter, United Kingdom

Corresponding author: Reinaldo Imbrozio Barbosa (reinaldo@inpa.gov.br)

Academic editor: Giannantonio Domina

ZooBank: [urn:lsid:zoobank.org:pub:9AFA35F3-ED8B-42BD-B669-F185C3995B40](https://www.zoobank.org/pub:9AFA35F3-ED8B-42BD-B669-F185C3995B40)

Abstract

Background

Ecotone has been defined as “a *multi-dimensional environmentally stochastic interaction zone between ecological systems with characteristics defined in space and time, and by the strength of the interaction*” (Hufkens et al. 2009). This is a known concept to define transitional zones between two or more ecological communities, ecosystems or biotic regions. Ecotone forests, dispersed in northern Brazilian Amazonia, are natural formations which have been largely affected by anthropogenic impacts, such as deforestation and fire. Maracá Ecological Station, State of Roraima, Brazil, is a protected area with extensive representations of ecotone forests in this region of the Amazonia. Forest inventories and floristic surveys are important as they extend our knowledge (1) of forest structure and tree species composition and (2) of tree and palm species ecology in this region of the Amazonia. Both improve our ability to predict changes in plant diversity, considering the future scenarios of climate change in comparison with previous surveys performed in Maracá.

New information

We present a forest inventory carried out in 129 plots (10 m x 50 m; 6.45 ha in total) dispersed in a grid (5 km x 5 km) located in a forest zone ecotone in the eastern part of Maracá Ecological Station. All stems (tree + palm) with diameter at breast height ≥ 10 cm were recorded, identified and measured. A total of 3040 stems were recorded (tree =

2815; palm = 225), corresponding to 42 botanic families and 140 identified species. Seven families and 20 genera contained unidentified taxa (12.2%). Sapotaceae (735 stems; 10 species), Leguminosae (409; 24) and Rubiaceae (289; 12) were the most abundant families. *Peltogyne gracilipes* Ducke (Leguminosae), *Pradosia surinamensis* (Eyma) T.D.Penn. (Sapotaceae) and *Ecclinusa guianensis* Eyma (Sapotaceae) were the species with the highest importance value index (~ 25%). The dominance ($\text{m}^2 \text{ha}^{-1}$) of these species corresponds to > 36% of the total value observed in the forest inventory. Our dataset provides complementary floristic and structure information on tree and palm in Maracá, improving our knowledge of this Amazonian ecotone forest.

Keywords

Amazon, Brazil, forest inventory, Maracá Island, palms, Roraima, tree species

Introduction

The Pan-Amazon or Continental Amazon has the largest tropical forest area on the planet with > 6 million km^2 (MapBiomas 2019, WWF 2019). Estimates based on > 1,100 permanent plots scattered throughout the region indicate that the tree richness ranges between 7-10,000 species occupying a great diversity of habitats (Cardoso et al. 2017, ter Steege et al. 2019). Brazilian Amazonia accounts for the largest physical area of this region (Salati and Vose 1984; > $5 \times 10^6 \text{ km}^2$). However, it has been long threatened by a series of anthropogenic impacts, such as the replacement of native forest by pasture and soybean planting (Almeida et al. 2016, Fearnside 2006), combined with an increase in burned areas of primary and secondary forests (Alencar et al. 2015, Aragão et al. 2018, Barni et al. 2015). Modern anthropogenic activities, associated with global warming, have a negative effect on tree diversity and reduce the conservation status of Amazonian forests (Barlow et al. 2016, Esquivel-Muelbert et al. 2019). Despite having large areas of ombrophilous forests, the ecotone forests are important ecological areas because they occupy the peripheral zone to the Amazon basin (Central Amazonia) and are considered as the forest formations most impacted by anthropogenic activities in the Southern (Marques et al. 2019, Nogueira et al. 2015) and Northern (Barni et al. 2015, Santos et al. 2013) Amazonian "arcs of deforestation". The accelerated increase of anthropogenic activities within the Amazonian ecotones has been a major contributor to the fact that the region as a whole is now approaching to the "tipping point", limiting forest growth and potentially favouring low-density environments when compared to those currently supported by the region (Lovejoy and Nobre 2018).

Several floristic surveys and forest inventories have been carried out in these peripheral regions, especially from 1980-1990 (Nelson and Oliveira 2001). However, this period was insufficient to accumulate realistic information on forest structure and floristic in view of the continuous advance of anthropogenic activities. This impedes the reliable characterisation of plant diversity in these peripheral Amazonian regions. One such area is the State of Roraima, located in the northern part of the Brazilian Amazonia. This

~225,000 km² area contains ecotone forest zones of great geocological importance because they are located between Guiana Shield (highland savannas and tropical dry forests) and Central Amazonia (lowland tropical forests) (Barbosa and Bacelar-Lima 2008, Oliveira et al. 2017), which gives them a unique phytogeographical and ecoevolutionary history, where a high degree of endemism is observed and unique vegetation sets exist (Nascimento and Proctor 1997, Milliken and Ratter 1998). However, the region has received few forest inventories and floristic surveys (Suppl. material 1); main of them are associated with the monodominant forests (Nascimento and Proctor 1997, Nascimento et al. 1997), campinas and campinaranas (Barbosa and Ferreira 2004, Damasco et al. 2013) and forest fragments dispersed in savannas (Sette-Silva 1993, Santos et al. 2013, Jaramillo 2015).

In an attempt to expand studies on plant diversity in this part of the Amazon, two PPBio (Biodiversity Research Program, <https://ppbio.inpa.gov.br>) 25-km² research grids were installed in areas defined as ecotone forests in Roraima taking into account the Brazilian Vegetation Classification System (Brazil-IBGE 2012): Maracá Ecological Station and Viruá National Park. Both are protected areas under Brazilian government management. Surveys of structure (vertical and horizontal) and tree species composition of the main forest types of these research grids have been expanded in Maracá (Nascimento et al. 2014, Nascimento et al. 2017, Villacorta 2017) and Viruá (Damasco et al. 2013, Barbosa et al. 2017). Accordingly, the current study provides preliminary data from a forest inventory carried out in the Maracá research grid. These data expand the scale of floristic and structural observations in this northern Amazonian ecotone zone from those initiated by Milliken and Ratter (1998) and Nascimento et al. (1997).

Project description

Title: *"Tree growth and mortality in Roraima ecotonal forests"* (Crescimento e mortalidade de árvores em florestas ecotonais de Roraima). The dataset is associated with the PhD thesis of Williamar Rodrigues Silva (Analysis of environmental conditions and climate variability on tree growth and mortality in ecotonal forests of Maracá Island, Roraima), PRONAT/UFRR, Boa Vista, Roraima, Brazil.

Personnel: Williamar Rodrigues Silva

Study area description: The dataset was constructed from a forest inventory conducted in the ecotone forests of eastern Maracá Island, state of Roraima, northern Brazilian Amazonia: 3.360086 N to 3.405148 N / -61.44169 W to -61.48583 W (Fig. 1 ; figure was constructed using QGIS 2019 free software). A mosaic of ombrophilous and seasonal (deciduous and semideciduous) forest types predominates in this region. Maracá Island is part of the Maracá Ecological Station, a Brazilian protected area (~ 101,000 ha), located between two channels of the Rio Uraricoera (Maracá and Santa Rosa). The study region is a continuum of ecotone forests that have contact with the largest savanna area in northern Brazilian Amazonia (Barbosa et al. 2007, Barbosa and Campos 2011). Although it was inhabited until the mid-1970s (Proctor and Miller 1998) and there has

been an increase in anthropogenic pressures in adjacent regions (Couto-Santos et al. 2014), the conservation status of Maracá remains excellent and the area may be considered an important representative of mature forest tree species diversity for the region. Eastern Maracá soil classes are directly related to drainage and relief (51-99 m a.s.l.). These local constraints determine different forest types that occupy such seasonal flooding-free areas as moist lowlands and rocky slopes (Villacorta 2017). Well-drained soils are situated in areas of higher reliefs (Ultisols) or on slopes dominated by base-rich rocky soils. Soils occupying the lower-lying areas are poorly drained with a dominance of typical hydromorphic sandy soils (Nortcliff and Robison 1998). Regional climate is characterised as a transition between Aw/Am climate subtypes by the Köppen classification (Barbosa 1997). Average annual rainfall is ~ 1900 mm (1986-2010), with a rainy season between April and September (140-420 mm month⁻¹) and a dry season between October and March (40-130 mm month⁻¹) (Couto-Santos et al. 2014).

Funding: CNPq (Proc. CNPq 403591/2016-3) funded data collection.

Sampling methods

Description: The floristic survey was carried in March and December 2017.

Sampling description: The PPBio research grid was installed between 2005-2006 in eastern Maracá Island (see details in <https://ppbio.inpa.gov.br/sitios/maraca/infra>). The grid is formed by 12 x 5-km trails (six parallel trails in N-S direction and six in E-W direction, forming a 1-km resolution grid (Magnusson et al. 2005, Pezzini et al. 2012). A floristic survey was conducted within the grid in two periods (March and December 2017) to sample tree and palm species richness and composition in the ecotone forest mosaic. The floristic survey was carried out in 129 plots (10 m x 50 m each one; 6.45 ha in total), distributed on the six E-W trails (Fig. 1; Suppl. material 2). Each plot was systematically arranged along each trail with a minimum distance of 150 m between them to avoid pseudoreplication. This method was adopted to better cover the great environmental heterogeneity, characteristic of ecotone forests. The floristic survey included measurements of stem diameter (diameter tape - precision 0.1 cm), collecting botanical material and general description of the individuals/species, following standardised tree measurement methods established for PPBio grids and modules (Castilho et al. 2004). Palm height was estimated with Android technology using the Measure Height App (Où 2014), while trees height was estimated by regression using a Maracá-specific allometric model (Barbosa et al. 2019).

Analysis: We calculated abundance (number of stems) and richness (number of species) for all arboreal stems ≥ 10 cm in diameter (tree and palm) recorded in the floristic inventory. Frequency and dominance (absolute and relative) were also calculated to estimate the importance value indexes for family (FIV) and species (IVI). All stems were classified by diameter size and total height classes to analyse the horizontal-vertical structure. We tabulated all floristic composition and diameter data (n = 129 plots) using a multiple interacting spreadsheet programme (Excel Office 365) and applied the

vegetation analysis methods described in Kent and Coker (1994) to calculate frequency, dominance and importance value indexes (family and species).

Quality control: All observed individuals were collected and morphotyped in the field, then subsequently botanically identified. Vouchers representing most of the inventoried species were deposited in the herbarium of the INPA (Instituto Nacional de Pesquisa da Amazônia), MIRR (Museu Integrado de Roraima) and UFRR (Universidade Federal de Roraima). The species identification was performed by Antônio Tavares Mello by comparison with exsiccates in the INPA Herbarium and by R.O. Perdiz and R. I. Barbosa, taking into account comparison with exsiccates in the INPA, UFRR and MIRR herbariums and material available via the digitised collection of project REFLORA (Reflora - Virtual Herbarium, available at <http://reflora.jbrj.gov.br/reflora/herbarioVirtual/>). Species scientific names were checked and corrected by comparison with data from Brazil Flora Group (BFG 2015). Family-level delineations followed APG-IV (2016).

Step description: The floristic survey described here was done in two periods (March and December 2017).

Geographic coverage

Description: Data was collected in 129 plots across PPBio-Maracá research grid located on the eastern end of Maracá Ecological Station (see Fig. 1). All permanent plots are located on coordinates: 3.40515 N and -61.48583 W; 3.36009 N and -61.44169 W (Datum WGS 84).

Taxonomic coverage

Description: We observed a total of 3040 stems (tree = 2815; palm = 225) in the floristic inventory (129 plots = 6.45 ha), corresponding to 42 families, 119 genera and 140 identified species (Table 1). Seven families and 20 genera contained unidentified taxa (12.2% of total), all of which occurred due to a lack of appropriate taxonomic material (individuals dormant or without fertile material) to provide a definite determination. Sapotaceae (735 stems; 10 species), Leguminosae (409; 24) and Rubiaceae (289; 12) were the most important families with the highest family importance values (FIV = 42.4%). *P. gracilipes* Ducke (Leguminosae), *P. surinamensis* (Eyma) T.D.Penn. (Sapotaceae) and *E. guianensis* Eyma (Sapotaceae) were the species with the highest importance value index; ~ 25% in total (Table 2). Dominance ($\text{m}^2 \text{ha}^{-1}$) of these species corresponds to > 36% of the total observed in the forest inventory.

Palm stems (225) were observed only amongst the 10-40 cm diameter size classes, with the main distribution concentrated in the 20-30 cm class (62.2%) (Table 3). Most tree stems fell within the 10-20 cm class (56.7%), with the largest diameter class (> 50 cm) representing 5.9% of measured stems. The species with the largest structural parameters were *Laetia procera* – Salicaceae (stem diameter = 89.3 cm; total height = 33.5 m; stems

= 1), *Pochota fendleri* (Seem) W.S. Alverson & M.C. Duarte – Malvaceae (63.4 cm; 25.6 m; 9) and *Anacardium giganteum* Hancock ex Engl. - Anacardiaceae (63.3 cm; 30.2 m; 2), all of these being encountered at low abundance. Thirty-four species were represented by a single stem.

Usage licence

Usage licence: Creative Commons Public Domain Waiver (CC-Zero)

IP rights notes: These data can be freely used, provided their source is cited.

Data resources

Data package title: Tree species composition in ecotone forests of the eastern Maracá Island, Roraima, northern Brazilian Amazonia: preliminary data

Resource link: https://ipt.sibbr.gov.br/sibbr/resource?r=maraca_comp_floristic

Alternative identifiers: <https://www.gbif.org/dataset/01d28467-87a1-4d64-ba40-4e3e1cc9091b>

Number of data sets: 1

Data set name: Composição de espécies arbóreas em florestas de ecótono do leste da Ilha de Maracá, Roraima, norte da Amazônia brasileira: dados preliminares (Tree species composition in ecotone forests of the eastern Maracá Island, Roraima, northern Brazilian Amazonia: preliminary data).

Data format: Darwin Core Archive DwC-A

Description: Occurrence of tree and palm species identified during a floristic survey in 129 plots installed in Maracá Island, Roraima, northern Brazilian Amazonia. Dataset consist in the occurrence.txt file containing the DwC-Attributes.

| Column label | Column description |
|-----------------|--|
| basisOfRecord | The specific nature of the data record. |
| language | Language of the resource. |
| institutionCode | Institution that has custody of the object or information about its registration. |
| collectionCode | The name or acronym of the collection or dataset from which the record is derived. |
| occurrenceID | A identifier for the occurrence (unique). |
| catalogNumber | An identifier (preferably unique) for the record within the dataset or collection. |
| habitat | Description of the habitat in which the event occurred. |
| continent | The Continent of the occurrence. |

| | |
|-------------------------|--|
| country | The Country of the occurrence. |
| stateProvince | The State or Province of the occurrence. |
| county | The County of the occurrence. |
| locality | The location-specific description. |
| decimalLatitude | The geographical latitude in decimal degrees of the geographical centre of a location. |
| decimalLongitude | The geographical longitude in decimal degrees of the geographical centre of a location. |
| geodeticDatum | The ellipsoid, geodetic datum or spatial reference system (SRS) in which the geographical coordinates given in decimalLatitude and decimalLongitude are based. |
| kingdom | Full scientific name of the kingdom in which the taxon is classified. |
| family | Full scientific name of the family in which the taxon is classified. |
| genus | Full scientific name of the genus in which the taxon is classified. |
| specificEpithet | The name of the species-specific epithet. |
| intraspecificEpithet | The name of the terminal or lower-level infraspecific epithet of the scientific name. |
| scientificName | The full scientific name. It must be the name of lowest level taxonomic rank that was determined. |
| vernacularName | Common or vernacular name. |
| taxonRemarks | Comments or notes about the taxon or name. |
| identificationQualifier | A brief phrase or standard term ("cf.", "aff.") to express the determiner's doubts about identification. |

Additional information

Discussion

Our floristic composition results for ecotone forests on eastern Maracá Island complement previous investigations carried out at the macro (Milliken and Ratter 1998) and micro (Thompson et al. 1992, Nascimento et al. 1997, Nascimento and Proctor 1997) sampling scale at this location. Although our study was not conducted to differentiate the forest types comprising this ecotone region, the broad dispersion of 129 small plots (0.05 ha each) over a wide area (25 km²), aids understanding of the various forest types present in the area. This occurred since small plots were better suited to the environmental variability scale of this ecotone region, because they covered specific sampling areas of each forest type comprising the ecotonal mosaic. Use of plots smaller than those commonly used in tropical forest inventories (0.5-1.0 ha; e.g. Phillips et al. 2016) may be an alternative for floristic surveys or forest inventories in regions with high environmental variability. However, sampling, using small plots in Maracá, followed basic rules: (i) sampling design maintained the independence of each sampling unit, (ii) large number of samples (> 100) to adequately represent the environment and (iii) annual tree

censuses. All these recommendations are important to reduce the coefficient of variation between samples and to avoid temporal measurement problems associated with the descriptors of floristic composition, dynamics and forest structure (Keller et al. 2001, Wagner et al. 2010).

Our study recorded stem density (471 stems ha⁻¹) as the basal area (26.3 m² ha⁻¹) values compatible with those of Thompson et al. (1992) (range 340-476 ind ha⁻¹; 21.7-26.7 m² ha⁻¹) and Nascimento et al. (2014) (408-512 ind ha⁻¹; 26.0-32.5 m² ha⁻¹), both studies also being performed in eastern Maracá, but using smaller sampling areas. Overall, our study indicates that the main families described in previous inventories, such as Arecaceae, Burseraceae, Chrysobalanaceae, Leguminosae, Rubiaceae and Sapotaceae (Milliken and Ratter 1998, Nascimento et al. 1997), retained the proportional representation they had in surveys two decades earlier – indicating the site had little anthropic impact in the intervening time period. Additionally, the similarity of our larger sample to smaller-scale efforts of the past indicates that compositional units repeat, underscoring the near-fractal nature of the vegetation mosaic. The main families observed in Maracá are common throughout the Amazon, are always present in forest inventories and floristic surveys and almost always have the largest number of species, so that they are considered to be hyperdominant in the region (ter Steege et al. 2013). For example, Leguminosae and Sapotaceae (Condé and Tonini 2013) and Burseraceae, Chrysobalanaceae and Leguminosae (Alarcón and Peixoto 2007) were also families with high importance value indices in other Roraima forest types. We emphasise that the importance of each family in these surveys differs from Maracá, indicating that the ecotone forests of this region have their own composition and dynamics, so differing from mosaics or forest types observed elsewhere in the State of Roraima.

As with the families, most plant species, described in past inventories, are also present in our list, especially those with higher IVI (*P. gracilipes*, *P. surinamensis*, *E. guianensis*), besides *Lecythis corrugata* subsp. *rosea* (Lecythidaceae), *Attalea maripa* (Arecaceae) and *Licania discolor* (Chrysobalanaceae), all of which were strongly represented in previous surveys. The case of *L. discolor* Pilg. is the most interesting because the individuals attributed to this species in our work were largely attributed to *L. kunthiana* Hook.f. in previous surveys, a very similar taxonomic species, but were far less abundant at Maracá. The proportion and spatial distribution of species inventoried in Maracá ecotone forests are conditioned by a variety of environmental filters. For example, *P. gracilipes* is a deciduous species endemic to this area of the northern Amazonia that can form monodominant agglomerations (Nascimento and Proctor 1997). However, the monodominance of this species occurs only in seasonally flooded areas in bottom lands, with high Fe⁺² concentration in the soil (Nascimento et al. 2017). Such conditions are non-existent or rare in fertile and flood-free soils (Villacorta 2017). These distinct *P. gracilipes*-associated environmental characteristics reveal an ecotone region where forest types are intertwined with topographic, edaphic and hydrological constraints.

The species *P. gracilipes* plays an important ecological role (IVI = 10.4%) and it has been used to delimit forest types on Maracá. For example, Milliken and Ratter (1998) used this species to define forests monodominant with *Peltogyne* as “*Peltogyne gracilipes* forest”.

Similarly, Nascimento et al. (1997) used this species to divide the Maracá ecotone zone into three forest types: (i) PRF (*Peltogyne*-rich Forest) or forests monodominated with *P. gracilipes*, (ii) FWP (Forest without *Peltogyne*) or types where this species do not occur and (iii) PPF (*Peltogyne*-poor Forest), which are mixed types with low *P. gracilipes* abundance. The three types correspond analogously to the Deciduous Seasonal Forest (= PRF), Semideciduous Seasonal Forest (= PPF) and Open Ombrophilous Forest (= FWP) of the Brazilian Vegetation Classification System (Brazil-IBGE 2012), all with distinct hydro-edaphic and topographic characteristics (Carvalho et al. 2018). These forest types definitions have enhanced the understanding of variation in biomass/carbon stocks estimates for the Maracá Island (Nascimento et al. 2007, Nascimento et al. 2014).

Conclusion

The results of this study agree with data from previous investigations, indicating that the environmental heterogeneity of the ecotone forests of eastern Maracá Island influences floristic richness and structural distinctions, with *P. gracilipes* abundance acting as a descriptor variable for forest types. Consequently, the floristic survey conducted by this study is important because it expands our knowledge of forest structure and tree species composition in ecotone zones of the northern Brazilian Amazonia, improving our ability to predict changes in species composition and plant diversity when we take into account comparisons between previous forest inventories performed in Maracá. Finally, this study contributes to the local floristic knowledge, complements the herbarium collections with new collections, allows the development of similar studies and also enables the elaboration of management plans for the conservation of the local biota.

Acknowledgements

This study was supported by two projects: (i) Crescimento e mortalidade de árvores em florestas ecotonais de Roraima: efeito das condicionantes ambientais e da variabilidade climática (Proc. CNPq n. 403591/2016-3) and (ii) Ecologia e Manejo dos Recursos Naturais das Florestas e Savanas de Roraima (PPI-INPA 015/122). The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) provided a fellowship for R.I. Barbosa (CNPq 304204/2015-3) and a Ph.D. scholarship to R.O. Perdiz (CNPq 142243/2015-9). W.R. Silva and H.L.S. Farias were supported by postgraduate fellowships from PELD Program (CNPq/CAPES/FAPs/BC-Fundo Newton; Proc. n. 441575/2016-1), C.D.A. Villacorta by the Organization of American States (OAS), A.S. Oliveira by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and L.C.S. Carvalho by CAPES/PNPD (Programa Nacional de Pós-Doutorado). Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) provided permits for the study (SISBIO n. 52071). Two reviewers provided helpful comments.

Author contributions

WRS, CDAV, HLSF, ASO, LCSC and RIB collected the data; ACC provided the figures; WRS, ROP and RIB curated the data; WRS and RIB analysed the data and wrote the paper.

References

- Alarcón JGS, Peixoto AL (2007) Florística e fitossociologia de um trecho de um hectare de floresta de terra firme, em Caracará, Roraima, Brasil. *Boletim do Museu Paraense Emílio Goeldi (Ciências Naturais)* 2 (2): 33-60. URL: http://scielo.iec.gov.br/scielo.php?script=sci_arttext&pid=S1981-81142007000200004
- Alencar A, Brando P, Asner G, Putz F (2015) Landscape fragmentation, severe drought and the new Amazon forest fire regime. *Ecological Applications* 25 (6). <https://doi.org/10.1890/14-1528.1>
- Almeida CA, Coutinho AC, Esquerdo JCDM, Adami M, Venturieri A, Diniz CG, Dessay N, Durieux L, Gomes AR (2016) High spatial resolution land use and land cover mapping of the Brazilian Legal Amazon in 2008 using Landsat-5/TM and MODIS data. *Acta Amazonica* 46 (3): 291-302. <https://doi.org/10.1590/1809-4392201505504>
- APG-IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181: 1-20. <https://doi.org/10.1111/boj.12385>
- Aragão L, Anderson LO, Fonseca MG, Rosan TM, Vedovato LB, Wagner FH, Silva CVJ, Silva Junior CHL, Arai E, Aguiar AP, Barlow J, Berenguer E, Deeter MN, Domingues LG, Gatti L, Gloor M, Malhi Y, Marengo JA, Miller JB, Phillips OL, Saatchi S (2018) 21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions. *Nature communications* 9 (1). <https://doi.org/10.1038/s41467-017-02771-y>
- Barbosa RI, Barbosa RI, Ferreira EFG, Castellon EG (1997) Distribuição das chuvas em Roraima. *Homem, Ambiente e Ecologia no Estado de Roraima*. Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil, 613 pp. [ISBN 8521100086].
- Barbosa RI, Ferreira CA (2004) Biomassa acima do solo de um ecossistema de “campina” em Roraima, norte da Amazônia Brasileira. *Acta Amazonica* 34: 577-586. <https://doi.org/10.1590/S0044-59672004000400009>
- Barbosa RI, Campos C, Pinto F, Fearnside P (2007) The “Lavrados” of Roraima: Biodiversity and Conservation of Brazil’s Amazonian Savannas. *Functional Ecosystems and Communities* 1 (1): 29-41. URL: [http://www.globalsciencebooks.info/Online/GSBOnline/images/0706/FEC_1\(1\)/FEC_1\(1\)29-41o.pdf](http://www.globalsciencebooks.info/Online/GSBOnline/images/0706/FEC_1(1)/FEC_1(1)29-41o.pdf)
- Barbosa RI, Bacelar-Lima CG (2008) Notes on plant diversity and phytophysionomies in Roraima through herbarium INPA's database. *Amazônia: Ciência & Desenvolvimento* 4 (7): 131-154.
- Barbosa RI, Campos C (2011) Detection and geographical distribution of clearing areas in the savannas (‘lavrado’) of Roraima using Google Earth web tool. *Journal of Geography and Regional Planning* 4 (3): 122-136. URL: <https://www.semanticscholar.org/paper/>

[Detection-and-geographical-distribution-of-clearing-Barbosa-Campos/a4ac528277a5712a6dd4b5316b3054ddb1097599](https://doi.org/10.1016/j.foreco.2017.04.026)

- Barbosa RI, Castilho CV, Perdiz RO, Damasco G, Rodrigues R, Fearnside PM (2017) Decomposition rates of coarse woody debris in undisturbed Amazonian seasonally flooded and unflooded forests in the Rio Negro-Rio Branco Basin in Roraima, Brazil. *Forest Ecology and Management* 397: 1-9. <https://doi.org/10.1016/j.foreco.2017.04.026>
- Barbosa RI, Ramírez-Narváez PN, Fearnside PM, Villacorta CDA, Carvalho LCS (2019) Allometric models to estimate tree height in northern Amazonian ecotone forests. *Acta Amazonica* 49 (2): 81-90. <https://doi.org/10.1590/1809-4392201801642>
- Barlow J, Lennox GD, Ferreira J, Berenguer E, Lees AC, Mac Nally R, Thomson JR, Ferraz SF, Louzada J, Oliveira VH, Parry L, Solar RR, Vieira IC, Aragao LE, Begotti RA, Braga RF, Cardoso TM, de Oliveira RCJ, Souza Jr CM, Moura NG, Nunes SS, Siqueira JV, Pardini R, Silveira JM, Vaz-de-Mello FZ, Veiga RC, Venturieri A, Gardner TA (2016) Anthropogenic disturbance in tropical forests can double biodiversity loss from deforestation. *Nature* 535 (7610): 144-7. <https://doi.org/10.1038/nature18326>
- Barni PE, Pereira VB, Manzi AO, Barbosa RI (2015) Deforestation and forest fires in Roraima and their relationship with phytoclimatic regions in the Northern Brazilian Amazon. *Environmental Management* 55 (5): 1124-1138. <https://doi.org/10.1007/s00267-015-0447-7>
- BFG (2015) Growing knowledge: an overview of Seed Plant diversity in Brazil. *Rodriguésia* 66 (4): 1085-1113. <https://doi.org/10.1590/2175-7860201566411>
- Brazil-IBGE (2012) Manual técnico da vegetação brasileira: sistema fitogeográfico, inventário das formações florestais e campestres, técnicas e manejo de coleções botânicas, procedimentos para mapeamentos. Instituto Brasileiro de Geografia e Estatística (IBGE), Rio de Janeiro, Brazil. 2nd. ed. 275 p. URL: ftp://geofpt.ibge.gov.br/documentos/recursos_naturais/manuais_tecnicos/manual_tecnico_vegetacao_brasileira.pdf
- Cardoso D, Sarkinen T, Alexander S, Amorim AM, Bittrich V, Celis M, Daly DC, Fiaschi P, Funk VA, Giacomini LL, Goldenberg R, Heiden G, Iganci J, Kelloff CL, Knapp S, Cavalcante de Lima H, Machado AFP, Dos Santos RM, Mello-Silva R, Michelangeli FA, Mitchell J, Moonlight P, de Moraes PLR, Mori SA, Nunes TS, Pennington TD, Pirani JR, Prance GT, de Queiroz LP, Rapini A, Riina R, Rincon CAV, Roque N, Shimizu G, Sobral M, Stehmann JR, Stevens WD, Taylor CM, Trovo M, van den Berg C, van der Werff H, Viana PL, Zartman CE, Forzza RC (2017) Amazon plant diversity revealed by a taxonomically verified species list. *Proceedings of the National Academy of Sciences of the United States of America* 114 (40): 10695-10700. <https://doi.org/10.1073/pnas.1706756114>
- Carvalho LCS, Fearnside PM, Nascimento MT, Barbosa RI (2018) Amazon soil charcoal: Pyrogenic carbon stock depends of ignition source distance and forest type in Roraima, Brazil. *Global Change Biology* 24 (9): 4122-4130. <https://doi.org/10.1111/gcb.14277>
- Castilho CV, Schiatti J, Freitas MA, Araújo MC, Coelho F, Magnusson W, Costa F (2004) Manual para medição e marcação de árvores em grades e módulos RAPELD do PPBio. Repositório de Dados do PPBio. PPBio Amazônia Ocidental, Manaus, AM, Brazil. URL: https://ppbio.inpa.gov.br/sites/default/files/Protocolo_estrutura_vegetacao_2014_0.pdf
- Condé TM, Tonini H (2013) Fitossociologia de uma floresta ombrófila densa na Amazônia Setentrional, Roraima, Brasil. *Acta Amazonica* 43 (3): 247-260. <https://doi.org/10.1590/S0044-59672013000300002>

- Couto-Santos F, Luizão F, Carneiro-Filho A (2014) The influence of the conservation status and changes in the rainfall regime on forest-savanna mosaic dynamics in Northern Brazilian Amazonia. *Acta Amazonica* 44 (2): 197-206. <https://doi.org/10.1590/S0044-59672014000200005>
- Damasco G, Vicentini A, Castilho C, Pimentel T, Nascimento HM (2013) Disentangling the role of edaphic variability, flooding regime and topography of Amazonian white-sand vegetation. *Journal of Vegetation Science* 24 (2). <https://doi.org/10.1111/j.1654-1103.2012.01464.x>
- Esquivel-Muelbert A, Baker TR, Dexter KG, Lewis SL, Brienen RJW, Feldpausch TR, Lloyd J, Monteagudo-Mendoza A, Arroyo L, Alvarez-Davila E, Higuchi N, Marimon BS, Marimon-Junior BH, Silveira M, Vilanova E, Gloor E, Malhi Y, Chave J, Barlow J, Bonal D, Davila Cardozo N, Erwin T, Fauset S, Herault B, Laurance S, Poorter L, Qie L, Stahl C, Sullivan MJ, Ter Steege H, Vos VA, Zuidema PA, Almeida E, Almeida de Oliveira E, Andrade A, Vieira SA, Aragao L, Araujo-Murakami A, Arets E, Aymard CG, Baraloto C, Camargo PB, Barroso JG, Bongers F, Boot R, Camargo JL, Castro W, Chama Moscoso V, Comiskey J, Cornejo Valverde F, Lola da Costa AC, Del Aguila Pasquel J, Di Fiore A, Fernanda Duque L, Elias F, Engel J, Flores Llampazo G, Galbraith D, Herrera Fernandez R, Honorio Coronado E, Hubau W, Jimenez-Rojas E, Lima AJN, Umetsu RK, Laurance W, Lopez-Gonzalez G, Lovejoy T, Aurelio Melo Cruz O, Morandi PS, Neill D, Nunez Vargas P, Pallqui Camacho NC, Parada Gutierrez A, Pardo G, Peacock J, Pena-Claros M, Penuela-Mora MC, Petronelli P, Pickavance GC, Pitman N, Prieto A, Quesada C, Ramirez-Angulo H, Rejou-Mechain M, Restrepo Correa Z, Roopsind A, Rudas A, Salomao R, Silva N, Silva Espejo J, Singh J, Stropp J, Terborgh J, Thomas R, Toledo M, Torres-Lezama A, Valenzuela Gamarra L, van de Meer PJ, van der Heijden G, van der Hout P, Vasquez Martinez R, Vela C, Vieira ICG, Phillips OL (2019) Compositional response of Amazon forests to climate change. *Global Change Biology* 25 (1): 39-56. <https://doi.org/10.1111/gcb.14413>
- Fearnside PM (2006) Desmatamento na Amazônia: dinâmica, impactos e controle. *Acta Amazonica* 36 (3): 395-400. <https://doi.org/10.1590/S0044-59672006000300018>
- Hufkens K, Scheunders P, Ceulemans R (2009) Ecotones in vegetation ecology: methodologies and definitions revisited. *Ecological Research* 24: 977-986. <https://doi.org/10.1007/s11284-009-0584-7>
- Jaramillo MMA (2015) Estrutura, biomassa arbórea e composição florística de ilhas de mata da savana de Roraima, Norte da Amazônia Brasileira. MSc. Thesis, UFRR - PRONAT, Boa Vista, Roraima, Brazil. URL: http://bdt.d.ibict.br/vufind/Record/UFRR_7f0b9955cdd438aabdb34036bc47394c
- Keller M, Palace M, Hurr G (2001) Biomass estimation in the Tapajós National Forest, Brazil: examination of sampling and allometric uncertainties. *Forest Ecology and Management* 154: 371-382. [https://doi.org/10.1016/S0378-1127\(01\)00509-6](https://doi.org/10.1016/S0378-1127(01)00509-6)
- Kent M, Coker P (1994) *Vegetation description and analysis: a practical approach*. Wiley & Sons, Chichester. [ISBN 0471948101]
- Lovejoy T, Nobre C (2018) Amazon tipping point. *Science Advances* 4 (2): eaat2340. <https://doi.org/10.1126/sciadv.aat2340>
- Magnusson W, Lima A, Luizão R, Luizão F, Costa FC, Castilho CV, Kinupp VF (2005) RAPELD: A modification of the Gentry Method for biodiversity surveys in long-term ecological research sites. *Biota Neotropica* 5 (2): 19-24. <https://doi.org/10.1590/S1676-06032005000300002>

- MapBiomias (2019) Amazonia - Dados de la cobertura del suelo. Rede MapBiomias URL: <http://plataforma.amazonia.mapbiomas.org/map>
- Marques E, Marimon-Junior BH, Marimon B, Matricardi ET, Mews H, Colli G (2019) Redefining the Cerrado–Amazonia transition: implications for conservation. *Biodiversity and Conservation* <https://doi.org/10.1007/s10531-019-01720-z>
- Milliken W, Ratter JA, Milliken W, Ratter JA (1998) The vegetation of the Ilha de Maracá. Maracá: the biodiversity and environment of an Amazonian rainforest. Wiley, Chichester, UK, 508 pp. [ISBN 0471979171].
- Nascimento M, Proctor J (1997) Population dynamics of five tree species in a monodominant Peltophyne forest and two other forest types on Maracá Island, Roraima, Brazil. *Forest Ecology and Management* 94 (1-3): 115-128. [https://doi.org/10.1016/S0378-1127\(96\)03968-0](https://doi.org/10.1016/S0378-1127(96)03968-0)
- Nascimento MT, Proctor J, Villela DM (1997) Forest structure, floristic composition and soils of an Amazonian monodominant forest on Maracá Island, Roraima, Brazil. *Edinburgh Journal of Botany* 54: 1-38. <https://doi.org/10.1017/S096042860000384X>
- Nascimento MT, Barbosa RI, Villela DM, Proctor J (2007) Above-ground biomass changes over an 11-year period in an Amazon monodominant forest and two other lowland forests. *Plant Ecology* 192 (2): 181-191. <https://doi.org/10.1007/s11258-007-9303-z>
- Nascimento MT, Carvalho LC, Barbosa RI, Villela DM (2014) Variation in floristic composition, demography and above-ground biomass over a 20-year period in an Amazonian monodominant forest. *Plant Ecology & Diversity* 7 (102): 293-303. <https://doi.org/10.1080/17550874.2013.772673>
- Nascimento MT, Barbosa RI, Dexter K, Castilho CV, Silva Carvalho LC, Villela DM (2017) Is the Peltophyne gracilipes monodominant forest characterised by distinct soils? *Acta Oecologica* 85: 104-107. <https://doi.org/10.1016/j.actao.2017.10.001>
- Nelson B, Oliveira A, Capobianco JPR, Veríssimo A, Moreira A, Sawyer D, Santos I, Pinto LP (2001) Área Botânica. Biodiversidade na Amazônia: avaliação e ações prioritárias para a conservação, uso sustentável e repartição de benefícios. Instituto Socioambiental, São Paulo, 21 pp. [ISBN 8585994134].
- Nogueira EM, Yanai AM, Fonseca FO, Fearnside PM (2015) Carbon stock loss from deforestation through 2013 in Brazilian Amazonia. *Global Change Biology* 21: 1271-1292. <https://doi.org/10.1111/gcb.12798>
- Nortcliff S, Robison D, Milliken W, Ratter J (1998) The soils of the Ilha de Maracá. Maracá: The Biodiversity & Environment of an Amazonian Rainforest. Wiley, Chichester, UK, 508 pp. [ISBN 0471979171].
- Oliveira RLC, Farias HLS, Perdiz RO, Scudeller VV, Barbosa RI (2017) Structure and tree species composition in different habitats of savanna used by indigenous people in the Northern Brazilian Amazon. *Biodiversity Data Journal* (5). <https://doi.org/10.3897/BDJ.5.e20044>
- Oü D (2014) Measure Height Application (Version 1.4). Google Play URL: https://play.google.com/store/apps/details?id=ee.deskis.android.height&hl=pt_BR
- Pezzini F, Melo PHA, Oliveira DMS, Amorim RX, Figueiredo FOG, Drucker DP, Rodrigues FRO, Zuquim G, Emilio T, Costa FRC, Magnusson WE, Sampaio AF, Lima AP, Garcia ARM, Manzatto AG, Nogueira A, Costa CP, Barbosa CEA, Bernardes C, Castilho CV, Cunha CN, Freitas CG, Cavalcante CO, Brandão DO, Rodrigues DJ, Santos ECPR, Baccaro FB, Ishida FY, Carvalho FA, Moulatlet GM, Guillaumet JB, Pinto JLPV, Schiatti J, Vale JD, Belger L, Verdade LM, Pansonato MP, Nascimento MT, Santos MCV, Cunha

- MS, Arruda R, Barbosa RI, Romero RL, Pansini S, Pimentel TP (2012) The Brazilian Program for Biodiversity Research (PPBio) Information System. *Biodiversity & Ecology* 4: 265-274. <https://doi.org/10.7809/b-e.00083>
- Phillips O, Baker T, Feldpausch T, Brienen R (2016) RAINFOR Field Manual for plot establishment and remeasurement. University of Leeds, Leeds, UK. URL: http://www.rainfor.org/upload/ManualsEnglish/RAINFOR_field_manual_version_2016.pdf
 - Proctor J, Miller R, Milliken W, Ratter J (1998) Human occupation on the Ilha de Maracá: preliminary notes. Maracá: The biodiversity & environment of an Amazonian Rainforest. Wiley, Chichester, UK, 508 pp. [ISBN 0471979171].
 - QGIS (2019) QGIS Geographic Information System (QGIS 3.8 Zanzibar). Open Source Geospatial Foundation Project URL: <http://qgis.osgeo.org>
 - Salati E, Vose P (1984) Amazon Basin: a system in equilibrium. *Science* 255 (4658): 129-138. <https://doi.org/10.1126/science.225.4658.129>
 - Santos NMC, Vale Jr. JF, Barbosa RI (2013) Florística e estrutura arbórea de ilhas de mata em áreas de savana do norte da Amazônia brasileira. *Boletim do Museu Paraense Emílio Goeldi (Ciências Naturais)* 8 (2): 205-221. URL: https://www.researchgate.net/publication/257382499_Floristic_and_tree_structure_of_forest_islands_in_savanna_areas_of_Northern_Brazilian_A
 - Sette-Silva EL (1993) Inventário preliminar das espécies arbóreas das florestas dos arredores de Boa Vista (Roraima) - uma abordagem fitossociológica. MSc. Thesis, INPA/FUA, Manaus, AM. URL: https://koha.inpa.gov.br/cgi-bin/koha/opac-detail.pl?biblionumber=5208&shelfbrowse_itemnumber=6313
 - Silva WR, Villacorta CD, Carvalho LC, Farias HL, Perdiz RO, Barbosa RI (2019) Tree species composition in ecotone forests of the eastern Maracá Island, Roraima, northern Brazilian Amazonia: preliminary data (version 1.3). Sistema de Informação sobre a Biodiversidade Brasileira - SiBBR: Occurrence dataset. GBIF <https://doi.org/10.15468/xa5lrb>
 - ter Steege H, Pitman NC, Sabatier D, Baraloto C, Salomao RP, Guevara JE, Phillips OL, Castilho CV, Magnusson WE, Molino JF, Monteagudo A, Nunez Vargas P, Montero JC, Feldpausch TR, Coronado EN, Killeen TJ, Mostacedo B, Vasquez R, Assis RL, Terborgh J, Wittmann F, Andrade A, Laurance WF, Laurance SG, Marimon BS, Marimon BHJ, Guimaraes Vieira IC, Amaral IL, Brienen R, Castellanos H, Cardenas Lopez D, Duivenvoorden JF, Mogollon HF, Matos FD, Davila N, Garcia-Villacorta R, Stevenson Diaz PR, Costa F, Emilio T, Levis C, Schiatti J, Souza P, Alonso A, Dallmeier F, Montoya AJ, Fernandez Piedade MT, Araujo-Murakami A, Arroyo L, Gribel R, Fine PV, Peres CA, Toledo M, Aymard CG, Baker TR, Ceron C, Engel J, Henkel TW, Maas P, Petronelli P, Stropp J, Zartman CE, Daly D, Neill D, Silveira M, Paredes MR, Chave J, Lima Filho Dde A, Jorgensen PM, Fuentes A, Schongart J, Cornejo Valverde F, Di Fiore A, Jimenez EM, Penuela Mora MC, Phillips JF, Rivas G, van Andel TR, von Hildebrand P, Hoffman B, Zent EL, Malhi Y, Prieto A, Rudas A, Ruschell AR, Silva N, Vos V, Zent S, Oliveira AA, Schutz AC, Gonzales T, Trindade Nascimento M, Ramirez-Angulo H, Sierra R, Tirado M, Umana Medina MN, van der Heijden G, Vela CI, Vilanova Torre E, Vriesendorp C, Wang O, Young KR, Baider C, Balslev H, Ferreira C, Mesones I, Torres-Lezama A, Urrego Giraldo LE, Zagt R, Alexiades MN, Hernandez L, Huamantupa-Chuquimaco I, Milliken W, Palacios Cuenca W, Pauletto D, Valderrama Sandoval E, Valenzuela Gamarra L, Dexter

- KG, Feeley K, Lopez-Gonzalez G, Silman MR (2013) Hyperdominance in the Amazonian tree flora. *Science* 342 (6156). <https://doi.org/10.1126/science.1243092>
- ter Steege H, Mota de Oliveira S, Pitman NCA, Sabatier D, Antonelli A, Guevara Andino JE, Aymard GA, Salomao RP (2019) Towards a dynamic list of Amazonian tree species. *Scientific reports* 9 (1). <https://doi.org/10.1038/s41598-019-40101-y>
 - Thompson J, Proctor J, Viana V, Milliken W, Ratter JA, Scott DA (1992) Ecological studies on a lowland evergreen rain forest on Maraca Island, Roraima, Brazil. I. Physical environment, forest structure and leaf chemistry. *Journal of Ecology* 80 (4): 689-703. <https://doi.org/10.2307/2260860>
 - Villacorta CDA (2017) Condicionantes edáficas para a monodominância do gênero *Peltogyne* (Fabaceae) na Ilha de Maracá, Roraima, norte da Amazônia brasileira. MSc Thesis, UFRR - PRONAT, Boa Vista, Roraima, 46 pp.
 - Wagner F, Rutishauser E, Blanc L, Herault B (2010) Effects of plot size and census interval on descriptors of forest structure and dynamics. *Biotropica* 42 (6): 664-671. <https://doi.org/10.1111/j.1744-7429.2010.00644.x>
 - WWF (2019) Amazon places: facts. World Wildlife Fund. Washington, DC URL: <https://www.worldwildlife.org/places/amazon>




Figure 1. Study area with emphasis on the PPBio research grid (Maracá), Roraima, northern Brazilian Amazonia.

Table 1.

List of the most important arboreal families observed for eastern Maracá Island, where: Abundance = number of stems \geq 10 cm in diameter; BA = total basal area (m^2); ReAb = relative abundance (%); ReFq = relative frequency (%); ReDo = relative dominance (%); FIV = family importance value (%) representing the mean between ReAb, ReFq and ReDo. Raw data can be accessed on the GBIF website (Silva et al. 2019).

| Family | Genus | Species | Abundance | AB m^2 | ReAb (%) | ReFq (%) | ReDo (%) | FIV (%) |
|------------------|-------|---------|-----------|----------|----------|----------|----------|---------|
| Sapotaceae | 4 | 10 | 735 | 4597.95 | 24.2 | 10.4 | 27.1 | 20.5 |
| Leguminosae | 18 | 24 | 409 | 3615.21 | 13.5 | 7.8 | 21.3 | 14.2 |
| Rubiaceae | 9 | 11 | 289 | 860.76 | 9.5 | 8.5 | 5.1 | 7.7 |
| Arecaceae | 6 | 6 | 225 | 1228.04 | 7.4 | 7.4 | 7.2 | 7.4 |
| Lecythidaceae | 5 | 6 | 239 | 880.84 | 7.9 | 7.2 | 5.2 | 6.7 |
| Burseraceae | 2 | 8 | 160 | 705.41 | 5.3 | 5.7 | 4.2 | 5.0 |
| Chrysobalanaceae | 7 | 7 | 116 | 831.58 | 3.8 | 5.1 | 4.9 | 4.6 |
| Apocynaceae | 2 | 4 | 114 | 402.16 | 3.8 | 4.9 | 2.4 | 3.7 |
| Moraceae | 7 | 8 | 91 | 344.19 | 3.0 | 4.5 | 2.0 | 3.2 |
| Annonaceae | 3 | 5 | 86 | 214.72 | 2.8 | 3.0 | 1.3 | 2.4 |
| other 33 | 56 | 52 | 576 | 3286.97 | 18.9 | 35.5 | 19.4 | 24.6 |
| Total | 119 | 141 | 3040 | 16967.83 | 100 | 100 | 100 | 100 |

Table 2.

List of the most important arboreal species (stem diameter ≥ 10 cm) observed on eastern Maracá Island, northern Brazilian Amazonia, where: Density = stem density (absolute and relative), Dominance = BA in m^2 (absolute and relative). Frequency = number of plots where the species was observed (relative), IVI = importance value index (mean between %Density, %Dominance, %Frequency). The raw data can be accessed on the GBIF website (Silva et al. 2019).

| Species | Family | Abundance (ind) | Density | | Dominance | | Frequency | IVI |
|---|------------------|--------------------|----------------------|------|---------------------------------|-------|-----------|-------|
| | | | ind ha ⁻¹ | % | m ² ha ⁻¹ | % | % | |
| <i>Peltogyne gracilipes</i> Ducke | Leguminosae | 299 | 46.4 | 9.84 | 4.7 | 17.92 | 3.39 | 10.38 |
| <i>Pradosia surinamensis</i> (Eyma) T.D.Penn. | Sapotaceae | 170 | 26.4 | 5.59 | 3.0 | 11.22 | 4.96 | 7.26 |
| <i>Ecclinusa guianensis</i> Eyma | Sapotaceae | 276 | 42.8 | 9.08 | 1.9 | 7.09 | 4.58 | 6.92 |
| <i>Attalea maripa</i> (Aubl.) Mart. | Arecaceae | 156 | 24.2 | 5.13 | 1.5 | 5.86 | 4.33 | 5.11 |
| <i>Lecythis corrugata</i> subsp. <i>rosea</i> (Spruce ex O.Berg) S.A.Mori | Lecythidaceae | 178 | 27.6 | 5.86 | 0.9 | 3.47 | 4.02 | 4.45 |
| <i>Alseis longifolia</i> Ducke | Rubiaceae | 160 | 24.8 | 5.26 | 0.9 | 3.56 | 3.58 | 4.13 |
| <i>Pouteria surumuensis</i> Baehni | Sapotaceae | 113 | 17.5 | 3.72 | 0.8 | 2.99 | 3.26 | 3.32 |
| <i>Pouteria hispida</i> Eyma | Sapotaceae | 85 | 13.2 | 2.80 | 1.0 | 3.73 | 3.01 | 3.18 |
| <i>Protium stevensonii</i> (Standl.) Daly | Burseraceae | 98 | 15.2 | 3.22 | 0.8 | 2.95 | 2.82 | 3.00 |
| <i>Licania discolor</i> Pilg. | Chrysobalanaceae | 72 | 11.2 | 2.37 | 1.0 | 3.80 | 2.70 | 2.96 |
| <i>Himatanthus articulatus</i> (Vahl) Woodson | Apocynaceae | 100 | 15.5 | 3.29 | 0.5 | 1.93 | 3.14 | 2.78 |
| <i>Pouteria venosa</i> (Mart.) Baehni | Sapotaceae | 62 | 9.6 | 2.04 | 0.4 | 1.38 | 2.20 | 1.87 |
| <i>Simaba orinocensis</i> Kunth | Simaroubaceae | 30 | 4.7 | 0.99 | 0.8 | 3.06 | 1.51 | 1.85 |
| <i>Pseudolmedia laevigata</i> Trécul | Moraceae | 48 | 7.4 | 1.58 | 0.2 | 0.71 | 2.13 | 1.47 |
| <i>Duroia eriopila</i> L.f. | Rubiaceae | 47 | 7.3 | 1.55 | 0.1 | 0.46 | 1.88 | 1.30 |
| <i>Astrocarium aculeatum</i> G.Mey | Arecaceae | 37 | 5.7 | 1.22 | 0.2 | 0.76 | 1.63 | 1.20 |
| <i>Pochota fendleri</i> (Seem) W.S. Alverson & M.C. Duarte | Malvaceae | 9 | 1.4 | 0.30 | 0.7 | 2.59 | 0.38 | 1.09 |
| <i>Gutteria schomburgkiana</i> Mart. | Annonaceae | 30 | 4.7 | 0.99 | 0.2 | 0.74 | 1.51 | 1.08 |
| <i>Duguetia lepdota</i> (Miq.) Pulle | Annonaceae | 47 | 7.3 | 1.55 | 0.1 | 0.43 | 1.19 | 1.06 |

| | | | | | | | | |
|--------------------------------|-----------|------|-------|-------|------|-------|-------|-------|
| <i>Quiina rhytidopus</i> Tull. | Ochnaceae | 26 | 4.0 | 0.86 | 0.1 | 0.21 | 1.51 | 0.86 |
| Other 120 | - | 997 | 154.6 | 32.80 | 6.6 | 25.15 | 46.26 | 34.74 |
| Total | 43 | 3040 | 471.3 | 100 | 26.3 | 100 | 100 | 100 |

Table 3.

Mean and standard deviation of structural parameters (stem diameter and total height) observed in ecotone forests of eastern Maracá Island, northern Brazilian Amazonia.

| Classes (cm) | Palm | | | Tree | | |
|--------------|------------|---------------|---------------|------------|---------------|---------------|
| | Height (m) | Diameter (cm) | Abundance (n) | Height (m) | Diameter (cm) | Abundance (n) |
| 10-20 | 15.0±4.0 | 16.8±2.6 | 32 | 13.5±1.9 | 13.8±2.8 | 1598 |
| 20-30 | 15.4±5.3 | 25.4±2.8 | 138 | 19.2±1.4 | 23.8±3.0 | 591 |
| 30-40 | 19.7±5.7 | 32.1±1.7 | 55 | 23.3±1.0 | 34.0±2.9 | 297 |
| 40-50 | | | | 26.2±0.8 | 43.9±3.0 | 159 |
| > 50 | | | | 29.8±2.0 | 62.5±13.1 | 170 |
| Total | 16.4±5.6 | 25.8±5.3 | 225 | 17.4±5.4 | 22.7±14.1 | 2815 |

Supplementary materials

Suppl. material 1: The main studies in Roraima ecotone areas involving forest inventories and floristic surveys.

Authors: Williamar Rodrigues Silva and Reinaldo Imbrozio Barbosa

Data type: List of studies

[Download file](#) (18.75 kb)

Suppl. material 2: Geographic coordinates (degree - lat/long; DATUM WGS84) and altitude (m a.s.l.) of sampling units (plotID) in the eastern Maracá Island.

Authors: Williamar Rodrigues Silva, Carlos Darwin Angulo Villacorta and Reinaldo Imbrozio Barbosa

Data type: geographic coordinates (lat/long)

[Download file](#) (4.45 kb)