

Intertidal insects associated with halophytic *Suaeda* (Amaranthaceae) in Japan: a case study in Saga, northern Kyushu

Akihito Kita^{‡,§}, Ayman Khamis Elsayed[‡], Makoto Tokuda^{‡,|}

[‡] Faculty of Agriculture, Saga University, Saga, Japan

[§] Saga Prefectural Space and Science Museum, Takeo, Japan

[|] The United Graduate School of Agricultural Sciences, Kagoshima University, Kagoshima, Japan

Corresponding author: Makoto Tokuda (tokudam@cc.saga-u.ac.jp)

Academic editor: David Bilton

Abstract

In contrast to a great diversity in insects in terrestrial and freshwater ecosystems, few known species have adapted to inhabit marine environments. In this study, we surveyed insects associated with halophytic plants of *Suaeda* (Amaranthaceae) distributed in intertidal zones, in northern Kyushu, Japan. On four Japanese native species of *Suaeda*, we found insects belonging to five orders and 18 species. Amongst them, the genus *Clanoneurum* (Diptera: Ephydriidae) and *Coleophora deviella* (Lepidoptera: Coleophoridae) were newly reported from Japan; and *Orthotylus* (*Melanotrichus*) *parvulus* (Hemiptera: Miridae) was newly recorded from Kyushu. The seasonal occurrence of several insects on *Suaeda* is reported.

Keywords

halophyte, insect fauna, intertidal zone, *Suaeda*

Introduction

Insects have achieved a great diversity in terrestrial and freshwater ecosystems, but they seldom inhabit marine environments (e.g. Cheng 1976, Ward 1992). Only some insect taxa are known to adapt to intertidal zones and very few in neritic and oceanic environments (Ward 1992, Ikawa et al. 2012, Kaiser et al. 2016).

In intertidal zones, Coleoptera are the most dominant group in terms of the number of genera reported and followed by Diptera, Collembola and Hemiptera (Ward 1992). These insects exhibit various adaptations to seawater, such as plastron respiration under seawater and behavioural avoidance of submergence (Hinton 1976, Ward 1992).

Adaptive mechanisms of intertidal insects to marine habitats are interesting study subjects to understand their survival strategies in harsh environments for many insects.

Some herbivorous insects are associated with halophytes growing in the intertidal zones (Foster and Treherne 1975, Dorchin 2001, Dorchin and Freidberg 2008, Cho et al. 2015, Elsayed et al. 2015). As herbivory critically affects some plant community structures in intertidal zones (Rand 2002, Pennings et al. 2007), further intensive studies of herbivores are important to understand their role in determining plant community structures in intertidal zones.

The genus *Suaeda* (Amaranthaceae) consists of approximately 100 species and most of them grow in coastal areas and tidal wetlands (Yonekura 2017). Several studies have reported insects on *Suaeda*, such as a psyllid (Hemiptera: Sternorrhyncha) associated with *Suaeda japonica* Makino in Korea (Cho et al. 2015); gall midges (Diptera: Cecidomyiidae) in the Holarctic Region and South America (Kieffer 1909, Kieffer and Jörgensen 1910, Felt 1918, Mamaev and Mirumian 1990, Dorchin 2001, Dorchin and Freidberg 2008, Skuhravá et al. 2014, Elsayed et al. 2015) and their hymenopteran parasitoids (Gates et al. 2018), a leaf beetle in North America (von Groll et al. 2022) and lepidopterans in various localities worldwide (Paik et al. 2013, Adamski et al. 2018, Karisch et al. 2020, Budashkin and Bidzilya 2021). However, information on insect fauna associated with *Suaeda* is still fragmental and comprehensive studies have never been conducted as far as we know.

The area facing the Ariake Sea in Saga Prefecture has the largest mudflats in Japan, owing to the largest tidal range in the country, which provides a suitable habitat for halophytes including *Suaeda* (Henmi et al. 2017, Tokuda 2019). In the present study, we periodically surveyed the insect fauna associated with four *Suaeda* species in Saga, Japan and examined the seasonal occurrence of several dominant insect species to reveal their ecological aspects.

Materials and Methods

Study plants

Four native species of *Suaeda* are known to be distributed in Japan (Yonekura 2017): *Suaeda glauca* (Bunge) Bunge distributed in Honshu and Kyushu, Japan as well as in eastern Siberia, Ussuri, Mongol, China and Korean Peninsula; *Suaeda maritima* (L.) Dumort. subsp. *asiatica* H. Hara distributed widely in East Asia including Honshu and Kyushu, Japan; *Suaeda malacosperma* H. Hara distributed in western Honshu and Kyushu, Japan, as well as in Korean Peninsula; and *S. japonica* distributed in northern Kyushu, Japan and Korean Peninsula. They are all annual plants (Yonekura 2017). We investigated insects associated with all four native *Suaeda* species in this study.

Periodical investigations

Field investigations were conducted at the following six census sites in Saga Prefecture: Benga, Imari City (*S. glauca*); Iida, Kashima City (*S. maritima asiatica*); Muta, Tara Town (*S. maritima asiatica*); Inuido, Saga City (*S. malacosperma*); Edo, Saga City (*S. japonica*); and Higashiyoka, Saga City (*S. japonica*) (Fig. 1). Amongst them, Benga faces the Genkai Sea, i.e. areas between the Japan Sea and the East China Sea and the other five sites are located along the coastline of the Ariake Sea. The investigations were conducted at two-week intervals from April to November 2015 and monthly from December 2015 to December 2016.

At each census site, insects on *Suaeda* plants were collected by an approximately five-minute sweeping on each census date (qualitative survey). In 2016, three quadrats (1 m × 1 m) were set in *Suaeda* communities and insects inhabiting there were visually investigated in each quadrat (quantitative survey) and collected by net sweeping. Collected insects were kept either as dried specimens or in 99% ethanol for future DNA analyses.

Inventory data of insects on each *Suaeda* species were based on both qualitative and quantitative surveys and the seasonal occurrence data of major species were on the qualitative survey.

Results and Discussion

Insect fauna

Throughout the field surveys, we found 18 insect species belonging to five orders from *Suaeda* (Table 1).

In Orthoptera, *Gampsocleis buergeri* de Haan (Tettigoniidae) and *Euparattix insularis* Bey-Bienko (Tetrigidae) were found respectively on *S. glauca* (in 2015) and *S. malacosperma*. Amongst them, *G. buergeri* is distributed in western Honshu and northern Kyushu, Japan and *E. insularis* is in Honshu, Shikoku, Kyushu, the Izu Islands, the Ogasawara Islands and Korean Peninsula (Ichikawa et al. 2006). Both species are polyphagous and not specialists of *Suaeda* plants.

In Hemiptera, *Aphis* sp. (Aphididae) was found on all four *Suaeda* species surveyed (Fig. 2A). This species is probably undescribed and a specialist of *Suaeda* plants (Y. Matsumoto, personal communication). *Orthotylus (Melanotrichus) parvulus* (Miridae) was found on *S. maritima asiatica* and *S. japonica* (Fig. 2E). This species is associated with *S. maritima asiatica* and *Salicornia europaea* (Amaranthaceae) and is distributed in the Palearctic Region, but in Japan, it was only known from Tsushima Island and Hyogo Prefecture, Honshu (Yasunaga et al. 2001, Yasunaga et al. 2016, Shishido and Yasunaga

2016). We newly report this species from Kyushu, as mentioned earlier and *S. japonica* is a new host record for this species.

In Coleoptera, *Baris scolopacea* Germar (Curculionidae) was found on *S. japonica* and *S. glauca* in 2015 (Fig. 2B). This weevil is distributed in Honshu, Shikoku, Kyushu, Tsushima, Amami-Oshima and Kuroshima in Japan, as well as in Korea and China (Yoshihara 2016). In western parts of Japan, *B. scolopacea* is associated with Amaranthaceae and induces galls on *Achyranthes* sp., *Chenopodium* spp. and *Dysphania anthelmintica* (L.) Mosyakin et Clemants (= *Ambrina anthelmintica* (L.) Spach) (Yoshihara 2016). In this study, we did not find galls on *Suaeda* plants. As we found stem galls on *Atriplex patens* (Litv.) Iljin (Amaranthaceae) growing close to *Suaeda*, the weevil is possibly responsible for them. Three widely distributed species of Coccinellidae in Japan were found on *S. japonica*, namely *Harmonia axyridis* (Pallas), *Coccinella septempunctata* L. and *Propylea japonica* (Thunberg). As these coccinellids are predatory species (Sakamoto 2018), they probably visit *Suaeda* plants to feed on aphids and other insects. Furthermore, *Medythia nigrobilineata* (Motschulsky) (Chrysomelidae) was found on *S. malacosperma* and *Eobia cinereipennis* (Motschulsky) (Oedemeridae) was on *S. glauca* (in 2015). *Medythia nigrobilineata* is distributed in Hokkaido, Honshu, Shikoku, Kyushu, Sado, Tsushima, the Goto Islands, the Korean Peninsula, China and eastern Siberia and is known to feed on Fabaceae (Morimoto and Hayashi 1986). As we found wild fabaceous plants near the census site of *S. malacosperma*, the species may accidentally visit *S. malacosperma* from them. *Eobia cinereipennis* is distributed in Hokkaido, Honshu, Shikoku, Kyushu, Izu Islands, Amami Islands, Ryukyus and Korea. Although this species is a flower-visiting species inhabiting seashores (Morimoto and Hayashi 1986), it was collected in June which is not the flowering season of *S. glauca*. For this reason, this species probably visited *S. glauca* by chance while visiting the flowers of other plants growing around *S. glauca*.

In Diptera, *Clanoneurum* sp. (Ephidridae) was found on *S. japonica* and *S. maritima asiatica* (Fig. 2C) and two species of Syrphidae, namely *Epistrophe balteata* de Geer and *Metasyrphus nitens* Zetterstedt, were found on *S. japonica*. The genus *Clanoneurum* is newly reported from Japan in this paper. This genus contains four species worldwide (GBIF Secretariat 2019). Further taxonomic studies are needed to confirm whether the species found in this study is undescribed. As the syrphids are found in October, the flowering season of *S. japonica*, they might visit the flowers of the plant. The pollination ecology of these *Suaeda* plants is an important study subject in the future.

In Lepidoptera, *Coleophora deviella* Zeller (Coleophoridae) was found on *S. japonica* and *S. maritima asiatica* (Fig. 2D). This species is known in the western Palaearctic Region (from Spain to southern Russia) and is associated with several species of Amaranthaceae including *S. maritima* (Anikin 1988, Ellis 2020). This is the first record of *C. deviella* from the eastern Palaearctic Region. Two species of polyphagous Noctuidae were found in this study; *Sarcopolia illoba* (Butler) was on *S. glauca* and *Spodoptera litura* (Fabricius) was on *S. glauca* and *S. japonica*. Both species are polyphagous species (Yoshimatsu et al. 2011). In addition, *Spoladea recurvalis* (Fabricius) (Crambidae) was found on *S. japonica*, *S. glauca* and *S. maritima asiatica* (Fig. 2F). This

species is polyphagous and known to be associated with Amaranthaceae and some other plants (Yoshimatsu and Miyata 2011). *Spoladea recurvalis* was previously collected over the East China Sea and, in Kyushu, large numbers of individuals have been collected in autumn along the coast, suggesting its long-distance migration habit (Yoshimatsu and Miyata 2011). In addition, larvae of an unidentified species of Geometridae were found on *S. japonica* and *S. maritima asiatica*.

Seasonal occurrence

Amongst insects found on *Suaeda* species, the seasonal occurrence of *Aphis* sp., *Clanoneurum* sp. and *C. deviella* was investigated in the quantitative survey in 2016 by counting individuals of *Aphis* sp., leaves mined by *Clanoneurum* sp. and larval cases formed by *C. deviella* on plants, respectively.

In Edo and Higashiyoka, the number of *Aphis* sp. individuals peaked in September (Fig. 3 A). In addition, another small peak was detected in July in Edo. As no individuals were found in May and from October to December, this species may exhibit host alternation, but hosts, other than *Suaeda*, are not yet known at present.

Mines produced by *Clanoneurum* sp. were found from July to October in Iida and Muta (Fig. 3B). At both census sites, two peaks were found in summer (July or August) and autumn (October). The first peak was larger than the second in Iida and vice versa in Muta. These results suggest that the species is bivoltine.

The number of larval cases formed by *C. deviella* gradually increased from summer to autumn and peaked in September or October in all localities (Fig. 3C), suggesting the univoltine life cycle of this species.

Conclusions

In this study, we recognised 18 insect species on *Suaeda* plants and investigated seasonal occurrence of several herbivorous species. Amongst the insects, *Aphis* sp. is probably an undescribed species; *O. parvulus* was newly recorded from Kyushu; the genus *Clanoneurum* and *C. deviella* were newly reported from Japan. As mentioned in the Introduction, faunistic studies of insects associated with *Suaeda* is limited worldwide, but our findings indicate diverse fauna of insects, especially halophyte-associated herbivores adapting to intertidal zones.

Acknowledgements

We thank S. Iwamura for his valuable information on *Suaeda* plants. Our thanks are extended to Y. Matsumoto, Y. Nakatani, Y. Sakamaki, M. Suwa and H. Yoshitake for identifying *Aphis* sp., *O. (M.) parvulus*, *Clanoneurum* sp., *B. scolopacea* and *C. deviella*, respectively and for giving us useful information about these insects. We are grateful to S.

Adachi-Fukunaga, A. Tetsuka, M. Okuzono and other members of the Laboratory of Systems Ecology, Faculty of Agriculture, Saga University for their kind help in field surveys.

Ethics and security

- This work is original research carried out by the authors.
- All authors agree with the contents of the manuscript and its submission to the journal.
- No part of the research has been published in any form elsewhere, unless it is fully acknowledged in the manuscript.
- The manuscript is not being considered for publication elsewhere while it is being considered for publication in this journal.
- Any research in the paper not carried out by the authors is fully acknowledged in the manuscript.
- All appropriate ethics and other approvals were obtained for the research.

Author contributions

Conceptualisation: [Makoto TOKUDA]; Methodology: [Akihito KITA], [Makoto TOKUDA]; Data Analysis: [Makoto TOKUDA]; Investigation: [Akihito KITA], [Ayman Khamis ELSAYED], [Makoto TOKUDA]; Data Curation: [Akihito KITA], [Makoto TOKUDA]; Writing-Original Draft Preparation: [Akihito KITA], [Makoto TOKUDA]; Writing-Review and Editing: [Akihito KITA], [Ayman Khamis ELSAYED], [Makoto TOKUDA]; Visualisation: [Akihito KITA], [Ayman Khamis ELSAYED], [Makoto TOKUDA]; Supervision: [Makoto TOKUDA]; Project Administration: [Makoto TOKUDA].

All authors have read and agreed to the published version of the manuscript.

Conflicts of interest

The authors declare no conflict of interest.

References

- Adamski D, Kula RR, Gates MW, Torr ns J, Fidalgo P, Buffington ML (2018) Immature stages of *Scrobipalpus patagonica* Povoln y, 1977 (Lepidoptera: Gelechiidae: Gnorimoschemini), a gall inquiline of *Suaeda divaricata* MOQ. (Amaranthaceae) in Argentina with a summary of its parasitoids. Proceedings of the Entomological Society of Washington 120 (4): 659-669. <https://doi.org/10.4289/0013-8797.120.4.659>
- Anikin VV (1988) The casebearers of the Volga-Ural inter-river region (Lepidoptera, Coleophoridae). Entomofauna 19: 33-44.

- Budashkin YI, Bidzilya OV (2021) Review of the genus *Aporiptura* Falkovitsh, 1972 (Lepidoptera, Coleophoridae) with description of a new species from Crimea. Entomological Review 101 (4): 564-574. <https://doi.org/10.1134/s0013873821040060>
- Cheng L (1976) Marine Insects. North-Holland Publishing Company, Amsterdam – Oxford & American Elsevier Publishing Company, New York, 581 pp.
- Cho G, Burckhardt D, Lee S (2015) *Rhodochlanis suaedicola* sp. nov. (Hemiptera: Sternorrhyncha: Psylloidea: Aphalaridae) associated with *Suaeda japonica* (Amaranthaceae) from Korea. Zootaxa 4028: 388-396. <https://doi.org/10.11646/zootaxa.4028.3.4>
- Dorchin N (2001) Gall midges (Diptera: Cecidomyiidae) infesting *Suaeda monoica* (Chenopodiaceae) in Israel. Proceedings of the Entomological Society of Washington 103 (3): 561-581.
- Dorchin N, Freidberg A (2008) The Chenopodiaceae-feeding gall midges (Diptera: Cecidomyiidae) of the Na'aman salt marsh, Israel. Zootaxa 1937: 1-22. <https://doi.org/10.11646/zootaxa.1937.1.1>
- Ellis WN (2020) Plant Parasites of Europe, leafminers, galls and fungi. <https://bladmineerders.nl/>. Accessed on: 2021-12-04.
- Elsayed AK, Skuhrová M, Karam HH, Elminshawy A, Al-Eryan MA (2015) New records and new species of gall midges (Diptera: Cecidomyiidae) developing on Chenopodiaceae in Egypt. Zootaxa 3904: 105-115. <https://doi.org/10.11646/zootaxa.3904.1.6>
- Felt EP (1918) New gall midges (Dipt.). Journal of Economic Entomology 11: 380-384. <https://doi.org/10.1093/jee/11.4.380>
- Foster WA, Treherne JE (1975) The distribution of an intertidal aphid *Pemphigus trehernei* Foster, on marine saltmarshes. Oecologia 21: 141-155. <https://doi.org/10.1007/BF00345557>
- Gates MW, Torrens J, Fidalgo P, Kula RR, Buffington ML, Adamski D (2018) The Gall Associates of *Asphondylia* poss. *suaedicola* Kieffer & Jörgensen (Diptera: Cecidomyiidae) on *Suaeda divaricata* Moq. (Amaranthaceae) in the Semiarid Argentina and Summary of Parasitic Hymenoptera Associated with *Suaeda* Worldwide. Neotropical Entomology 47 (5): 598-609. <https://doi.org/10.1007/s13744-017-0557-4>
- GBIF Secretariat (2019) *Clanoneurum* Becker, 1903. GBIF Backbone Taxonomy. <https://doi.org/10.15468/39omei>. Accessed on: 2021-12-04.
- Henmi Y, Fuchimoto D, Kasahara Y, Shimanaga M (2017) Community structures of halophytic plants, gastropods and brachyurans in salt marshes in Ariake and Yatsushiro seas of Japan. Plankton and Benthos Research 12 (4): 224-237. <https://doi.org/10.3800/pbr.12.224>
- Hinton HE (1976) Respiratory adaptations of marine insects. In: Cheng L (Ed.) Marine Insects. North-Holland Publishing Company, Amsterdam – Oxford & American Elsevier Publishing Company, New York, 43–78 pp.
- Ichikawa A, Ito F, Kano Y, Kawai M, Tominaga O, Murai T (2006) Orthoptera of the Japanese archipelago in Color. Hokkaido University Press, Sapporo, 687 pp. [In Japanese].
- Ikawa T, Nozoe Y, Yamashita N, Nishimura N, Ohnoki S, Yusa K, Hoshizaki S, Komaba M, Kawakubo A (2012) Life histories of two endangered sea skaters *Halobates matsumurai* Esaki and *Asclepios shiranui* (Esaki) (Hemiptera: Gerridae: Halobatinae). Psyche 2012: 1-7. <https://doi.org/10.1155/2012/261071>

- Kaiser T, Poehn B, Szkiba D, Preussner M, Sedlazeck F, Zrim A, Neumann T, Nguyen L, Betancourt A, Hummel T, Vogel H, Dorner S, Heyd F, von Haeseler A, Tessmar-Raible K (2016) The genomic basis of circadian and circalunar timing adaptations in a midge. *Nature* 540: 69-73. <https://doi.org/10.1038/nature20151>
- Karisch T, Fowler L, Stevens N, Dutton AJ (2020) Three new *Opogona* species with wing reduction from St Helena Island (Lepidoptera, Tineidae). *Metamorphosis* 31: 33-39.
- Kieffer JJ (1909) Contributions à la connaissance des insectes gallicoles. *Bulletin de la Société d'Histoire Naturelle de Metz* 3 (2): 1-35.
- Kieffer JJ, Jörgensen P (1910) Gallen und Gallentiere aus Argentinien. *Centralblatt für Bakteriologie, Parasitenkunde und Infektionskrankheiten* 27 (2): 362-444.
- Mamaev BM, Mirumian LS (1990) Description of the gall midge *Arafavilla terteriani* gen. et. sp. n. (Diptera, Cecidomyiidae), inducing galls on *Suaeda altissima* (L.) Pall. in Armenia. *Doklady Akademii Nauk Armyanskoï SSR* 90: 139-141.
- Morimoto K, Hayashi N (1986) The coleoptera of Japan in color. Vol. 1. Hoikusha Publishers Co., Ltd, Osaka, 450 pp. [In Japanese].
- Paik CH, Lee GH, Choi MY, Noh TH, Shim HK (2013) Overwintering site and occurrence dynamics of *Scrobipalpa salinella* (Zeller) (Lepidoptera: Gelechiida). *Korean Journal of Applied Entomology* 52 (2): 71-74. <https://doi.org/10.5656/KSAE.2012.12.0.078>
- Pennings S, Zimmer M, Dias N, Sprung M, Davé N, Ho C, Kunza A, McFarlin C, Mews M, Pfaunder A, Salgado C (2007) Latitudinal variation in plant-herbivore interactions in European salt marshes. *Oikos* 116 (4): 543-549. <https://doi.org/10.1111/j.0030-1299.2007.15591.x>
- Rand TA (2002) Variation in insect herbivory across a salt marsh tidal gradient influences plant survival and distribution. *Oecologia* 132: 549-558. <https://doi.org/10.1007/s00442-002-0989-2>
- Sakamoto Y (2018) *The Handbook of Lady-birds*. Bun-ichi Co., Ltd., Tokyo, 88 pp. [In Japanese].
- Shishido T, Yasunaga T (2016) New record of *Orthotylus (Melanotrichus) choii* Josifov from Japan, with a taxonomic review of *Melanotrichus* Reuter (Heteroptera, Miridae, Orthotylinae). *Rostria* 60: 21-29. [In Japanese with English summary].
- Skuhrová M, Skuhrový V, Elsayed AK (2014) Gall midges (Diptera: Cecidomyiidae) of Egypt-annotated list and zoogeographical analysis. *Acta Societatis Zoologicae Bohemicae* 78: 241-268.
- Tokuda M (2019) Natural environments and endangered organisms in the Rokkaku River including tidal flat areas of the Ariake Sea. *Kasen (River)* 879: 65-68. [In Japanese].
- von Groll E, Moura L, Carvalho G (2022) Revision, morphometry and cladistics of *Erynephala* (Coleoptera: Chrysomelidae: Galerucinae: Galerucini). *Zoologischer Anzeiger* 296: 1-32. <https://doi.org/10.1016/j.jcz.2021.11.001>
- Ward JV (1992) *Aquatic insect ecology 1. Biology and habitat*. Jone Wiley & Sons, Inc., New York, 438 pp.
- Yasunaga T, Takai M, Kawasawa T (2001) *A field guide to Japanese bugs II – Terrestrial heteropterans*. Zenkoku Noson Kyoiku Kyokai Publishing Co., Ltd., Tokyo, 350 pp. [In Japanese]. [ISBN ISBN978-4-88137-089-6]
- Yasunaga T, Yamada K, Ishikawa T (2016) Family Miridae. In: *The Editorial Committee of Catalogue of the Insects of Japan (Ed.) Catalogue of the Insects of Japan volume 4 Paraneoptera*. The Entomological Society of Japan, Fukuoka, 376-421 pp. [In Japanese].

- Yonekura K (2017) *Amaranthaceae*. In: Ohashi H, Kadota H, Murata J, Yonekura K, Kihara H (Eds) *Wild flowers of Japan, revised edition vol. 4. Malvaceae ~ Apocynaceae*. Heibonsha Ltd., Tokyo, 128-142 pp. [In Japanese].
- Yoshihara K (2016) *The insects of Japan Vol. 6. Coleoptera, Curculionidae, Barididnae*. Touka Shobo, Fukuoka, 171 pp.
- Yoshimatsu S, Miyata A (2011) *Crambidae*. In: Komai F, Yoshiyasu Y, Nasu Y, Saito T (Eds) *A Guide to the lepidoptera of Japan*. Tokai University Press, Hadano, 741-770 pp. [In Japanese].
- Yoshimatsu S, Miyata A, Owada M, Yoshiyasu Y, Nasu Y (2011) *Noctuidae*. In: Komai F, Yoshiyasu Y, Nasu Y, Saito T (Eds) *A Guide to the lepidoptera of Japan*. Tokai University Press, Hadano, 885-930 pp. [In Japanese].




Figure 1.

Map of census sites. *Suaeda* plants were surveyed at the following sites: *S. glauca* in Benga, *S. japonica* in Edo and Higatayoka, *S. malacosperma* in Inuido and *S. maritima asiatica* in Iida and Muta.




Figure 2.

Insects found on *Suaeda* plants. **A** A nymph of *Aphis* sp. (Hemiptera: Aphididae) found on *S. maritima asiatica* in Iida; **B** An adult of *Baris scolopacea* (Coleoptera: Curculionidae) found on *S. glauca* in Benga; **C** An adult of *Clanoneurum* sp. (Diptera: Ephidridae) found on *S. japonica* in Edo; **D** An adult of *Coleophora deviella* (Lepidoptera: Coleophoridae) found on *S. japonica* in Edo; **E** An adult of *Orthotylus (Melanotrichus) parvulus* (Hemiptera: Miridae) found on *S. maritima asiatica* in Muta; and **F** An adult of *Spoladea recurvalis* (Lepidoptera: Crambidae) found on *S. glauca* in Benga.




Figure 3.

Seasonal changes in densities of major insects on *Suaeda*. **A** Seasonal changes in the number of *Aphis* sp. individuals (per plant) on *Suaeda* plants in 2016; **B** Seasonal changes in the number of mines (per plant) produced by *Clanoneurum* sp. larvae on *Suaeda* plants in 2016 and **C** Seasonal changes in the number of larval cases (per plant) formed by *Coleophora deviella* on *Suaeda* plants in 2016.

Table 1.

Insects found on *Suaeda* species and their feeding habit. Abbreviations of *Suaeda* plants are as follows: SG, *S. glauca*; SJ, *S. japonica*; SMRT, *S. maritima asiatica*; and SMRC, *S. maracosperma*. P: present on the plant.

Order	Family	Species	Feeding habit	SG	SJ	SMRT	SMRC
Orthoptera	Tettigoniidae	<i>Gampsocleis buergeri</i>	Omnivore	P			
	Tetrigidae	<i>Euparatettix insularis</i>	Herbivore				P
Hemiptera	Aphididae	<i>Aphis</i> sp.	Herbivore		P	P	P
	Miridae	<i>Orthotylus (Melanotrichus) parvulus</i>	Herbivore	P	P	P	
Coleoptera	Curculionidae	<i>Baris scolopacea</i>	Herbivore	P	P		
	Coccinellidae	<i>Coccinella septempunctata</i>	Predator		P		
		<i>Harmonia axyridis</i>	Predator		P		
		<i>Propylea japonica</i>	Predator		P		
	Chrysomelidae	<i>Medythia nigrobilineata</i>	Herbivore				P
Oedemeridae	<i>Eobia cinereipennis</i>	Herbivore	P				
Diptera	Ephydriidae	<i>Clanoneurum</i> sp.	Herbivore		P		P
	Syrphidae	<i>Metasyrphus nitens</i>	Herbivore		P		
		<i>Epistrophe balteata</i>	Herbivore		P		
Lepidoptera	Coleophoridae	<i>Coleophora deviella</i>	Herbivore		P	P	
	Noctuidae	<i>Sarcopolia illoba</i>	Herbivore	P			
	Noctuidae	<i>Spodoptera litura</i>	Herbivore	P	P		
	Crambidae	<i>Spoladea recurvalis</i>	Herbivore	P		P	
	Geometridae	gen. sp. (unidentified)	Herbivore		P		P
Number of insect species found				7	12	4	5