

***Geastrum suae* sp. nov. (Gastraceae, Basidiomycota) a new species from Yunnan Province, China**

Zheng-Quan Zhang[‡], Chao-Hai Li^{‡§}, Lin Li^{‡¶||}, Hong-Wei Shen^{‡¶||}, Jun He[#], Xi-Jun Su[‡], Zong-Long Luo[‡]

[‡] College of Agriculture and Biological Science, Dali University, Dali, China

[§] College of Pharmacy, Dali University, Dali, China

[|] Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai, Thailand

[¶] School of Science, Mae Fah Luang University, Chiang Rai, Thailand

[#] Biotechnology and Germplasm Resources Institute, Yunnan Academy of Agricultural Sciences, Kunming, China

Corresponding author: Zong-Long Luo (luozonglongfungi@163.com)

Academic editor: Alfredo Vizzini

Abstract

Background

Geastrum is the largest genus of Gastraceae and is widely distributed all over the world. Four specimens which belong to *Geastrum* were collected during our scientific expedition to Cangshan Mountain, Yunnan, China. Based on morphological characteristics and phylogenetic analysis, a new species was introduced.

New information

Geastrum suae is characterised by its large basidiomata (height 35–70 mm, diameter 18–37 mm) with long stipe (height 10–45 mm), smooth pink exoperidium and sessile globose endoperidial body. Phylogenetic analysis has been carried out, based on the internal transcribed spacer (ITS) and large subunit ribosomal ribonucleic acid (nrLSU) sequence data. The illustration and description for the new taxa are provided.

Keywords

Gastraceae, ITS, nrLSU, taxonomy, phylogeny

Introduction

Gastrum Pers. is the largest genus of Geastraceae and was established by Persoon (1794). *Gastrum* is commonly known as the earthstars with worldwide distribution and the most species-diverse in the family Geastraceae. Up to now, there are 140 valid species in this genus (Wijayawardene et al. 2022, Zhou et al. 2022, Cabral et al. 2022, Wang and Bau 2023). *Gastrum* clearly differs from *Myriostoma* by a single endodermal stoma (Sousa et al. 2014). Due to the non-splitting ectoderm and the poorly-developed endoperidium being different from *Gastrum*, researchers thought that *Radiigera* is one of the genera closely related to *Gastrum* (Sunhede 1989, de Toledo and Castellano 1996). Later, some studies have found that specimens in *Radiigera* are nested in *Gastrum* (Hosaka et al. 2006, Hosaka and Castellano 2008, da Silva et al. 2013), but the relationship between these two genera has not been studied in depth until Jeppson et al. (2013) classified the species of *Radiigera* into the genus *Gastrum*. Species of this genus are distributed globally, especially in temperate and tropical regions, such as Brazil-Amazon and Europe (de León 1968, da Silva et al. 2013, Jeppson et al. 2013, Zamora et al. 2014, Cabral et al. 2014a, Cabral et al. 2014b, Sousa et al. 2015, Crous et al. 2016, Cabral et al. 2017, Crous et al. 2017, Sousa et al. 2017, Crous et al. 2018a, Crous et al. 2018b, Assis et al. 2019, Finy et al. 2021, Rodrigues et al. 2021). However, the taxonomic relationship under the genus was chaotic (Zamora et al. 2013) until Zamora et al. (2014) divided it into 14 Sections using polygenic analysis, viz. Sect. *Campestria*, *Corollina*, *Elegantly*, *Exareolata*, *Fimbriata*, *Fornicata*, *Gastrum*, *Hariotia*, *Hieronymia*, *Myceliostroma*, *Papillata*, *Pseudoilmbata*, *Schmidelia* and *Trichaster*.

In China, the early systematic report of *Gastrum* can be found in "Fungi in China" (Deng 1963) and "The Confluence of Chinese Fungi" (Dai 1979). Zhou et al. (2007) detailed descriptions of 16 species of *Gastrum* in China in "Flora Fungorum Sinicorum-Geastraceae and Nidulariaceae". Later, three new records and nine new species were reported (Han and Bau 2016, Zhou et al. 2022, Wang and Bau 2023).

Four specimens which belong to *Gastrum* were collected during our scientific expedition to Cangshan Mountain, Yunnan, China. Morphological and phylogenetic analysis revealed that these specimens are the same species and are different from other species in *Gastrum*. Therefore, we introduced it as a new species and provided the detailed description and illustration.

Materials and methods

Morphological description

Macro-morphological descriptions were based on fresh specimens, which were photographed in the field with notes and laboratory supplemental measurements. The colour is compared with the standard colours in the colorhexa website (<https://colorhexa.com>)

www.colorhexa.com). Micro-morphological data were obtained from the fresh specimens and observed by using a light microscope, following Accioly et al. (2019). Sections were studied at magnification of up to 1000 \times using a NiKon eclipse Ni microscope and phase contrast illumination and scanning electron microscope (SEM) analysis was done under a Shimadzu SSX-550. Preparation of the material examined under SEM followed da Silva et al. (2011). Microscopic features and measurements were made from slide preparations stained with 5% potassium hydroxide (KOH). Basidiospore features, hyphal system, colour, sizes and shapes were recorded and photographed. Measurements were made using the Image Framework v.0.9.7 to represent variation in the size of basidiospores, 5% of measurements were excluded from each end of the range and extreme values are given in parentheses.

The abbreviation for spore measurements (n/m/p) denote “n” spores measured from “m” basidiocarps of “p” specimens. Basidiospore dimensions (and “Q” values) are given as (a) b–av–c (d), where “a” represents the minimum, “d’ the largest, “av” the average “b” and “c” covers a minimum of 90% of the values. “Q” is the length/width ratio of a spore inside view and “Qm” for the average of all spores \pm standard deviation. Voucher specimens are deposited in the Herbarium of Cryptogams, Kunming Institute of Botany Academia Sinica (KUN-HKAS).

DNA extraction, PCR amplification and sequencing

The DNA extractions were performed from a small piece of the dried basidioma by using Trelief™ Plant Genomic DNA Kit from Tsingke Biotechnology Co., Ltd (Beijing, China). Two DNA regions were amplified: the internal transcribed spacer nuclear ribosomal DNA (ITS), nuclear ribosomal large subunit (nrLSU) with the primer pairs ITS1F/ITS4 and LR0R/LR5, respectively (Table 1).

PCR reactions (25 μ l) contained mixture: 12.5 μ l 2X SanTaq PCR Master Mix (including MgCl₂, dNTP, Taq DNA Polymerase, PCR buffer, loading etc.), 1 μ l each of primer, 2 μ l DNA solution and 9.5 μ l sterilised distilled H₂O. The PCR cycling for ITS and nrLSU was as follows: initial denaturation at 94°C for 5 min, followed by 35 cycles at 94°C for 30 sec, 53°C for 30 sec and 72°C for 50 sec and a final extension of 72°C for 10 min. The PCR products were visualised via UV light after electrophoresis on 1% agarose gels stained with ethidium bromide. Successful PCR products were sent to Sangon Biotech Limited Company (Shanghai, China), using forward PCR primers. When sequences have heterozygous INDELS or ambiguous sites, samples were sequenced bidirectionally to make contigs of the amplified regions or verify the ambiguous sites. Raw DNA sequences were assembled and edited in Sequencher 4.1.4 and the assembled DNA sequences were deposited in GenBank (Table 2).

Sequence alignment

Sequence data of two partial loci, internal transcribed spacer region (ITS) and the large subunit ribosomal RNA gene (nrLSU) were analysed. All the sequences, except those

which were obtained from this study, were selected from GenBank for phylogenetic analyses (Table 2). Sequences were aligned using the online version of MAFFT v.7 (<http://mafft.cbrc.jp/alignment/server/>) (Katoh and Standley 2013) and adjusted using BioEdit v.7.0.9 (Hall 1999) by hand to allow maximum alignment and minimise gaps. Ambiguous regions were excluded from the analyses and gaps were treated as missing data. AliView 1.19-beta was used to convert the alignment fasta file to Phylip and Nexus format for phylogenetic analysis. Phylogenetic analyses were obtained from Maximum Likelihood (ML) and Bayesian Inference (BI).

Molecular phylogenetic analyses

The Maximum Likelihood (ML) and Bayesian Inference (BI) methods were used to analyse the combined dataset of ITS and nrLSU sequences. ML analysis was conducted with RAxML-HPC2 on the CIPRES Science Gateway (Miller et al. 2010), involving 100 ML searches; all model parameters were estimated by the programme. The ML bootstrap values (ML-BS) were obtained with 1000 rapid bootstrapping replicates.

Bayesian analysis was performed with MrBayes v.3.2 (Ronquist et al. 2012), with the best-fit model of sequence evolution estimated with MrModelTest 2.3 (Nylander et al. 2008) to evaluate posterior probabilities (PP) (Rannala and Yang 1996, Zhaxybayeva and Gogarten 2002) by Markov Chain Monte Carlo (MCMC) sampling. Six simultaneous Markov chains were run for 100,000,000 generations, trees were sampled every 500th generation and 200,000 trees were obtained. The first 50,000 trees, representing the burn-in phase of the analyses, were discarded, while the remaining 150,000 trees were used for calculating posterior probabilities in the majority rule consensus tree (the critical value for the topological convergence diagnostic is 0.01).

The phylogenetic tree was visualised with FigTree version 1.4.0 (Rambaut 2012) and made in Adobe Illustrator CS5 (Adobe Systems Inc., USA). Sequences derived in this study were deposited in GenBank (<http://www.ncbi.nlm.nih.gov>).

Taxon treatment

Geastrum suae Z.Q. Zhang C.H. Li & Z.L. Luo, sp. nov.

- MycoBank [MB845193](#)

Materials

Holotype:

- a. scientificName: *Geastrum suae*; kingdom: Fungi; phylum: Basidiomycota; class: Agaricomycetes; order: Geastrales; family: Geastraceae; genus: *Geastrum*; verbatimElevation: 2160 m; locationRemarks: China, Yunnan Province, Dali City, Cangshan Mountain; verbatimLatitude: 25°43'36.97"N; verbatimLongitude: 100°07'16.46"E; year: 2020; month: September; day: 4; habitat: Terrestrial; fieldNotes: grows in groups on the ground in mixed coniferous and broad-leaved forests, with thick

humus; recordNumber: SJ582; recordedBy: Zheng-Quan Zhang; type: KUN-HKAS 123795; occurrenceID: 6D676216-9572-5A8D-A7C3-4C445C671395

Paratype:

- a. scientificName: *Geastrum suae*; kingdom: Fungi; phylum: Basidiomycota; class: Agaricomycetes; order: Geastrales; family: Geastraceae; genus: *Geastrum*; verbatimElevation: 2221 m; locationRemarks: China, Yunnan Province, Dali City, Cangshan Mountain; verbatimLatitude: 25°40'16.38"N; verbatimLongitude: 100°09'08.42"E; year: 2020; month: October; day: 14; habitat: Terrestrial; fieldNotes: grows in groups on the ground in mixed coniferous and broad-leaved forests, with thick humus; recordNumber: MB015; recordedBy: Chao-Hai Li; type: KUN-HKAS 123796; occurrenceID: F58D4D0C-33BB-541B-A92B-7AF436F12F49

Other materials:

- a. scientificName: *Geastrum suae*; kingdom: Fungi; phylum: Basidiomycota; class: Agaricomycetes; order: Geastrales; family: Geastraceae; genus: *Geastrum*; verbatimElevation: 2208 m; locationRemarks: China, Yunnan Province, Dali City, Cangshan Mountain; verbatimLatitude: 25°40'28"N; verbatimLongitude: 100°08'59"E; year: 2021; month: September; day: 3; habitat: Terrestrial; fieldNotes: grows in groups on the ground in mixed coniferous and broad-leaved forests, with thick humus; recordNumber: SJ2501; recordedBy: K. Wang; type: KUN-HKAS 123793; occurrenceID: 9213A508-19C3-5A9C-8E34-8DCDD9D4C170
- b. scientificName: *Geastrum suae*; kingdom: Fungi; phylum: Basidiomycota; class: Agaricomycetes; order: Geastrales; family: Geastraceae; genus: *Geastrum*; verbatimElevation: 2350 m; locationRemarks: China, Yunnan Province, Yangbi County, Cangshan Mountain; verbatimLatitude: 25°41'59"N; verbatimLongitude: 100°02'00"; year: 2021; month: October; day: 1; habitat: Terrestrial; fieldNotes: grows in groups on the ground in mixed coniferous and broad-leaved forests, with thick humus; recordNumber: SJ2500; recordedBy: G. H. Yang; type: KUN-HKAS 123794; occurrenceID: CE65C858-1CD7-53B3-B545-F1750391FB8E

Description

Unexpanded basidiomata 13–28 mm, cylindrical to ellipsoidal, very light grey (#fdfdfd) to very pale red (#ffe6e6) with a slight protrusion, rough. Expanded basidiomata height 35–70 mm, diameter 18–37 mm, deep saccate, **Exoperidium** splitting into 6, arched, not hygrometric, prosthecae length 23–35 mm, diameter 5–13 mm, exoperidium attached to the rhizomorphs. Rhizomorphs with 0.1–5.4 µm hyphae, fibrous and transparent, white (#ffffff). **Mycelial layer** 49.5–59.0 µm, consisting of transparent hyphae (1.0–3.5 µm) with thin walls and no septum, curved. **Fibrous layer** 6.5–16.5 µm, transparent, curved, thick-walled hyphae (1.1–5.0 µm) smooth, transparent to cream (#ffffdd), pure red (#e60000) to dark red (#9a0000) when stained with Congo red. **Pseudoparenchymatous layer** 2.5–19.3 × 2.7–30.4 µm, irregular shape, mycelium is transparent when fresh, pure orange (#ffa500) to moderate pink (#cc6691) when stained with Congo red, the thickness of the pseudoparenchyma layer is about 1.0–1.3 mm, very soft pink (#d98ca0). **Endoperidial body** 11–23 mm, globose, sessile, very light grey (#fdfdfd) to dark grey (#a0a0a0), with lighter reticulation. Endoperidial surface with some protruding hyphae, endoperidium is interwoven by transparent hyphae, fibrous. **Peristome**

fibrillose, unpleated, wide conical, with obvious oral margin ring. Columella obvious very light grey (#f4f4f4 to #e0e0e0). **Eucapillitium hyphae** 1.0–5.5 μm , thick-walled, with distinct cavities, smooth, the ends tapering and are bluntly rounded (Fig. 1).

Basidiospores globose: Holotype (40/2/1) 4.5–5.3–6.0 \times (4.5)5.0–5.4–6.0 μm , Q = (0.80)0.83–1.12(1.14), Qm= 0.98 \pm 0.08, n = 40, including spines truncated at the apex ornamentation, with 0.2–0.5 μm high warts, ornamentation isolated or coalescing crest-like warts. Basidia not observed.

Diagnosis

Geastrum suae is characterised by long stipes and larger basidiomata; Pseudoparenchymatous layer is pink, smooth; globose endoperidial body, grey; the ends of eucapillitium hyphae taper and are bluntly rounded; and they live in groups.

Etymology

The species is named suae (Lat.), in memory of the Chinese mycologist Prof. Hong-Yan Su, who kindly helped the authors in many ways and sadly passed away on 3 May 2022 during the preparation of the current paper.

Habit

It grows in groups on the ground in mixed coniferous and broad-leaved forests where there are *Alnus nepalensis* and *Pinus yunnanensis*, with thick humus. Currently, it is known only from Cangshan Mountain.

Analysis

Phylogenetic analysis

Firstly, we constructed the ML tree of *Geastrum* genus, based on ITS (1–540 bp) and nrLSU (541–1498 bp) genes and found that *G. suae* is in Sect. *Mycelioatroma*. The Maximum Likelihood bootstrap values (ML) equal to or greater than 70% are given above each node (Fig. 2), with the Final ML Optimisation Likelihood: -24127.230142. The aligned matrix had 856 distinct alignment patterns, with 6.78% completely undetermined characters or gaps. The base frequency and rate are as follows: A = 0.274187, C = 0.208839, G = 0.265219, T = 0.251755; rate AC = 1.202699, AG = 3.054698, AT = 1.472914, CG = 0.671195, CT = 5.726232, GT = 1.000000; gamma distribution shape: α = 0.269052. Therefore, we constructed the ML tree and Bayesian tree of Sect. *Mycelioatroma*, based on ITS and nrLSU genes and clarified the position of *G. suae* in this Section. The dataset is composed of ITS and nrLSU genes, comprising a total of 1478 characters including gaps, ITS (1–591 bp) and nrLSU (592–1478 bp), including 35 taxa with *Myriostoma coliforme* (MA-Fungi 83759) as the outgroup taxon (Fig. 3). The best fit model for the combined 2-gene dataset estimated and applied in the Bayesian analysis was GTR+I+G, lset nst = 6, rates = invgamma; prset statefreqpr = dirichlet

(1,1,1,1). The phylogenetic analysis of ML and BI produces similar topology. The combined dataset analysis of RAxML generates a best-scoring tree (Fig. 3), with the Final ML Optimisation Likelihood value of -7513.207751. The aligned matrix had 584 distinct alignment patterns, with 21.33% completely undetermined characters or gaps. The base frequency and rate are as follows: A = 0.272494, C = 0.207593, G = 0.257821, T = 0.262093; rate AC = 1.093594, AG = 2.765430, AT = 1.755140, CG = 0.441983, CT = 5.721217, GT = 1.000000; gamma distribution shape: α = 0.243957. Bootstrap support values with ML greater than 70% and Bayesian posterior probabilities (PP) greater than 0.95 are given above the nodes (Fig. 3).

Phylogenetic analysis showed that four new collections of *G. suae* clustered together with high bootstrap support and are sister to *G. rubellum* with good bootstrap support (74% ML/1 PP Fig. 3).

Discussion

Geastrum suae can be easily recognised by the basidiomata with pink neat, smooth 6-lobed ectoderm, globose sessile endoperidium and longer prosthecae.

In the phylogenetic inferences, *Geastrum suae* is sister to *G. rubellum*, which is known from the biome Tropical and Subtropical Moist Broadleaf Forests in Brazil (Accioly et al. 2019) (Fig. 3). Morphologically, both species share similar characteristics of the mesopodal basidiomata, but *G. rubellum* has reddish to brownish exoperidium with longer exoperidium hairs. *G. suae* hardly has such hairs and the reddish pseudoparenchymatous layer in *G. rubellum* also clearly differentiates *G. suae*. Not only that, but *G. rubellum* also has reddish to brownish exoperidium with a verrucose to hairy mycelial layer, while the exoperidium of *G. suae* is almost smooth. Their size is different, the expanded basidiomata saccate of *G. rubellum* being 10 mm high \times 8.5–30 mm wide, while *G. suae* is 35–70 mm high \times 18–37 mm wide. The warts on the basisiospore of *G. suae* are shorter than those of *G. rubellum*. The pseudoparenchymatous layer of *G. rubellum* is pure (or mostly pure) pink (#fa007d) when fresh, brownish-grey when dried, but is very pale red (#ffcccd) for *G. suae*. The ITS comparison between our specimen (KUN-HKAS 123795) and *G. rubellum* (LIP: PAM/MART 12.100) revealed a 53 bp difference in a total of 542 bp. The nrLSU comparison between *G. suae* (KUN-HKAS 123795) and *G. rubellum* (LIP: PAM/MART 12.100) revealed 11 bp difference in a total of 809 bp (Accioly et al. 2019). It is worth noting that *G. rubellum* is distributed in the Neotropics (Accioly et al. 2019). Combined with the above analysis, we introduce *Geastrum suae* as a new species.

Acknowledgements

The research was financed by the National Natural Science Foundation of China (Project No. 32060006). This study was also supported by the Yunnan Fundamental Research Projects (grant No. 202201BC070001). The authors thank Kai Wang for the assistance in

sample collection, thank Dan-Feng Bao for sharing the knowledge of phylogeny and thank Long-Li Li for giving instruction for microscopic measurement.

References

- Accioly T, Sousa JO, Moreau PA, Lécuru C, da Silva BDB, Roy M, Gardes M, Baseia IG, Martín MP (2019) Hidden fungal diversity from the Neotropics: *Gastrum hirsutum*, *G. schweinitzii* (Basidiomycota, Geastrales) and their allies. PLOS One 14 (2): e0211388. <https://doi.org/10.1371/journal.pone.0211388>
- Assis NM, Freitas-Neto JF, Sousa JO, Barbosa FR, Baseia IG (2019) *Gastrum hyalinum* (Basidiomycota, Geastraceae), a new species from Brazilian Southern Amazon. Studies in Fungi 4 (1): 83-89. <https://doi.org/10.5943/sif/4/1/11>
- Cabral TS, da Silva BDB, Marinho P, Baseia IG (2014a) *Gastrum rusticum* (Geastraceae, Basidiomycota), a new earthstar fungus in the Brazilian Atlantic rainforest - a molecular analysis. Nova Hedwigia 98: 265-272. <https://doi.org/10.1127/0029-5035/2013/0158>
- Cabral TS, Silva BDB, Ishikawa NK, Alfredo DDS, Neto RB, Clement CR, Baseia IG (2014b) A new species and new records of gasteroid fungi (Basidiomycota) from Central Amazonia, Brazil. Phytotaxa 183 (4): 239-25. <https://doi.org/10.11646/phytotaxa.183.4.3>
- Cabral TS, Sousa J, da Silva BDB, Martín MP, Clement CR, Baseia IG (2017) A remarkable new species of *Gastrum* with an elongated branched stipe. Mycoscience 58 (5): 344-350. <https://doi.org/10.1016/j.myc.2017.03.004>
- Cabral TS, Sousa JDO, da Silva BDB, Martín MP, Clement CR, Goulart Baseia I (2022) Updates on *Gastrum* sect. *Exareolata*, with a description of a striking new species from the Neotropics. Plant Systematics and Evolution 308: 23. <https://doi.org/10.1007/s00606-022-01814-2>
- Crous PW, Wingfield MJ, Burgess TI, Hardy GESJ, Crane C, Barrett S, Cano-Lira JF, Le Roux JJ, Thangavel R, Guarro J, Stchigel AM, Martín MP, Alfredo DS, Barber PA, Barreto RW, Baseia IG, Cano-Canals J, Cheewangkoon R, Ferreira RJ, Gené J, Lechat C, Moreno G, Roets F, Shivas RG, Sousa JO, Tan YP, Wiederhold NP, Abell SE, Accioly T, Albizu JL, Alves JL, Antoniolli ZI, Aplin N, Araújo J, Arzanlou M, Bezerra JDP, Bouchara J, Carlavilla JR, Castillo A, Castroagudín VL, Ceresini PC, Claridge GF, Coelho G, Coimbra VRM, Costa LA, da Cunha KC, da Silva SS, Daniel R, de Beer ZW, Dueñas M, Edwards J, Enwistle P, Fiuza PO, Fournier J, García D, Gibertoni TB, Giraud S, Guevara-Suarez M, Gusmão LFP, Haituk S, Heykoop M, Hirooka Y, Hofmann TA, Houbraken J, Hughes DP, Kautmanová I, Koppel O, Koukol O, Larsson E, Latha KPD, Lee DH, Lisboa DO, Lisboa WS, López-Villalba Á, Maciel JLN, Manimohan P, Manjón JL, Marincowitz S, Marney TS, Meijer M, Miller AN, Olariaga I, Paiva LM, Piepenbring M, Poveda-Molero JC, Raj KNA, Raja HA, Rougeron A, Salcedo I, Samadi R, Santos TAB, Scarlett K, Seifert KA, Shuttleworth LA, Silva GA, Silva M, Siqueira JPZ, Souza-Motta CM, Stephenson SL, Sutton DA, Tamakeaw N, Telleria MT, Valenzuela-Lopez N, Viljoen A, Visagie CM, Vizzini A, Wartchow F, Wingfield BD, Yurchenko E, Zamora JC, Groenewald JZ (2016) Fungal Planet description sheets: 469-557. Persoonia 37: 218-403. <https://doi.org/10.3767/003158516X694499>
- Crous PW, Wingfield MJ, Burgess TI, Carnegie AJ, Hardy GESJ, Smith D, Summerell BA, Cano-Lira JF, Guarro J, Houbraken J, Lombard L, Martín MP, Sandoval-Denis M,

Alexandrova AV, Barnes CW, Baseia IG, Bezerra JDP, Guarnaccia V, May TW, Hernández-Restrepo M, Stchigel AM, Miller AN, Ordoñez ME, Abreu VP, Accioly T, Agnello C, Agustín Colmán A, Albuquerque CC, Alfredo DS, Alvarado P, Araújo-Magalhães GR, Arauzo S, Atkinson T, Barili A, Barreto RW, Bezerra JL, Cabral TS, Camello Rodríguez F, Cruz RHSF, Daniëls PP, da Silva BDB, de Almeida DAC, de Carvalho Júnior AA, Decock CA, Delgat L, Denman S, Dimitrov RA, Edwards J, Fedosova AG, Ferreira RJ, Firmino AL, Flores JA, García D, Gené J, Giraldo A, Góis JS, Gomes AAM, Gonçalves CM, Gouliamova DE, Groenewald M, Guéorguiev BV, Guevara-Suarez M, Gusmão LFP, Hosaka K, Hubka V, Huhndorf SM, Jadan M, Jurjević Ž, Kraak B, Kučera V, Kumar TKA, Kušan I, Lacerda SR, Lamertthon S, Lisboa WS, Loizides M, Luangsa-Ard JJ, Lysková P, Mac Cormack WP, Macedo DM, Machado AR, Malysheva EF, Marinho P, Matočec N, Meijer M, Mešić A, Mongkolsamrit S, Moreira KA, Morozova OV, Nair KU, Nakamura N, Noisripoon W, Olariaga I, Oliveira RJV, Paiva LM, Pawar P, Pereira OL, Peterson SW, Prieto M, Rodríguez-Andrade E, Rojo De Blas C, Roy M, Santos ES, Sharma R, Silva GA, Souza-Motta CM, Takeuchi-Kaneko Y, Tanaka C, Thakur A, Smith MT, Tkalcic Z, Valenzuela-Lopez N, van der Kleij P, Verbeken A, Viana MG, Wang XW, Groenewald JZ (2017) Fungal Planet description sheets: 625-715.

Persoonia 39: 270-467. <https://doi.org/10.3767/persoonia.2017.39.11>

- Crous PW, Wingfield MJ, Burgess TI, Hardy GESJ, Gené J, Guarro J, Baseia IG, García D, Gusmão LFP, Souza-Motta CM, Thangavel R, Adamčík S, Barili A, Barnes CW, Bezerra JDP, Bordallo JJ, Cano-Lira JF, de Oliveira RJV, Ercole E, Hubka V, Iturrieta-González I, Kubátová A, Martín MP, Moreau P, Morte A, Ordoñez ME, Rodríguez A, Stchigel AM, Vizzini A, Abdollahzadeh J, Abreu VP, Adamčíková K, Albuquerque GMR, Alexandrova AV, Álvarez Duarte E, Armstrong-Cho C, Banniwa S, Barbosa RN, Bellanger J, Bezerra JL, Cabral TS, Caboř M, Caicedo E, Cantillo T, Carnegie AJ, Carmo LT, Castañeda-Ruiz RF, Clement CR, Čmoková A, Conceição LB, Cruz RHSF, Damm U, da Silva BDB, da Silva GA, da Silva RMF, de A Santiago ALCM, de Oliveira LF, de Souza CAF, Déniel F, Dima B, Dong G, Edwards J, Félix CR, Fournier J, Gibertoni TB, Hosaka K, Iturriaga T, Jadan M, Jany J, Jurjević Ž, Kolařík M, Kušan I, Landell MF, Leite Cordeiro TR, Lima DX, Loizides M, Luo S, Machado AR, Madrid H, Magalhães OMC, Marinho P, Matočec N, Mešić A, Miller AN, Morozova OV, Neves RP, Nonaka K, Nováková A, Oberlies NH, Oliveira-Filho JRC, Oliveira TGL, Papp V, Pereira OL, Perrone G, Peterson SW, Pham THG, Raja HA, Raudabaugh DB, Řehulka J, Rodríguez-Andrade E, Saba M, Schauflerová A, Shivas RG, Simonini G, Siqueira JPZ, Sousa JO, Stajsic V, Svetasheva T, Tan YP, Tkalcic Z, Ullah S, Valente P, Valenzuela-Lopez N, Abrinbana M, Viana Marques DA, Wong PTW, Xavier de Lima V, Groenewald JZ (2018a) Fungal Planet description sheets: 716-784. Persoonia 40: 239-393. <https://doi.org/10.3767/persoonia.2018.40.10>
- Crous PW, Luangsa-Ard JJ, Wingfield MJ, Carnegie AJ, Hernández-Restrepo M, Lombard L, Roux J, Barreto RW, Baseia IG, Cano-Lira JF, Martín MP, Morozova OV, Stchigel AM, Summerell BA, Brandrud TE, Dima B, García D, Giraldo A, Guarro J, Gusmão LFP, Khamsuntorn P, Noordeloos ME, Nuankaew S, Pinruan U, Rodríguez-Andrade E, Souza-Motta CM, Thangavel R, van Iperen AL, Abreu VP, Accioly T, Alves JL, Andrade JP, Bahram M, Baral H, Barbier E, Barnes CW, Bendiksen E, Bernard E, Bezerra JDP, Bezerra JL, Bizio E, Blair JE, Bulyonkova TM, Cabral TS, Caiafa MV, Cantillo T, Colmán AA, Conceição LB, Cruz S, Cunha AOB, Darveaux BA, da Silva AL, da Silva GA, da Silva GM, da Silva RMF, de Oliveira RJV, Oliveira RL, De Souza JT,

- Dueñas M, Evans HC, Epifani F, Felipe MTC, Fernández-López J, Ferreira BW, Figueiredo CN, Filippova NV, Flores JA, Gené J, Ghorbani G, Gibertoni TB, Glushakova AM, Healy R, Huhndorf SM, Iturrieta-González I, Javan-Nikkhah M, Juciano RF, Jurjević Ž, Kachalkin AV, Keochanpheng K, Krisai-Greilhuber I, Li Y, Lima AA, Machado AR, Madrid H, Magalhães OMC, Marbach PAS, Melanda GCS, Miller AN, Mongkolsamrit S, Nascimento RP, Oliveira TGL, Ordoñez ME, Orzes R, Palma MA, Pearce CJ, Pereira OL, Perrone G, Peterson SW, Pham THG, Piontelli E, Pordel A, Quijada L, Raja HA, Rosas de Paz E, Ryvarden L, Saitta A, Salcedo SS, Sandoval-Denis M, Santos TAB, Seifert KA, Silva BDB, Smith ME, Soares AM, Sommai S, Sousa JO, Suetrong S, Susca A, Tedersoo L, Telleria MT, Thanakitpipattana D, Valenzuela-Lopez N, Visagie CM, Zapata M, Groenewald JZ (2018b) Fungal Planet description sheets: 785-867. Persoonia 41: 238-417. <https://doi.org/10.3767/persoonia.2018.41.12>
- Dai FL (1979) Chinese Fungus Collection. Science Press, Beijing. [In Chinese].
 - da Silva BDB, Sousa JO, Baseia IG (2011) Discovery of *Gastrum xerophilum* from the Neotropics. Mycotaxon 118: 355-359. <https://doi.org/10.5248/118.355>
 - da Silva BDB, Cabral TS, Marinho P, Ishikawa NK, Baseia IG (2013) Two new species of *Gastrum* (Gastraceae, Basidiomycota) found in Brazil. Nova Hedwigia 96: 445-456. <https://doi.org/10.1127/0029-5035/2013/0089>
 - de León PP (1968) A revision of the Gastraceae. Fieldiana Botany 31: 303-349.
 - Deng SQ (1963) Fungal in China. Science Press, Beijing. [In Chinese].
 - de Toledo LD, Castellano M (1996) A revision of the genera *Radiigera* and *Pyrenogaster*. Mycologia 88 (5): 963-88. <https://doi.org/10.2307/3760983>
 - Finy P, Papp V, Knapp D, Bóka K, Kovács G, Dima B (2021) *Gastrum dolomiticum*, a new earthstar species from Central Europe. Plant Systematics and Evolution 307 (4): 43. <https://doi.org/10.1007/s00606-021-01766-z>
 - Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts. Molecular Ecology 2 (2): 113-118. <https://doi.org/10.1111/j.1365-294x.1993.tb00005.x>
 - Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95-98. https://doi.org/10.14601/Phytopathol_Mediterr-14998u1.29
 - Han BX, Bau T (2016) New records of *Gastrum* from China. Mycosistema 35 (11): 1416-1424. [In Chinese]. <https://doi.org/10.13346/j.mycosistema.150234>
 - Hosaka K, Bates ST, Beever RE, Castellano MA, Colgan W, Domínguez LS, Nouhra ER, Geml J, Giachini AJ, Kenney SR, Simpson NB, Spatafora JW, Trappe JM (2006) Molecular phylogenetics of the gomphoid-phalloid fungi with an establishment of the new subclass Phallomycetidae and two new orders. Mycologia 98 (6): 949-59. <https://doi.org/10.3852/mycologia.98.6.949>
 - Hosaka K, Castellano MA (2008) Molecular phylogenetics of Geastrales with special emphasis on the position of Sclerogaster. Bulletin of the National Museum of Nature and Science 34: 161-173.
 - Jeppson M, Nilsson RH, Larsson E (2013) European earthstars in Geastraceae (Geastrales, Phallomycetidae) – a systematic approach using morphology and molecular sequence data. Systematics and Biodiversity 11 (4): 437-465. <https://doi.org/10.1080/14772000.2013.857367>

- Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30 (4): 772-780. <https://doi.org/10.1093/molbev/mst010>
- Miller M, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010 Gateway Computing Environments Workshop (GCE)8. <https://doi.org/10.1109/gce.2010.5676129>
- Nylander JA, Wilgenbusch JC, Warren DL, Swofford DL (2008) AWTY (are we there yet?): a system for graphical exploration of MCMC convergence in Bayesian phylogenetics. *Bioinformatics* 24 (4): 581-583. <https://doi.org/10.1093/bioinformatics/btm388>
- Persoon CH (1794) Neuer Versuch einer systematischen Einteilung der Schwämme. *Neues Magazin für die Botanik* 1: 63-128.
- Rambaut A (2012) FigTree version 1.4.0. <http://tree.bio.ed.ac.uk/software/figtree/>
- Rannala B, Yang Z (1996) Probability distribution of molecular evolutionary trees: A new method of phylogenetic inference. *Journal of Molecular Evolution* 43 (3): 304-311. <https://doi.org/10.1007/bf02338839>
- Rodrigues DJ, Barbosa FR, Noronha JC, Carpanedo RS, Tourinho AN, Battiroldi LD (2021) Biodiversidade da Estação Ecológica do Rio Ronuro - Biodiversity of the Rio Ronuro Ecological Station. Fundação UNISELVA
- Ronquist F, Teslenko M, van der Mark P, Ayres D, Darling A, Höhna S, Larget B, Liu L, Suchard M, Huelsenbeck J (2012) MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a Large Model Space. *Systematic Biology* 61 (3): 539-542. <https://doi.org/10.1093/sysbio/sys029>
- Sousa J, da Silva BDB, Alfredo D, Baseia I (2014) New records of Geastraceae (Basidiomycota, Phallomycetidae) from Atlantic Rainforest remnants and relicts of Northeastern Brazil. *Darwiniana, nueva serie* 2 (2): 207-221. <https://doi.org/10.14522/darwiniana/2014.22.595>
- Sousa J, Suz L, García M, Alfredo D, Conrado L, Marinho P, Ainsworth AM, Baseia I, Martín M (2017) More than one fungus in the pepper pot: Integrative taxonomy unmasks hidden species within *Myriostoma califorme* (Geastraceae, Basidiomycota). *PLOS One* 12 (6): e0177873. <https://doi.org/10.1371/journal.pone.0177873>
- Sousa JO, Baracho GS, Baseia IG (2015) *Geastrum laevigatum*: a new earthstar fungus with uncommon smooth spores. *Mycosphere* 6 (4): 501-507. <https://doi.org/10.5943/mycosphere/6/4/12>
- Sunhede S (1989) Geastraceae (Basidiomycotina): Morphology, ecology, and systematics with special emphasis on the North European species. *Syn. Fungorum* 1: 1-534.
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of bacteriology* 172 (8): 4238-4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Wang X, Bau T (2023) Seven New Species of the Genus *Geastrum* (Geastrales, Geastraceae) in China. *Journal of Fungi* 9 (2): 251.
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols* 1 (1): 315-322. <https://doi.org/10.1016/b978-0-12-372180-8.50042-1>
- Wijayawardene N, Hyde K, Dai D, Sánchez-García M, Goto B, Saxena R, Erdoğdu M, Selçuk F, Rajeshkumar K, Aptroot A, Błaszkowski J, Boonyuen N, da Silva G, de Souza

F, Dong W, Ertz D, Haelewaters D, Jones E, Karunarathna S, Kirk P, Kukwa M, Kumla J, Leontyev D, Lumbsch H, Maharachchikumbura S, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro J, Oehl F, Pawłowska J, Pem D, Pfiegl W, Phillips A, Pošta A, He M, Li J, Raza M, Sruthi O, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkalc̆ec Z, Tokarev Y, Wanasinghe D, Wijesundara D, Wimalaseana S, Madrid H, Zhang G, Gao Y, Sánchez-Castro I, Tang L, Stadler M, Yurkov A, Thines M (2022) Outline of Fungi and fungus-like taxa – 2021. *Mycosphere* 13 (1): 53-453. <https://doi.org/10.5943/mycosphere/13/1/2>

- Zamora JC, Calonge FDD, Martín MP (2013) New sources of taxonomic information for earthstars (*Gastrum*, Geastraceae, Basidiomycota): phenoloxidases and rhizomorph crystals. *Phytotaxa* 132 (1): 1-20. <https://doi.org/10.11646/phytotaxa.132.1.1>
- Zamora JC, de Diego Calonge F, Hosaka K, Martín MP (2014) Systematics of the genus *Gastrum* (Fungi: Basidiomycota) revisited. *Taxon* 63 (3): 477-497. <https://doi.org/10.12705/633.36>
- Zhaxybayeva O, Gogarten JP (2002) Bootstrap, Bayesian probability and maximum likelihood mapping: exploring new tools for comparative genome analyses. *BMC Genomics* 3 (1). <https://doi.org/10.1186/1471-2164-3-4>
- Zhou H, Li JQ, Hou CL (2022) Two new species of the genus *Gastrum* from the Yanshan Mountains in China. *Mycosistema* 41 (1): 1-16. [In Chinese]. <https://doi.org/10.13346/j.mycosistema.210194>
- Zhou TS, Chen YH, Zhao LZ, Fu H, Yang B (2007) Mycology of China. 36. Science Press, Beijing. [In Chinese].




Figure 1.

Geastrum suae (KUN-HKAS 123795, holotype). **a** fresh unexpanded fruiting bodies; **b, c** fresh mature fruiting bodies; **d** mycelial layer, fibrous layer and pseudoparenchymatous layer; **e** hyphae of mycelial layer; **f** pseudoparenchymatous layer (cells in the stack); **g, h** eucapillitium hyphae; **i-k** basidiospores (LM); **l-n** basidiospores (SEM). Scale bars: a = 10 mm; b, c = 20 mm; d = 80 μ m; f, g, i-k = 10 μ m; h = 70 μ m; l = 1 μ m; m, n = 500 nm.



Figure 2.

Phylogenetic tree of *Geastrum* species and related taxa, based on ITS and nrLSU sequence data.




Table 1.

Amplification primers information used in this study.

Gene	Primer	Primer sequence (5'-3')	References
ITS	ITS1F	CTTGGTCATTTAGAGGAAGTAA	Gardes and Bruns (1993)
	ITS4	TCCTCCGCTTATTGATATGC	White et al. (1990)
nrLSU	LR0R	ACCCGCTGAACCTTAAGC	Vilgalys and Hester (1990)
	LR5	ATCCTGAGGGAAACTTC	Vilgalys and Hester (1990)

Table 2.

Species, specimens, Collection locality and GenBank accession numbers of sequences used in this study (newly-generated sequences are indicated in bold).

Species	Strain/Voucher	Collection locality	GenBank Accession No.	
			ITS	nrLSU
<i>Geastrum mirabile</i>	strain: 228-394	Japan	AB509736	-
<i>Geastrum javanicum</i>	TNS:TKG-GE-90902	Japan	JN845100	JN845218
<i>Geastrum mirabile</i>	TNS:KH-JPN10-714	Japan	JN845109	JN845227
<i>Geastrum parvistriatum</i>	MA-Fungi 69583	Spain	JN943160	JN939560
<i>Geastrum parvistriatum</i>	Herb. Zamora 272	Spain	JN943162	JN939572
<i>Geastrum striatum</i>	Herb. Zamora 257	Spain	JN943164	JN939557
<i>Geastrum campestre</i>	Herb. Zamora 283	Spain	JN943167	JN939575
<i>Geastrum aff. arenarium</i>	Herb. Zamora 76	Spain	KF988338	KF988470
<i>Geastrum lageniforme</i>	Herb. Zamora 316	Spain	KF988339	KF988514
<i>Geastrum cf. calceum</i>	UFRN-Fungos 723	Brazil	KF988340	KF988477
<i>Geastrum cf. calceum</i>	MA-Fungi 83761	Argentina	KF988341	KF988478
<i>Geastrum aff. hariotii</i>	Börge Petterson 2070	Mozambique	KF988342	KF988507
<i>Geastrum cf. saccatum</i>	Herb. Sunhede 7749	Australia	KF988343	KF988556
<i>Geastrum hieronymi</i>	MA-Fungi 83767	Argentina	KF988344	KF988509
<i>Geastrum cf. stipitatum</i>	Herb. Zamora 528	Brazil	KF988345	KF988576
<i>Geastrum albonigrum</i>	MA-Fungi 36140-2	Panama	KF988349	KF988468
<i>Geastrum aff. arenarium</i>	MA-Fungi 68191	Spain	KF988350	KF988469
<i>Geastrum cf. arenarium</i>	MA-Fungi 83760	Argentina	KF988351	KF988471
<i>Geastrum argentinum</i>	LPS 48446	Argentina	KF988352	KF988472
<i>Geastrum argentinum</i>	MA-Fungi 82605	Argentina	KF988353	KF988473
<i>Geastrum berkeleyi</i>	MA-Fungi 74668	Spain	KF988354	KF988474
<i>Geastrum berkeleyi</i>	Herb. Sunhede 7724	Sweden	KF988355	KF988475
<i>Geastrum berkeleyi</i>	Herb. Zamora 504	Sweden	KF988356	KF988476
<i>Geastrum campestre</i>	Herb. Sunhede 7575	Sweden	KF988357	KF988479
<i>Geastrum campestre</i>	MICH 28566	USA	KF988358	KF988480
<i>Geastrum corollinum</i>	MA-Fungi 5746	Spain	KF988359	KF988481
<i>Geastrum corollinum</i>	Herb. Sunhede 7744	Sweden	KF988360	KF988482
<i>Geastrum coronatum</i>	Herb. Zamora 266	Spain	KF988361	KF988483
<i>Geastrum coronatum</i>	Herb. Zamora 522	Sweden	KF988362	KF988484
<i>Geastrum coronatum</i>	MICH 28567	USA	KF988363	KF988485
<i>Geastrum aff. coronatum</i>	MICH 72012	USA	KF988364	KF988486
<i>Geastrum aff. coronatum</i>	MICH 72014	USA	KF988365	KF988487

<i>Geastrum elegans</i>	Herb. Zamora 189	Spain	KF988366	KF988488
<i>Geastrum elegans</i>	UPS F-560810	Sweden	KF988367	KF988489
<i>Geastrum entomophilum</i>	MA-Fungi 70785	Brazil	KF988368	KF988490
<i>Geastrum fimbriatum</i>	Herb. Zamora 234	Spain	KF988369	KF988491
<i>Geastrum fimbriatum</i>	Herb. Sunhede 7739	Sweden	KF988370	KF988492
<i>Geastrum flexuosum</i>	UPS F-119844	Sweden	KF988371	KF988493
<i>Geastrum floriforme</i>	MA-Fungi 69173	Spain	KF988372	KF988494
<i>Geastrum floriforme</i>	Herb. Zamora 453	Spain	KF988373	KF988495
<i>Geastrum fornicatum</i>	Herb. Zamora 255	Spain	KF988374	KF988496
<i>Geastrum fornicatum</i>	MA-Fungi 30749	Spain	KF988375	KF988497
<i>Geastrum fuscogleba</i>	NY Trappe 1071	USA	KF988376	KF988498
<i>Geastrum fuscogleba</i>	NY Trappe 9500	USA	KF988377	KF988499
<i>Geastrum glaucescens</i>	MA-Fungi 83762	Argentina	KF988378	KF988500
<i>Geastrum glaucescens</i>	MA-Fungi 83763	Argentina	KF988379	KF988501
<i>Geastrum aff. glaucescens</i>	MA-Fungi 83764	Argentina	KF988380	KF988502
<i>Geastrum hariotii</i>	MA-Fungi 83765	Argentina	KF988381	KF988504
<i>Geastrum aff. hariotii</i>	MA-Fungi 78296	Brazil	KF988382	KF988505
<i>Geastrum aff. hariotii</i>	MA-Fungi 78289	Brazil	KF988383	KF988506
<i>Geastrum hieronymi</i>	MA-Fungi 83766	Argentina	KF988384	KF988508
<i>Geastrum kotlabae</i>	MA-Fungi 39563	Spain	KF988385	KF988510
<i>Geastrum kotlabae</i>	Herb. Zamora 440	Spain	KF988386	KF988511
<i>Geastrum aff. kotlabae</i>	MA-Fungi 33300	Tanzania	KF988387	KF988512
<i>Geastrum lageniforme</i>	Herb. Zamora 207	Spain	KF988388	KF988513
<i>Geastrum aff. lageniforme</i>	MA-Fungi 83768	Argentina	KF988389	KF988516
<i>Geastrum aff. lageniforme</i>	COFC Hama 327	Niger	KF988390	KF988517
<i>Geastrum aff. lageniforme</i>	MA-Fungi 83770	Argentina	KF988391	KF988518
<i>Geastrum aff. lageniforme</i>	MA-Fungi 83769)	Argentina	KF988392	KF988519
<i>Geastrum aff. lageniforme</i>	MA-Fungi 78398	Portugal	KF988393	KF988520
<i>Geastrum aff. lageniforme</i>	Herb. Ribes 221210-01	Spain	KF988394	KF988521
<i>Geastrum melanocephalum</i>	Herb. Zamora 34	Spain	KF988395	KF988522
<i>Geastrum melanocephalum</i>	Herb. Sunhede 7737	Sweden	KF988396	KF988523
<i>Geastrum michelianum</i>	Herb. Sunhede 7738	Sweden	KF988397	KF988524
<i>Geastrum michelianum</i>	Herb. Zamora 227	Spain	KF988398	KF988525
<i>Geastrum aff. michelianum</i>	MA-Fungi 83771	Argentina	KF988399	KF988527
<i>Geastrum minimum</i>	Herb. Zamora 191	Spain	KF988400	KF988528
<i>Geastrum minimum</i>	Herb. Sunhede 7746	Sweden	KF988401	KF988529
<i>Geastrum minimum</i>	MICH 72010	USA	KF988402	KF988530

<i>Geastrum minimum</i>	MICH 28119	Spain	KF988403	KF988531
<i>Geastrum minimum</i>	MA-Fungi 31530	USA	KF988404	KF988532
<i>Geastrum minimum</i>	MA-Fungi 86669	Sweden	KF988405	KF988533
<i>Geastrum morganii</i>	Herb. Lebeuf HRL0177	Canada	KF988406	KF988534
<i>Geastrum aff. morganii</i>	Herb. Zamora 367	Spain	KF988407	KF988535
<i>Geastrum aff. morganii</i>	Herb. Zamora 525	Spain	KF988408	KF988536
<i>Geastrum aff. morganii</i>	MA-Fungi 83772	Argentina	KF988409	KF988537
<i>Geastrum aff. morganii</i>	MA-Fungi 83773	Argentina	KF988410	KF988538
<i>Geastrum ovalisporum</i>	MA-Fungi 47184	Bolivia	KF988411	KF988539
<i>Geastrum pectinatum</i>	Herb. Zamora 252	Spain	KF988412	KF988540
<i>Geastrum pectinatum</i>	UPS F-560803	Sweden	KF988413	KF988541
<i>Geastrum pectinatum</i>	UPS F-09935 (161483)	Tanzania	KF988414	KF988542
<i>Geastrum pectinatum</i>	MA-Fungi 83774	Argentina	KF988415	KF988543
<i>Geastrum pleosporum</i>	MA-Fungi 56971	Cameroon	KF988416	KF988544
<i>Geastrum pouzarii</i>	MA-Fungi 2944	Czechoslovakia	KF988417	KF988545
<i>Geastrum pouzarii</i>	Herb. Sunhede 7494	Czechoslovakia	KF988418	KF988546
<i>Geastrum pseudolimbatum</i>	Herb. Zamora 231	Spain	KF988419	KF988547
<i>Geastrum pseudolimbatum</i>	UPS F-560804	Sweden	KF988420	KF988548
<i>Geastrum quadrifidum</i>	Herb. Zamora 170	Spain	KF988421	KF988549
<i>Geastrum quadrifidum</i>	MA-Fungi 86671	Sweden	KF988422	KF988550
<i>Geastrum quadrifidum</i>	MICH 72512	USA	KF988423	KF988551
<i>Geastrum rufescens</i>	Herb. Zamora 253	Spain	KF988424	KF988552
<i>Geastrum rufescens</i>	Herb. Zamora 274	Spain	KF988425	KF988553
<i>Geastrum cf. saccatum</i>	MA-Fungi 47185-2	Bolivia	KF988426	KF988554
<i>Geastrum cf. saccatum</i>	MA-Fungi 83775	Argentina	KF988427	KF988555
<i>Geastrum cf. saccatum</i>	UPS F-530056	Japan	KF988428	KF988558
<i>Geastrum cf. saccatum</i>	MA-Fungi 83777	Argentina	KF988429	KF988559
<i>Geastrum cf. saccatum</i>	Herb. Zamora 260	Spain	KF988430	KF988560
<i>Geastrum cf. saccatum</i>	Herb. Zamora 461	Spain	KF988431	KF988561
<i>Geastrum cf. saccatum</i>	COFC Hama 343	Niger	KF988432	KF988562
<i>Geastrum cf. saccatum</i>	MA-Fungi 83778	Argentina	KF988433	KF988563
<i>Geastrum schmidelli</i>	Herb. Zamora 279	Spain	KF988434	KF988564
<i>Geastrum schmidelli</i>	UPS F-560805	Sweden	KF988435	KF988565
<i>Geastrum cf. schweinitzii</i>	S Henrik Kylin 1983 30.X	Papua New Guinea	KF988436	KF988566
<i>Geastrum cf. schweinitzii</i>	MA-Fungi 83779	Argentina	KF988437	KF988567
<i>Geastrum cf. schweinitzii</i>	MA-Fungi 36141	Panama	KF988438	KF988568
<i>Geastrum cf. schweinitzii</i>	MA-Fungi 83780	Argentina	KF988439	KF988569

<i>Geastrum smardae</i>	Herb. Lebeuf HRL 0160	Canada	KF988440	KF988573
<i>Geastrum smardae</i>	Herb. Zamora 527	Spain	KF988441	KF988574
<i>Geastrum smithii</i>	MA-Fungi 83783	Argentina	KF988442	KF988575
<i>Geastrum striatum</i>	MA-Fungi 86672	Sweden	KF988443	KF988577
<i>Geastrum "triplex"</i>	UPS F-014630 (213863)	Madagascar	KF988444	KF988578
<i>Geastrum "triplex"</i>	MA-Fungi 83784	Argentina	KF988445	KF988579
<i>Geastrum cf. velutinum</i>	MA-Fungi 83785	Argentina	KF988446	KF988581
<i>Geastrum cf. velutinum</i>	MA-Fungi 83786	Argentina	KF988447	KF988582
<i>Geastrum cf. velutinum</i>	Herb. Ribes 311207-62	Spain	KF988448	KF988583
<i>Geastrum cf. velutinum</i>	MA-Fungi 83787	Peru	KF988449	KF988584
<i>Geastrum violaceum</i>	BAFC 51671	Argentina	KF988450	KF988585
<i>Geastrum violaceum</i>	MA-Fungi 82487	Argentina	KF988451	KF988586
<i>Geastrum sp.1</i>	MA-Fungi 83788	Argentina	KF988452	KF988587
<i>Geastrum sp.1</i>	MA-Fungi 83789	Argentina	KF988453	KF988588
<i>Geastrum sp.2</i>	MA-Fungi 31143	Spain	KF988454	KF988589
<i>Geastrum sp.2</i>	MA-Fungi 37546	Spain	KF988455	KF988590
<i>Geastrum sp.3</i>	MA-Fungi 83790	Argentina	KF988456	KF988591
<i>Geastrum sp.4</i>	MA-Fungi 83791	Peru	KF988457	KF988592
<i>Geastrum sp.5</i>	Herb. Zamora 145	Spain	KF988458	KF988593
<i>Geastrum sp.5</i>	Herb. Zamora 450	Spain	KF988459	KF988594
<i>Geastrum sp.6</i>	MA-Fungi 83792	Argentina	KF988460	KF988595
<i>Geastrum sp.7</i>	MA-Fungi 83793	Argentina	KF988461	KF988596
<i>Geastrum sp.7</i>	MA-Fungi 83794	Argentina	KF988462	KF988597
<i>Geastrum sp.8</i>	MA-Fungi 83795	Argentina	KF988463	KF988598
<i>Geastrum hirsutum</i>	UFRN-Fungos 1214	Brazil	KJ127029	-
<i>Geastrum javanicum</i>	UFRN-Fungos 1215	Brazil	KJ127031	-
<i>Geastrum minutisporum</i>	CORD14	Argentina	KM260664	-
<i>Geastrum minutisporum</i>	CORD15	Argentina	KM260665	-
<i>Geastrum pusillipilosum</i>	UFRN:Fungos 2315	Brazil	KX761175	KX761176
<i>Geastrum pusillipilosum</i>	UFRN:Fungos 2759	Brazil	KX761177	KX761178
<i>Geastrum piquiriunense</i>	UFRN:Fungos:2892	Brazil	MH260269	MH260270
<i>Geastrum hirsutum</i>	INPA:259950	Brazil	MH634993	MH635026
<i>Geastrum rubropusillum</i>	UFRN:Fungos:2308	Brazil	MH634994	MH635027
<i>Geastrum baculicrystallum</i>	UFRN:Fungos:2835	Brazil	MH634995	MH635028
<i>Geastrum brunneocapillatum</i>	UFRN:Fungos:2286	Brazil	MH634996	MH635029
<i>Geastrum rubellum</i>	UFRN:Fungos:2844	Brazil	MH634999	MH635031
<i>Geastrum neoamericanum</i>	UFRN:Fungos:2302	Brazil	MH635001	MH635040

<i>Geastrum courtecuissei</i>	LIP:FH 2004090503	Guadeloupe	MH635003	MH635033
<i>Geastrum rubellum</i>	LIP:CL/MART 8067B	Martinique	MH635009	-
<i>Geastrum rubellum</i>	LIP:PAM/MART 12.100	Martinique	MH635010	MH635037
<i>Geastrum neoamericanum</i>	LIP:JLC 12030103	French	MH635014	MH635038
<i>Geastrum suae</i>	HKAS 123795 (Holotype)	China	ON529511	ON529515
<i>Geastrum suae</i>	HKAS 123794	China	ON529512	ON529516
<i>Geastrum suae</i>	HKAS 123793	China	ON529513	ON529517
<i>Geastrum suae</i>	HKAS 123796 (Paratype)	China	ON529514	ON529518
<i>Geastrum hariotii</i>	MA-Fungi 80070	Dominican Republic	-	KF988503
<i>Geastrum aff. lageniforme</i>	MA-Fungi 79056	Brazil	-	KF988515
<i>Geastrum cf. saccatum</i>	MA-Fungi 83776	Argentina	-	KF988557
<i>Geastrum cf. schweinitzii</i>	S Henrik Kylin 842	Fiji	-	KF988570
<i>Geastrum cf. velutinum</i>	MA-Fungi 73247	India	-	KF988580
<i>Geastrum michelianum</i>	Herb. Ribes 231208-31	Spain	-	KF988526
<i>Geastrum setiferum</i>	MA-Fungi 83781	Argentina	-	KF988571
<i>Geastrum setiferum</i>	MA-Fungi 83782	Argentina	-	KF988572
<i>Geastrum velutinum</i>	BJTC 221	China	-	MZ509382
<i>Geastrum velutinum</i>	BJTC 598	China	MZ508877	-
<i>Geastrum yanshanense</i>	BJTC 381	China	MZ508878	MZ509383
<i>Geastrum yanshanense</i>	BJTC 057	China	MZ508879	MZ509384
<i>Geastrum yanshanense</i>	BJTC 255	China	MZ508880	-
<i>Schenella pityophila</i>	Herb. Zamora 530	Spain	KF988346	KF988464
<i>Schenella pityophila</i>	Herb. Zamora 531	Spain	KF988347	KF988465
<i>Myriostoma coliforme</i>	MA-Fungi 83759	Argentina	KF988348	KF988467