

# Bats of Georgia - an occurrence dataset from 1835 through 2022

Ioseb Natradze<sup>‡</sup>, Alexander Buknikashvili<sup>‡</sup>, Giorgi Sheklashvili<sup>‡</sup>, Levan Mumladze<sup>‡</sup>

<sup>‡</sup> Institute of Zoology of Ilia State University, Tbilisi, Georgia

Corresponding author: Ioseb Natradze ([ioseb.natradze@iliauni.edu.ge](mailto:ioseb.natradze@iliauni.edu.ge))

Academic editor: Krizler Tanalgo

## Abstract

## Background

In Georgia, currently, 30 species of bats are recorded from four families and eleven genera. Although the oldest record of bats is from 1835 and continues until today, there are no comprehensive data available for bat diversity and distribution in Georgia. Thus, we aimed to fill that gap and make complete, expertly curated literature and our own published data openly available (through GBIF) for researchers and conservationists.

## New information

In this publication, out of 1987 records, 1243 (62.4%) are new and unpublished data. Generally, out of all records, 34% are literature and museum data and 66% are data collected by us. Additionally, for the first time in the history of the study of bats in Georgia, we initiated surveys in forested areas of the country.

## Keywords

Bats, Chiroptera, dataset, distribution, Caucasus, Georgia

## Introduction

Georgia is a part of the Caucasus biodiversity hotspot - one of 36 biodiversity hotspots recognised in the world, with incredible landscape and species diversity (Myers et al. 2000; Mittermeier et al. 2011; Noss et al. 2014). Although Georgia, as a part of the Caucasus, was distinguished as a biodiversity hotspot 22 years ago, the diversity and distribution of animal species remains poorly investigated (Mumladze et al. 2020). Even the inventory of vertebrate taxa (that were used by Myers et al. (2000) along with plants to

delineate biodiversity hotspots) is not yet satisfactorily done, leaving room to improve the understanding of species diversity, distribution and ecological requirements.

The oldest documented record of bats in Georgia dates back to 1835 when Nordmann (1840) reported four species. This was followed by surveys conducted by Kolenati in 1843 (Kolenati 1860). After that and for nearly 40 years, no new data were collected until the beginning of the 20<sup>th</sup> century (Radde 1899; Satunin 1903; Satunin 1908; Satunin 1912; Satunin 1915). As a result, 16 species for Georgia were indicated. Bat research even in the 20<sup>th</sup> century was not active, rather was sporadic and information was scattered in different Georgian and Russian publications: Chkhikvishvili (1926); Ognev 1928; Kuzyakin (1950); Janashvili (1953); Janashvili (1963); and these publications report 22 species for Georgia. Sporadic information about bat species occurrence is also given in the publications by non-bat researchers including Papava 1949; Papava 1953; Papava 1960; Matsaberidze 1961; Avaliani (1963); Matsaberidze and Khotenovsky 1966; Matsaberidze and Khotenovskii 1967; Avaliani (1969a); Avaliani (1969b); Avaliani (1970); Avaliani (1973); Perov 1975; Avaliani (1976); Matsaberidze (1976); Perov 1980; Perov (1983); Strelkov (1983); Strelkov (1988); and Shidlovsky (2013).

Systematic surveys of the bat fauna of Georgia were started at the end of the 20<sup>th</sup> century and resulted in a number of publications (Bukhnikashvili and Kandaurov 2002; Ivanitsky 2002; Bukhnikashvili et al. 2004; Gazaryan and Bukhnikashvili 2005; Bukhnikashvili et al. 2008; Gazaryan et al. 2008; Yavruyan et al. 2008; Bukhnikashvili et al. 2009; Ivanitsky 2010; Gazaryan et al. 2017; Ivanitsky 2017; Ivanitsky 2018; Imnadze et al. 2020; Urushadze et al. 2021). As a result of these studies, the number of bat species recorded in Georgia increased to 26 (Bukhnikashvili 2004). However, a considerable part of the results of bat surveys conducted during the last 25 years are either published in grey literature (i.e. publicly unavailable project reports) or kept unpublished by the authors. In addition to distribution data for other species, our unpublished data also provides records of four additional species (*Rhinolophus blasii*, *Myotis alcaethoe*, *Myotis davidii* and *Tadarida teniotis*) in Georgia that have not been previously documented in the literature. Thus, the goal of the present publication was to consolidate all available data about bat records of Georgia from 1835 through to 2022 into a comprehensive dataset and make it available through global and open-source databases such as GBIF (GBIF.org 2023) in order to facilitate further research and conservation of bats in Georgia.

## Sampling methods

**Description:** The dataset, prepared by Natradze et al. (2023), contains information about 1987 records of 30 species of four families and 11 genera collected from 1835 through to 2023 in country of Georgia (Fig. 1).

**Sampling description:** These records are based on literature published in Georgian, Russian and English languages, as well as data collected in the field by the authors during the last three decades. To collect field data, we employed various methods, including mist-netting, harp traps, visual inspection of both artificial and natural

underground and overground habitats and other shelters (Dietz and Kiefer 2016). Additionally, we used hand-held bat detectors, specifically the Pettesson D240x ultrasound bat model. In our dataset, we use data obtained through hand-held bat detectors, which were validated by visually confirming the presence of the bat. To ensure high data quality, all recorded data were included only if bats were identified at the species level.

**Quality control:** In the dataset, 37.4% of records are based on literature. For each of the literature records, we retrieved as much information as possible, such as sampling date, location, closest populated area, habitat etc. For most of the literature data (especially old ones), no exact geographic coordinates were given. However, since the vernacular names of sampling areas (i.e. names of subterranean objects) along with habitat descriptions were provided in many cases, we were able to exactly georeference a large number of sampling locations for 55.9% of literature records. On the other hand, not all records in literature are supplied with sampling dates and we were able to retrieve information on sampling dates for only 81.6% of literature records.

In the dataset, we provide location common names for all records. Geographic coordinates with 4 m accuracy are given for most of the records (74%), while for 26% (all literature data) of records, we have coordinates without accurate information. Additionally, 9% of records are given without the collecting dates. Record summary by species is given in Table 1, while the species records with metadata are provided at the GBIF web portal (Natradze et al. 2023).

Our database contains several cases that require further clarification regarding the identity of certain species, including: (i) some records of *Myotis nattereri* may actually pertain to *M. tschuliensis*, as suggested by Çoraman et al. (2019) and Kruskop and Solovyeva (2020); (ii) the identification of *M. davidii/mystacinus* may be erroneous due to their cryptic nature; and (iii) the potential existence of another species, *Miniopterus pallidus*, in eastern Georgia, as proposed by Šrámek et al. (2012). However, additional research, including DNA analysis, is required to confirm the identity of these species. Any modifications resulting from these investigations will be reflected in the subsequent version of the dataset.

## Geographic coverage

**Description:** The presented bat distribution dataset originated from the whole Georgian territory. Georgia (Fig. 1), covering an area of 69,700 km<sup>2</sup>, is located on the southern slopes of the Great Caucasus Mountain Range, Lesser Caucasus Mountains on the isthmus between the Black and Caspian Seas. It contains lowlands between the above-mentioned mountain ranges which include the Colchis lowland in the west (along the Black Sea Coast) and the Kura River lowland in the east. The land of Georgia covers an elevation range from sea level to approximately 5,184 m at Mount Shkhara. Two-thirds of the country is mountainous with an average height of 1200 m a.s.l.

Due to its diverse geography, the climate of the region varies greatly, from very humid lowlands and mountain forests in the west to dry forests and semi-deserts in the east and glaciated nival belts in the north. There are two zoogeographic subzones and three zoogeographic provinces in Georgia: the Circumboreal subzone (the Caucasus district of the European forest province) and Mediterranean subzone (the Anterior Asia district of the Mediterranean province and the Kura district of the Iran-Turan province) (Verestchagin 1959). Georgia has about 72 types of landscapes (Beruchashvili 2000): humid sub-tropic landscapes are in the western part; alpine landscapes are spread in the northern and north-eastern part; the typical Middle East treeless uplands are presented in the southern part; and semi-deserts of the Turanian type in the southeast part of the country.

The studied territory contains diverse bat habitats. Along with various kinds of forests (temperate broadleaf, evergreen and dry forests), particularly relevant are the western Great Caucasus slopes, which are represented by a number of Karst massifs that provide a large and diverse (yet only partly explored) subterranean environment suitable for bat species.

**Coordinates:** 40.946 and 43.818 Latitude; 39.660 and 46.933 Longitude.

## Temporal coverage

**Notes:** The dataset includes data collected from 1835 through to 2023. For all other records, time coverage could be divided into the following time periods, data collected in (i) 19<sup>th</sup>, (ii) 20<sup>th</sup> and (iii) 21<sup>st</sup> centuries. In the 19<sup>th</sup> century, there are 53 records which make up 2.7% of all records; in the 20<sup>th</sup> century, there are 231 records which make up 11.6% of all records and in the 21<sup>st</sup> century, there are 1523 records, which make up 76.7%. From the 1523 records, made in the 21<sup>st</sup> century, 1239 records are new, unpublished records, which make up 62.4% of all records.

## Usage licence

**Usage licence:** Open Data Commons Attribution License

## Data resources

**Data package title:** Bats of Georgia

**Resource link:** <https://www.gbif.org/dataset/8e1c23ba-5618-4bba-8bf-a195cce8dda0>

**Number of data sets:** 1

**Data set name:** batsofgeorgia

**Data format:** Excel table

## Data format version: V1.2

Column label	Column description
occurrenceID	Unique identifier of record.
kingdom	The full scientific name of the kingdom in which the taxon is classified.
phylum	The full scientific name of the phylum in which the taxon is classified.
class	The full scientific name of the class in which the taxon is classified.
order	The full scientific name of the order in which the taxon is classified.
family	The full scientific name of the subfamily in which the taxon is classified.
scientificName	Species full scientific (Latin) name including authorship and year.
locality	The specific description of the place of collection.
eventDate	Collection event date.
countryCode	Standard ISO 3166-1-alpha-2 country code
decimalLatitude	The geographic latitude (in decimal degrees).
decimalLongitude	The geographic longitude (in decimal degrees).
geodeticDatum	Geographic coordinates reference system EPSG.
coordinateUncertaintyInMetres	Coordinate measurement accuracy (metres in case of GPS recordings, NA - if manually georeferenced). However, see the field "dataGeneralisations" for further details
minimumElevationInMetres	Minimum elevation above sea level.
maximumElevationInMetres	Maximum elevation above sea level.
associatedReferences	Source for the particular record.
georeferenceSources	The system used during the georeferencing.
dataGeneralisations	According to the "Agreement on the Conservation of Populations of European Bats" (Eurobats, United Nations Environment Programme, Eurobats, May 2019), (Retrieved 7 August 2019), we intentionally reduced the precision of geographic coordinates in the dataset, while more precise information is available upon request.
basisOfRecord	The specific nature of the data record.
institutionCode	The code of the institution where data are stored.
collectionCode	The code of the collection.

## Additional information

Field data were collected under the permissions #2722/01; 2302/01; R/057-21, issued by the Ministry of Environmental Protection and Agriculture of Georgia.

## Acknowledgements

This work was supported by the Shota Rustaveli National Foundation of Georgia (SRNSFG) [FR-19-2295]. We would like to thank Mr. Andro Kandaurov (Institute of Zoology of Ilia State University) and Dr. Suren Gazaryan (UNEP/EUROBATS secretariat) for their input in the building of the dataset. We would like to thank Dr. David Bloom (VertNet: Sebastopol, California, US) who provided invaluable assistance with regard to GBIF IPT use and data preparation.

## References

- Avaliani R (1963) Data on distribution of some bats in Georgia. Proceedings of the Georgian Academy of Sciences 30 (1): 53-54. [In Georgian].
- Avaliani R (1969a) On the studies of Khulo district (mountain Adjara) Mammals. Proceedings of Simon Janashia State Museum of Georgia 22-23: 233-248. [In Georgian].
- Avaliani R (1969b) Materials to study mammals of high mountain fauna of West Georgia (Tsageri District). Proceedings of Simon Janashia State Museum of Georgia 22-23: 222-232. [In Georgian].
- Avaliani R (1970) Materials to study small mammals of West Georgia (Sachkhere district). Proceedings of Simon Janashia State Museum of Georgia 26-27: 153-168. [In Georgian].
- Avaliani R (1973) Materials to study small mammals of West Georgia (Zestaponi district). Proceedings of Simon Janashia State Museum of Georgia 24-25: 159-177. [In Georgian].
- Avaliani R (1976) On studies of small mammals (Insectivorous, Bats and Rodents) of Borjomi District. Proceedings of Simon Janashia State Museum of Georgia 29: 277-291. [In Georgian].
- Beruchashvili N (2000) Landscape diversity of Georgia and geographical analysis of landscape diversity of the world. In: Beruchashvili N, Kushlin A, Zazanashvili N (Eds) Biological and landscape diversity of Georgia. WWF, Tbilisi, 221-250 pp.
- Bukhnikashvili A, Kandaurov A (2002) The annotated list of mammals of Georgia. Proceedings of the Institute of Zoology 21: 319-340.
- Bukhnikashvili A (2004) On cadastre of small mammals (Insectivora, Chiroptera, Lagomorpha, Rodentia) of Georgia. Universal, Tbilisi, 132 pp. [ISBN 99928-0-8667]
- Bukhnikashvili A, Kandaurov A, Natradze I (2004) Records of Bats in Georgia over the last 140 years. Plecotus et al. 7: 41-57. [In Russian].
- Bukhnikashvili A, Kandaurov A, Natradze I (2008) Bats conservation action plan for Georgia. Universal, Tbilisi, 102 pp. [In Georgian].

- Bukhnikashvili A, Gazaryan S, Kandaurov A, Natradze I, Rakhmatulina I, Yavruyan E (2009) Current Status of Chiroptera Conservation in the Caucasus. In: Zazanashvili N, Mallon D (Eds) Status and Protection of Globally Threatened Species in the Caucasus. Tbilisi, 232 pp. [ISBN 978-9941-0-2203-6].
- Chkhikvishvili I (1926) Vertebrate nomenclature of Georgia. Proceedings of the Georgian Academy of Sciences 3: 1-80.
- Çoraman E, Dietz C, Hempel E, Ghazaryan A, Levin E, Presetnik P, Zagmajster M, Mayer F (2019) Reticulate evolutionary history of a Western Palaearctic Bat Complex explained by multiple mtDNA introgressions in secondary contacts. Journal of Biogeography 46 (2): 343-354. <https://doi.org/10.1111/jbi.13509>
- Dietz C, Kiefer A (2016) Bats of Britain and Europe. Bloomsbury USA, 404 pp. [ISBN 978-1-4729-3576-2]
- Gazaryan S, Bukhnikashvili A (2005) Preliminary data on the status of the Leisler's bat (*Nyctalus leisleri*) in the Caucasus. Nyctalus (N.F) Berlin 10 (3-4): 261-266.
- Gazaryan S, Bukhnikashvili A, Kandaurov A, Natradze I (2008) New status of the *Myotis daubentonii* in the Caucasus. Abstracts of the 11th European Bat Research Symposium. 11th European Bat Research Symposium, Cluj-Napoca, Romania, 18 - 22 August. Cluj-Napoca, 50 pp.
- Gazaryan S, Natradze I, Bukhnikashvili A (2017) First record of *Tadarida teniotis* in Georgia. In: Hudson A, Lina P (Eds) 14th European Bat Research Symposium - EBRS 2017 Abstract book. 14th European Bat Research Symposium, Donostia, The Basque Country, 1-5 August 2017. Donostia, 219 pp. [ISBN 978-84-697-4575-5].
- GBIF.org (2023) GBIF Home Page. <https://www.gbif.org/>. Accessed on: 2023-1-12.
- Imnadze T, Natradze I, Zhgenti E, Malania L, Abazashvili N, Sidamonidze K, Khmaladze E, Zakalashvili M, Imnadze P, Arner R, Motin V, Kosoy M (2020) Identification of a novel *Yersinia enterocolitica* strain from bats in association with a bat die-off that occurred in Georgia (Caucasus). Microorganisms 8 (7). <https://doi.org/10.3390/microorganisms8071000>
- Ivaniitsky A (2002) On the fauna of bats (Chiroptera) of Abkhazia. Plecotus et al. 5: 57-62. [In Russian].
- Ivaniitsky A (2010) New data on the bat fauna (Chiroptera) of Abkhazia. Plecotus et al. 13: 59-68.
- Ivaniitsky A (2017) Ecological and faunal characteristics of bats (Chiroptera) of Abkhazia and adjacent territories (Thesis for the degree of candidate of biological sciences. Institute of Ecology of the Volga Basin, Russian Academy of Sciences, Togliatti, Russia, 222 pp.
- Ivaniitsky A (2018) Bats (chiroptera) of Abkhazia and adjacent territories (fauna, ecology, zoogeography, protection). ARIAL1-153. [In Russian].
- Janashvili A (1953) The key of the mammals of Georgia. Stalin Publishing house of Tbilisi State University. Tbilisi1-216. [In Georgian].
- Janashvili A (1963) The animal world of Georgia. Vertebrata. Metsniereba, Tbilisi, 460 pp. [In Georgian].
- Kolenati F (1860) Monographie der Europaischen Chiroptera. Druck von Rudolf Rohrer Erben, Brünn.
- Kruskop S, Solovyeva E (2020) Validating the relationships: which species of *Myotis* "*nattereri*" group (Chiroptera: Vespertilionidae) actually inhabits the Caucasus. Mammalia 85 (1): 90-99. <https://doi.org/10.1515/mammalia-2019-0146>

- Kuzyakin A (1950) Bats. Sovetskaya Nauka, Moscow, 443 pp. [In Russian].
- Matsaberidze G (1961) On studies of bats helminthofauna in Georgia. 12 Scientific Conference for aspirant and post-graduate students and young researchers, Tbilisi. Tbilisi State University, Tbilisi, 134-139 pp. [In Georgian].
- Matsaberidze G, Khotenovsky I (1966) New trematode *Ophiosacculus eptesicus* species nov from bats of Eastern Georgia. Materials for the fauna of Georgia. Metsniereba, Tbilisi, 192 pp.
- Matsaberidze G, Khotenovskii I (1967) On the fauna of trematoda of chiropterans of Georgia. Animals and Plants Helminthofauna of Georgia 1 (1): 83-94. [In Georgian].
- Matsaberidze G (1976) Helminths of micromammalians of Georgia. Metsniereba, 234 pp.
- Mittermeier R, Turner W, Larsen F, Brooks T, Larsen C (2011) Global biodiversity conservation: The critical role of hotspots. In: Zachos FE, Hubel JC (Eds) Biodiversity hotspots: Distribution and protection of conservation priority areas. Springer, Berlin, 546 pp. [https://doi.org/10.1007/978-3-642-20992-5\\_1](https://doi.org/10.1007/978-3-642-20992-5_1)
- Mumladze L, Japoshvili B, Anderson E (2020) Faunal biodiversity research in the Republic of Georgia: a short review of trends, gaps, and needs in the Caucasus biodiversity hotspot. *Biologia* 75 (9): 1385-1397. <https://doi.org/10.2478/s11756-019-00398-6>
- Myers N, Mittermeier RA, Mittermeier CG, Fonseca GA, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403: 853-858. <https://doi.org/10.1038/35002501>
- Natradze I, Bukhnikashvili A, Sheklashvili G, Mumladze L (2023) Bats of Georgia. Institute of Zoology, Ilia State University. Occurrence dataset. GBIF. Release date: 2023-2-23. URL: <https://doi.org/10.15468/c2spsh>
- Nordmann A (1840) Voyage dans la Russie méridionale et la Crimée, par la Hongrie, la Valachie et la Moldavie, exécuté en 1837, sous la Direction de M. Anatole de Demidoff. Observations sur la Faune pontique. III. Ernest Bourdin et Comp., Paris, 756 pp.
- Noss R, Platt W, Sorrie B, Weakley A, Means DB, Costanza J, Peet R (2014) How global biodiversity hotspots may go unrecognized: lessons from the North American Coastal Plain. *Diversity and Distributions* 21 (2): 236-244. <https://doi.org/10.1111/ddi.12278>
- Ognev S (1928) Mammals of the Eastern Europe and Northern Asia: Insectivores and Bats. State Publishin House, Moscow, 351 pp. [In Russian].
- Papava A (1949) To distribution and to way of life of bats in Georgia. *Bulletin MOIP* 54 (3): 39-41. [In Gorgian].
- Papava A (1953) The key of the mammals of Georgia. Science-Methodical Cabinet, Tbilisi, 139 pp. [In Georgian].
- Papava A (1960) The key of the ammals of Georgia. Tsodna, Tbilisi, 160 pp. [In Georgian].
- Perov M (1975) Bats of Georgia. Fauna and its protection in the republics of Transcaucasia. Academy of Sciences of the Armenian SSR, Yerevan [In Russian].
- Perov M (1980) To study on bats of Georgia. *Terriological issues, "Nauka" Bats (Chiroptera)*: 59-63. [In Russian].
- Perov M (1983) Concerning the protection of bats. *Problems of ecology and environmental protection*. Tbilisi: 213-221.
- Radde G (1899) The collections of the Caucasus Museum. *Caucasian Museum*, Tbilisi, 520 pp. [In Russian].
- Satunin K (1903) Review of the study of mammals of the Caucasian region. *Proceedings of the Caucasian Department of Russian Geographical Society* 24 (2): 1-63. [In Russian].



- Satunin K (1908) The first addition to the list mammal of Caucasian Region. Proceedings of the Caucasian Department of Russian Geographical Society 26 (4): 1-30. [In Russian].
- Satunin K (1912) Cave bats of Abkhazia. Proceedings of the Caucasian Department of Russian Geographical Society 21 (1): 47-48. [In Russian].
- Satunin K (1915) Mammals of Caucasian Region (Chiroptera, Insectivora, Carnivora). Letters of the Caucasian Museum 1 (1): 1-410. [In Russian].
- Shidlovsky M (2013) Insectivora and Chiroptera (a faunistic overview). Lasha Kvichia private publishing, Tbilisi, 61 pp.
- Šrámek J, Gvoždík V, Benda P (2012) Hidden diversity in bent-winged bats (Chiroptera: Minopteridae) of the Western Palearctic and adjacent regions: implications for taxonomy. Zoological Journal of the Linnean Society 167 (1): 165-190. <https://doi.org/10.1111/j.1096-3642.2012.00870.x>
- Strelkov P (1983) Localities of records of *Myotis brandtii* Eversmann, 1845 and *Myotis mystacinus* Kuhl, 1819 (Chiroptera, Vespertilionidae), data from museums of USSR. Proceedings of the Institute of Zoology of the Academy of Sciences of the USSR 119: 38-42. [In Russian].
- Strelkov P (1988) Brown (*Plecotus auritus*) and grey (*P. austriacus*) bats (Chiroptera, Vespertilionidae) in the USSR. Zoologicheskyy Zhurnal 67 (1): 90-101. [In Russian].
- Urushadze L, Babuadze G, Shi M, Escobar L, Mauldin M, Natradeze I, Machablishvili A, Kutateladze T, Imnadze P, Nakazawa Y, Velasco-Villa A (2021) A cross sectional sampling reveals novel coronaviruses in bat populations of Georgia. Viruses 14 (1). <https://doi.org/10.3390/v14010072>
- Verestchagin N (1959) The Mammals of the Caucasus. History of Formation of the Fauna. AS USSR, Moscow, 703 pp.
- Yavruyan E, Rakhmatulina I, Bukhnikashvili A, Kandaurov A, Ntradze I, Gazaryan S (2008) Bats conservation action plan for the Caucasus. Universal, Tbilisi, 87 pp.




Figure 1.  
Bat records in Georgia from 1835 through to 2023.

Table 1.

Number of records for each species.

#	Species	Total Number of records	Literature & Museum data	Our published & unpublished data	Unpublished data
	<b>Chiroptera</b>				
	<b>Rhinolophidae</b>				
	<b>Rhinolophus</b>				
1	<i>Rhinolophus ferrumequinum</i> (Schreber, 1774)	185	58	127	84
2	<i>Rhinolophus hipposideros</i> (André, 1797)	167	83	84	53
3	<i>Rhinolophus euryale</i> Blasius, 1853	51	20	31	23
4	<i>Rhinolophus blasii</i> Peters, 1866	6	0	6	5
5	<i>Rhinolophus mehelyi</i> Matschie, 1901	5	5	0	0
	<b>Vespertilionidae</b>				
	<b>Myotis</b>				
6	<i>Myotis blythii</i> (Tomes, 1857)	120	34	86	61
7	<i>Myotis bechsteinii</i> Kuhl, 1817	25	4	21	21
8	<i>Myotis daubentonii</i> (Kuhl, 1817)	41	6	35	33
9	<i>Myotis natereri</i> (Kuhl, 1817)	65	12	53	51
10	<i>Myotis emarginatus</i> (Geoffroy, 1806)	72	15	57	35
11	<i>Myotis alcathoe</i> von Helversen & Heller, 2001	25	1	24	24
12	<i>Myotis brandtii</i> (Eversmann, 1845)	28	3	25	24
13	<i>Myotis davidii</i> (Peters, 1869)	3	0	3	3
14	<i>Myotis mystacinus</i> (Kuhl, 1817)	89	33	56	54
	<b>Nyctalus</b>				
15	<i>Nyctalus noctula</i> (Schreber, 1774)	89	16	73	72
16	<i>Nyctalus leisleri</i> (Kuhl, 1817)	87	10	77	68
17	<i>Nyctalus lasiopterus</i> (Schreber, 1780)	23	7	16	8
	<b>Eptesicus</b>				
18	<i>Eptesicus nilssonii</i> (von Keyserling & Blasius, 1839)	4	1	3	2
19	<i>Eptesicus serotinus</i> (Schreber, 1774)	160	48	112	105
	<b>Pipistrellus</b>				

20	<i>Pipistrellus pipistrellus</i> (Schreber, 1774)	289	64	225	218
21	<i>Pipistrellus pygmaeus</i> (Leach, 1825)	74	0	74	70
22	<i>Pipistrellus kuhlii</i> (Kuhl, 1817)	83	11	72	70
23	<i>Pipistrellus nathusii</i> (von Keyserling & Blasius, 1839)	17	15	2	2
	<b>Hypsugo</b>				
24	<i>Hypsugo savii</i> (Bonaparte, 1837)	23	4	19	17
	<b>Barbastella</b>				
25	<i>Barbastella barbastellus</i> (Schreber, 1774)	63	8	55	49
	<b>Plecotus</b>				
26	<i>Plecotus auritus</i> (Linnaeus, 1758)	60	29	31	29
27	<i>Plecotus macrobularis</i> Kuzyakin, 1965	14	5	9	8
	<b>Vespertilio</b>				
28	<i>Vespertilio murinus</i> Linnaeus, 1758	38	15	23	23
	<b>Miniopteridae</b>				
	<b>Miniopterus</b>				
29	<i>Miniopterus schreibersii</i> (Kuhl, 1817)	65	35	30	18
	<b>Molossidae</b>				
	<b>Tadarida</b>				
30	<i>Tadarida teniotis</i> (Rafinesque, 1814)	14	1	13	13