

Chemo-organo-autotrophic degradation of aromatic hydrocarbons indicates a new type of bacterial metabolism

Rainer U. Meckenstock[‡], Isabelle Heker[‡], Christian Seitz[§], Lisa Voskuhl[‡], Wolfgang Eisenreich[§]

[‡] University of Duisburg-Essen, Essen, Germany

[§] Technical University of Munich, Munich, Germany

Corresponding author: Rainer U. Meckenstock (rainer.meckenstock@uni-due.de)

Abstract

The sulfate-reducing culture N47 can grow with naphthalene and has a complete tricarboxylic acid cycle (TCA) and Wood-Ljungdahl pathway (WLP) while other organisms have only either of them. Here, we wanted to elucidate why N47 has two complete pathways potentially able to oxidize acetyl-CoA. Enzyme activities were measured in cell extracts indicating a fully functional rTCA and WLP. To elucidate the carbon flux through the pathways, cells were grown with ¹³C-labeled naphthalene or ¹³C-bicarbonate buffer. Amino acids and fatty acids were analysed for position specific ¹³C-incorporation with GC-MS, which indicated that in catabolism, acetyl-CoA from naphthalene was fully oxidized to CO₂ via the WLP. Acetyl-CoA for anabolism of amino acids, fatty acids and carbohydrates was surprisingly not coming from the substrate naphthalene but is generated *de novo* by CO₂-fixation, making N47 a chemoorganoautotrophic microorganism. This indicates that chemoorganoauto-trophy can also occur with complex substrates but probably requires a complete WLP and rTCA in anaerobic microorganisms.

Keywords

Central metabolism, chemo-organo-auto trophy, TCA-cycle, Wood-Ljungdahl pathway, anaerobic hydrocarbon degradation, sulfate-reduction

Presenting author

Rainer U. Meckenstock

Presented at

ISEB-ISSM 2023 joint meeting

Hosting institution

University of Duisburg-Essen, Environmental Microbiology and Biotechnology

Conflicts of interest

The authors have declared that no competing interests exist.