

The Shiny Cowbird, *Molothrus bonariensis* (Gmelin, 1789) (Aves: Icteridae), at 2,800 m asl in Quito, Ecuador

Verónica Crespo-Pérez[‡], C. Miguel Pinto^{§,¶}, Juan Manuel Carrión[¶], Rubén D. Jarrín-E[‡], Cristian Poveda[‡], Tjitte de Vries[‡]

[‡] Pontificia Universidad Católica del Ecuador, Escuela de Ciencias Biológicas, Av. 12 de Octubre y Roca, Quito, Ecuador

[§] Instituto de Ciencias Biológicas, Escuela Politécnica Nacional, Quito, Ecuador

[¶] Smithsonian Institution, National Museum of Natural History, Division of Mammals, 1000 Constitution Ave NW, Washington DC, United States of America

[¶] Fundación Zoológica del Ecuador, Quito, Ecuador

Corresponding author:

Academic editor: Christopher Heckscher

Abstract

Background

The Shiny Cowbird, *Molothrus bonariensis* Gmelin, 1789, is a brood parasite of hundreds of small-bodied birds that is native to South American lowlands. Within the last 100 years this species has been expanding its range throughout the Caribbean, towards North America, but has rarely been seen above 2,000 m asl.

New information

Here, we present records of Shiny Cowbirds in Quito, a city located 2,800 m above sea level that harbors a bird community typical of the Andean valleys. We found two juvenile individuals parasitizing two different pairs of Rufous-collared Sparrow (*Zonotrichia capensis* Müller, 1776). This report constitutes an altitudinal range expansion of reproductive populations of ca. 500m, which may have been prompted by anthropogenic disturbance.

Keywords

Brood parasite, Climate change, Habitat fragmentation, *Molothrus bonariensis*, Rufous-collared Sparrow, Shiny Cowbird, university campus, *Zonotrichia capensis*

Introduction

The Shiny Cowbird, *Molothrus bonariensis* Gmelin, 1789, is a brood parasite of hundreds of small-bodied birds, and the Rufous-collared Sparrow, *Zonotrichia capensis* Muller, 1776 is one of its main hosts (Mason 1986, Lowther 2013). It is native to South American lowlands (Bird Life International and NatureServe 2012), but within the last 100 years has been expanding its distribution throughout the Caribbean, towards North America (Post et al. 1993) with some isolated records from Costa Rica and Mexico (Kluza 1998, Sandoval et al. 2010) (Fig. 1). The species occupies lowlands and it has rarely been seen above 2,000 m asl (Hilty and Brown 1986, Lowther 2011); however, there are notorious records in Bolivia and Peru above 3,000 m asl (Miller 1917, Fjeldsa and Krabbe 1990, Jaramillo and Burke 1999, Balderrama 2006, Schulenberg et al. 2007). Particularly in Ecuador, it has been mostly recorded between 900 and 1,000 m asl, but also regularly higher to ca. 1,400 to 1,600 m asl (Ridgely and Greenfield 2001, Ridgely and Greenfield 2006). There have been isolated records in the highlands (as high as 2,700 m asl) but there are no reports of these birds breeding at such altitudes. There also are records in the Pichincha province (up to 2,700 m asl) (Fjeldsa and Krabbe 1990, Ridgely and Greenfield 2001), and vocal recordings available from the southern city of Loja (2,100 m asl) and from lake San Pablo in the north of the country (2,700 m asl) in the sound library Xeno-Canto (www.xeno-canto.org) (Fig. 2). In the year 2000 the Shiny Cowbird was first spotted in the inter Andean valley of Cumbayá, at 2,300 m asl, 7 km to the east of Quito, Ecuador (Carrion 2001, Carrion pers. obs.). These birds are now regular inhabitants of that area and may be observed there year round, sometimes in groups of more than 50 individuals (Carrion pers. obs.).

Here, we present records of juvenile Shiny Cowbirds at 2,800m asl in Quito—Ecuador's capital city located in the inter Andean valley, with a population of over 1.6 million people. The bird community of the city is typical of the inter Andean valleys (Carrion 2001), with many species of hummingbirds and other species like the Eared Dove, *Zenaida auriculata* Des Murs, 1847, the Golden Grosbeak, *Pheucticus chrysogaster* Lesson, 1832, the Great Thrush, *Turdus fuscater* Lafresnaye & d'Orbigny, 1837, the Vermilion Flycatcher, *Pyrocephalus rubinus* Boddaert, 1783, and the Black Flowerpiercer, *Diglossa humeralis* Fraser, 1840. The new record presented in this contribution highlights the dispersal capabilities of *M. bonariensis*, revealing that this species is able to colonize a city at high altitude.

Materials and methods

In the campus of the Pontificia Universidad Católica del Ecuador (PUCE), Quito, Ecuador (0°12'40" S, 78°29'28" W), we observed two juveniles of *M. bonariensis* interacting with *Zonotrichia capensis* Müller, 1776, between 11 April and 15 May 2015. One of the juveniles of *M. bonariensis* was larger and more developed than the other, and each was associated with a different *Z. capensis* pair. We observed the juveniles of *M. bonariensis*

displaying food begging behaviors to adult *Z. capensis* (which included chasing the sparrows on the ground), and vocalizing intensely on bushes and tree branches. The adults of *Z. capensis* were observed feeding the Shiny Cowbirds on several occasions (Fig. 3a, b) but they were not observed feeding fledglings of their own species.

Taxon treatment

Molothrus bonariensis Gmelin, 1789

Material

- a. vernacularName: Shiny Cowbird, Vaquero brillante; genus: *Molothrus*; specificEpithet: *bonariensis*; scientificNameAuthorship: Gmelin, 1789; continent: South America; country: ECUADOR; stateProvince: Pichincha; municipality: Quito; verbatimElevation: 2800 m; verbatimCoordinateSystem: decimal degrees; decimalLatitude: -0.2111111; decimalLongitude: -78.49111; geodeticDatum: WGS84; eventDate: 2015-04-11; habitat: urban university campus; individualCount: 2; lifeStage: juvenile; recordedBy: Verónica Crespo-Pérez; C. Miguel Pinto; identifiedBy: Tjitte de Vries; occurrenceID: CEE0C3DD-CB42-5BBE-9026-6E9CE3C79AE0

Diagnosis

The observed individuals of Shiny Cowbird matched the morphological and behavioral characteristics of the species (Fig. 3a, b, c). The most remarkable characteristic was the dull grayish coloration on the dorsum, but paler on the venter and throat, and the greyish white eyebrows (Hilty and Brown 1986, Restall et al. 2006). For Ecuador, three subspecies have been documented: *M. b. aequatorialis* in the north, *M. b. occidentalis* in the south, and *M. b. riparius* in the east (Restall et al. 2006). The juveniles that we observed have a clear coloration (Fig. 3a, b) that matches that of the subspecies *M. b. occidentalis*, and not that of *M. b. aequatorialis*—which is darker—as would be expected for the location of Quito in the north of the country. Nevertheless, juvenile coloration might not match that of adults and more studies should be conducted for subspecific assignment perhaps using an integrative taxonomic approach combining molecular, morphological and biogeographic information.

Identification keys

Key to the species of <i>Molothrus</i>		
1	Male and female weight more than 100 g	<i>Molothrus oryzivorus</i>
–	Male and female weight less than 100 g	2

2	Males iridescent black with a brown head	<i>Molothrus ater</i>
–	Males iridescent black without contrasting coloration of head	3
3	Males with red eyes during breeding season	<i>Molothrus aeneus</i>
–	Males with brown eyes during breeding season	4
4	Both sexes are dimorphic in coloration, males have a violet gloss	<i>Molothrus bonarensis</i>
–	Coloration of both sexes is similar, and males are less glossy than <i>M. bonarensis</i>	<i>Molothrus rufoaxillaris</i>

Discussion

In recent years the Shiny Cowbird has colonized the Caribbean and the east coast of the United States (Fig. 1). Particularly in Ecuador, this species seemed to be restricted to below ca. 2,000m asl (Hilty and Brown 1986, Ridgely and Greenfield 2001, Restall et al. 2006, Lowther 2011), with isolated higher altitude records (Fjeldsa and Krabbe 1990, Ridgely and Greenfield 2001, Xeno-canto Foundation 2012) (Fig. 2), and had never before been reported in the high-altitude city of Quito. Surely, this inability to colonize the city was puzzling since it has a very long history of habitat alteration, and one of the Shiny Cowbird's most common hosts, *Z. capensis*, does well in high elevations and is abundant in Quito (Carrion 2001, Ridgely and Greenfield 2001). Even though there is strong evidence suggesting habitat fragmentation is the main driver of *M. bonariensis* range expansions (Arendt and Vargas Mora 1984, Post et al. 1993, Hansen et al. 2005) we believe that in this case, climate change might be playing an important role as well. Recent climate warming may have relaxed climatic conditions allowing this species to expand towards higher elevations. There is, in fact, evidence of temperature increases in the Ecuadorian Andes (e.g., Vuille et al. 2008), which may be related to range expansions such as the one presented in this contribution. Climate induced distributional shifts have been fairly well documented especially in species with high dispersal ability like birds, insects and marine invertebrates (Parmesan 2006). Several authors have reported range expansions of species towards higher altitudes (Paulson 2001, Parmesan 2006, Moritz et al. 2008, Chen et al. 2009, Chen et al. 2011). Along the Andes there are several examples of upslope expansion of distributions of plants and birds (Weng et al. 2007, Feeley et al. 2010, Forero-Medina et al. 2011, Morueta-Holme et al. 2015). Also, according to recent evidence, tropical species respond more strongly than temperate ones to warming temperatures, with range shifts that match local temperature increase more closely than in temperate-zone montane species (Freeman and Freeman 2014).

Avian brood parasites, including cuckoos and cowbirds, often reduce the reproductive success of their hosts (Tuero et al. 2007). Costs associated with such parasitism promote the evolution of host antiparasitic defenses – such as responding aggressively towards cowbirds to prevent them from gaining access to the nest, burying or ejecting parasite

eggs, or abandoning parasitized nests – and create a coevolutionary arms race between hosts and parasites (Astie and Reboresda 2005). Nevertheless, a number of hosts, even commonly parasitized hosts like the Rufous-collared sparrow, have not evolved antiparasite defenses, despite the associated fitness costs, due probably to morphological or ecological constraints (Carro and Fernández 2013). In addition, in the case of invasive parasites, native hosts may be naïve to the parasite and lack defenses or appropriate mechanisms to counteract its negative effects (Taraschewski 2006, Fassbinder-Orth et al. 2013). The level of threat posed by invasive Cowbirds on native bird communities apparently depends on the local species present (i.e., whether they lack defenses against brood parasitism), and is closely related to other human disturbances, mainly habitat fragmentation (Rothstein and Peer 2005, Peer et al. 2013). In the specific case of the Shiny Cowbird, several studies have found negative effects of this parasite, especially for already endangered and/or endemic bird species (Post 2011, Price et al. 2011, Domínguez et al. 2014). In fact, there is evidence that suggests that Shiny Cowbird parasitism, coupled with habitat loss, almost drove the Pale-headed Brush-Finch, *Atlapetes pallidiceps* Sharpe, 1900, a critically endangered endemic to south-central Ecuador, to extinction (Oppel et al. 2004, Krabbe et al. 2010). Therefore, the Shiny Cowbird range expansion presented in this study may be of concern, especially for native, endemic, or endangered species that may be parasitized by this species. More studies are urgently needed to assess the threat posed by this invasive species to native avian fauna in the high Andes in order to guide management or control measures for this invasive brood parasite.

Acknowledgements

We thank the efforts of the Pontificia Universidad Católica del Ecuador to keep green areas in campus where observations of this kind can be conducted. We would also like to thank the Editor Christopher Heckscher, Juan Fernando Freile, and an anonymous reviewer, whose constructive critiques and comments greatly improved the quality of this contribution. We thank Edith Montalvo for introducing us to the Xeno-Canto website. CMP thanks Kristofer Helgen, Meri Price and Don Wilson for facilitating his visit to Ecuador in April 2015.

Author contributions

VCP, CMP, RDJ, CP, and TdV collected the data; RDJ and VCP took the photographs; VCP drew the maps; VCP and CMP wrote the text with contributions from JMC and TdV; VCP, CMP, JMC, RDJ, CP and TdV read and agree with the content of the manuscript.

References

- Arendt WJ, Vargas Mora TA (1984) Range Expansion of the Shiny Cowbird in the Dominican Republic. *Journal of Field Ornithology* 55 (1): 104-107. URL: <http://www.jstor.org/stable/4512862>
- Astie A, Reboreda J (2005) Creamy-bellied Thrush defenses against Shiny Cowbird brood parasitism. *The Condor* 107 (4): 788-796. <https://doi.org/10.1650/7733.1>
- Balderrama JA (2006) Diversidad, endemismo y conservación de la ornitofauna del Parque Nacional Tunari (Cochabamba, Bolivia). *Ecología de Bolivia* 41 (2): 149-170. [In Spanish]. URL: http://www.scielo.org.bo/scielo.php?script=sci_arttext&pid=S1605-25282006001000006
- Bird Life International and NatureServe (2012) Bird Species Distribution Maps of the World. <http://maps.iucnredlist.org/map.html?id=22724345>. Accessed on: 2015-7-30.
- Carrion JM (2001) Aves de Quito, Retratos y encuentros. SIMBIOE, Quito, 135 pp.
- Carro M, Fernández G (2013) Can nest predation explain the lack of defenses against cowbird brood parasitism in the Rufous-collared Sparrow (*Zonotrichia capensis*)? *The Auk* 130 (3): 408-416. <https://doi.org/10.1525/auk.2013.12164>
- Chen I, Hill J, Shiu H, Holloway J, Benedick S, Chey VK, Barlow H, Thomas C (2011) Asymmetric boundary shifts of tropical montane Lepidoptera over four decades of climate warming. *Global Ecology and Biogeography* 20 (1): 34-45. <https://doi.org/10.1111/j.1466-8238.2010.00594.x>
- Chen I-C, Shiu H-J, Benedick S, Holloway JD, Chey VK, Barlow HS, Hill JK, Thomas CD (2009) Elevation increases in moth assemblages over 42 years on a tropical mountain. *Proceedings of the National Academy of Sciences* 106 (5): 1479-1483. <https://doi.org/10.1073/pnas.0809320106>
- Domínguez M, Reboreda JA, Mahler B (2014) Impact of Shiny Cowbird and botfly parasitism on the reproductive success of the globally endangered Yellow Cardinal *Gubernatrix cristata*. *Bird Conservation International* 25 (3): 294-305. <https://doi.org/10.1017/s095927091400015x>
- Fassbinder-Orth C, Barak V, Brown C (2013) Immune Responses of a Native and an Invasive Bird to Buggy Creek Virus (Togaviridae: Alphavirus) and Its Arthropod Vector, the Swallow Bug (*Oeciacus vicarius*). *PLoS ONE* 8 (2): e58045. <https://doi.org/10.1371/journal.pone.0058045>
- Feeley K, Silman M, Bush M, Farfan W, Cabrera KG, Malhi Y, Meir P, Revilla NS, MNRQ, Saatchi S (2010) Upslope migration of Andean trees. *Journal of Biogeography* 38 (4): 783-791. <https://doi.org/10.1111/j.1365-2699.2010.02444.x>
- Fjeldsa J, Krabbe N (1990) *Birds of the High Andes: A Manual to the Birds of the Temperate Zone of the Andes and Patagonia, South America*. Apollo Booksellers, Copenhagen, 880 pp.
- Forero-Medina G, Terborgh J, Socolar SJ, Pimm S (2011) Elevational Ranges of Birds on a Tropical Montane Gradient Lag behind Warming Temperatures. *PLoS ONE* 6 (12): e28535. <https://doi.org/10.1371/journal.pone.0028535>
- Freeman BG, Freeman AMC (2014) Rapid upslope shifts in New Guinean birds illustrate strong distributional responses of tropical montane species to global warming.

- Proceedings of the National Academy of Sciences 111 (12): 4490-4494. <https://doi.org/10.1073/pnas.1318190111>
- Hansen A, Knight R, Marzluff J, Powell S, Brown K, Gude P, Jones K (2005) Effects of exurban development on biodiversity: patterns, mechanisms, and research needs. *Ecological Applications* 15 (6): 1893-1905. <https://doi.org/10.1890/05-5221>
 - Hilty S, Brown W (1986) A guide to the birds of Colombia. Princeton University Press, New Jersey, 996 pp.
 - Jaramillo A, Burke P (1999) New world blackbirds: the icterids. Christopher Helm Publishers Ltd, London, 432 pp.
 - Kluz DA (1998) First record of Shiny Cowbird (*Molothrus bonariensis*) in Yucatán, Mexico. *Wilson Bulletin-Morgantown Then Columbus-Ornithological Bulletin* 110: 429-430.
 - Krabbe N, Juiña M, Sornoza AF (2010) Marked population increase in Pale-headed Brush-finch *Atlapetes pallidiceps* in response to cowbird control. *Journal of Ornithology* 152 (2): 219-222. <https://doi.org/10.1007/s10336-010-0567-z>
 - Lowther PE (2011) Shiny Cowbird (*Molothrus bonariensis*). *Neotropical Birds Online*. http://neotropical.birds.cornell.edu/portal/species/overview?p_p_spp=672716. Accessed on: 2015-7-30.
 - Lowther PE (2013) Lists of victims and hosts of the parasitic cowbirds, Version 26 August 2013. http://fieldmuseum.org/sites/default/files/Molothrus_hosts-26aug2013.pdf. Accessed on: 2015-7-30.
 - Mason P (1986) Brood parasitism in a host generalist, the Shiny Cowbird: I. The quality of different species as hosts. *The Auk* 103: 52-60. URL: <http://www.jstor.org/stable/4086963>
 - Miller LE (1917) Field notes on *Molothrus bonariensis* and *M. badius*. *Bulletin of the American Museum of Natural History* 37: 579-592.
 - Moritz C, Patton JL, Conroy CJ, Parra JL, White GC, Beissinger SR (2008) Impact of a century of climate change on small-mammal communities in Yosemite National Park, USA. *Science* 322 (5899): 261-264. <https://doi.org/10.1126/science.1163428>
 - Morueta-Holme N, Engemann K, Sandoval-Acuña P, Jonas J, Segnitz RM, Svenning J (2015) Strong upslope shifts in Chimborazo's vegetation over two centuries since Humboldt. *Proceedings of the National Academy of Sciences* 112 (41): 12741-12745. <https://doi.org/10.1073/pnas.1509938112>
 - Opiel S, Schaefer HM, Schmidt V, Schroder B (2004) Cowbird parasitism of Pale-headed Brush-finch *Atlapetes pallidiceps*: implications for conservation and management. *Bird Conservation International* 14 (2): 63-75. <https://doi.org/10.1017/s0959270904000103>
 - Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics* 37 (1): 637-669. <https://doi.org/10.1146/annurev.ecolsys.37.091305.110100>
 - Paulson D (2001) Recent Odonata records from southern Florida - effects of global warming? *International Journal of Odonatology* 4 (1): 57-69. <https://doi.org/10.1080/13887890.2001.9748159>
 - Peer B, Rivers J, Rothstein S (2013) Cowbirds, conservation, and coevolution: potential misconceptions and directions for future research. *Chinese Birds* 4 (1): 15-30. <https://doi.org/10.5122/cbirds.2013.0009>
 - Post W (2011) Yellow-shouldered Blackbird (*Agelaius xanthomus*). http://neotropical.birds.cornell.edu/portal/species/overview?p_p_spp=666636. Accessed on: 2016-4-08.

- Post W, Cruz A, McNair DB (1993) The North American Invasion Pattern of the Shiny Cowbird. *Journal of Field Ornithology* 64: 32-41. URL: <http://www.jstor.org/stable/4513767>
- Price M, Lee V, Hayes W (2011) Population status, habitat dependence, and reproductive ecology of Bahama Orioles: a critically endangered synanthropic species. *Journal of Field Ornithology* 82 (4): 366-378. <https://doi.org/10.1111/j.1557-9263.2011.00340.x>
- Restall R, Rodner C, Lentino M (2006) *Birds of northern South America, vol. 2: an identification guide*. Christopher Helm Publishers Ltd, London, 656 pp.
- Ridgely R, Greenfield PJ (2001) *The Birds of Ecuador: Status, Distribution, and Taxonomy*. Ithaca, Cornell University Press, 880 pp.
- Ridgely R, Greenfield PJ (2006) *Aves del Ecuador: Guía de campo*. Jocotoco, Quito, 812 pp.
- Rothstein S, Peer B (2005) Conservation Solutions for Threatened and Endangered Cowbird (*Molothrus* spp.) Hosts: Separating Fact from Fiction. *Ornithological Monographs* 57: 98-114. <https://doi.org/10.2307/40166817>
- Sandoval L, Sánchez C, Biamonte E, Zook JR, Sánchez JE, Martínez D, Loth D, O'Donahoe J (2010) Recent records of new and rare bird species in Costa Rica. *Bulletin of the British Ornithologists' Club* 130: 237-245.
- Schulenberg TS, Stotz DF, Lane DF, O'Neill JP, Parker TA (2007) *Birds of Peru*. Christopher Helm Publishers Ltd, London, 659 pp.
- Taraschewski H (2006) Hosts and parasites as aliens. *Journal of Helminthology* 80 (2): 99-128. <https://doi.org/10.1079/joh2006364>
- Tuero DT, Fiorini VD, Reboreda JC (2007) Effects of Shiny Cowbird *Molothrus bonariensis* parasitism on different components of House Wren *Troglodytes aedon* reproductive success. *Ibis* 149 (3): 521-529. <https://doi.org/10.1111/j.1474-919x.2007.00676.x>
- Vuille M, Francou B, Wagnon P, Juen I, Kaser G, Mark B, Bradley R (2008) Climate change and tropical Andean glaciers: Past, present and future. *Earth-Science Reviews* 89: 79-96. <https://doi.org/10.1016/j.earscirev.2008.04.002>
- Weng C, Hooghiemstra H, Duivenvoorden JF (2007) Response of pollen diversity to the climate-driven altitudinal shift of vegetation in the Colombian Andes. *Philosophical Transactions of the Royal Society B: Biological Sciences* 362 (1478): 253-262. <https://doi.org/10.1098/rstb.2006.1985>
- Xeno-canto Foundation (2012) Xeno-canto: Sharing bird sounds from around the world. <http://www.xeno-canto.org/species/Molothrus-bonariensis>. Accessed on: 2016-4-08.




Figure 1.

Distribution of the Shiny Cowbird in the Americas, including the Caribbean, modified from Post et al. (1993) and BirdLife International and NatureServe (2014). Areas in blue represent the native range, while areas in red represent the invaded range.




Figure 2.

Distribution of the Shiny Cowbird in Ecuador, modified from Ridgely and Greenfield (2006). Areas in blue represent areas below 2000 m asl where the shiny cowbird has been previously reported (Ridgely and Greenfield 2006), black dots represent isolated, higher altitude records (up to 2700 m asl) (Ridgely and Greenfield 2006, Xeno-canto Foundation 2012), and the red dot marks the city of Quito, where we report the presence of the Shiny Cowbird.




Figure 3.

Juvenile individuals of Shiny cowbird, *Molothrus bonariensis*, and foster *Zonotrichia capensis* parent, photographed at the PUCE campus in Quito.

a: Foster *Zonotrichia capensis* parent (left) approaching juvenile Shiny cowbird (right). Photographed with a Nikon D800 camera and a Nikon AF-S NIKKOR 400mm f/2.8G ED VR lens by RDJ.

b: Foster *Zonotrichia capensis* parent (left) feeding juvenile Shiny cowbird (right). Photographed with a Nikon D800 camera and a Nikon AF-S NIKKOR 400mm f/2.8G ED VR lens by RDJ.

c: Juvenile of Shiny Cowbird showing the grey coloration, eye band and paler underparts. Photographed with a cell phone by VCP.