

# Mystery mushroom malingerers: *Megaselia marquezii* Hartop et al. 2015 (Diptera: Phoridae)

Brian V. Brown<sup>‡</sup>, Emily A. Hartop<sup>‡</sup>

<sup>‡</sup> Natural History Museum of Los Angeles County, Los Angeles, United States of America

Corresponding author: Brian V. Brown ([bbrown@nhm.org](mailto:bbrown@nhm.org))

Academic editor: Pierfilippo Cerretti

## Abstract

A mysterious female phorid fly, known for many years to be associated with fungal sporophores ("mushrooms") is identified as *Megaselia marquezii* Hartop et al. 2015. Male and female flies were collected emerging from the fungus *Psathyrella candolleana* (Fr.) Maire, and females were observed swarming over the sporophores.

## Keywords

Natural history, Diptera, fungivore, urban, biodiversity, citizen science

## Introduction

The genus *Megaselia* Rondani 1856 (Diptera: Phoridae) is immense, with over 1000 described species and certainly thousands more unrecognized. In addition to their species-richness, the lifestyles of various species are so diverse and little-studied that any addition to this knowledge is of great significance.

For nearly thirty years, one of us (BVB) has been aware of an unusual species of *Megaselia* whose females were associated with fungal fruiting bodies. These females have an unusual ovipositor that would lead them to key unconvincingly to *Plastophora* Brues 1905 (now mostly subsumed in *Myriophora* Brown 1993) in the key of Borgmeier (1963). Males were unknown, and had not been found on the fungi. The distinctive females (or ones similar to them) were found in eastern Canada, Costa Rica, and Los Angeles, California, USA.

Meanwhile, a large inventory project in Los Angeles has led to an unprecedented knowledge of the urban phorids of this city (Brown and Hartop 2016, Hartop et al. 2015, Hartop et al. 2016a, Hartop et al. 2016b). About 100 species, mostly of *Megaselia*, are known from Los Angeles, but many were new to science and had nothing known of their

lifestyle. Matching a lifestyle with a species previously known only from a name is a significant accomplishment.

## Material and methods

Specimens were collected from the back garden of a commercial bed and breakfast establishment in downtown Los Angeles, near the campus of the University of Southern California (coordinates 34.02 degrees North, 118.17 degrees West). On 20 April 2017, large numbers of fruiting bodies of the fungus *Psathyrella candolleana* (Fr.) Maire were noted in the garden. Adult flies that were flying above or running on the fungi (as in Fig. 1) were collected into 70% ethanol. Specimens were collected from two locations in the garden, one of which yielded teneral specimens that were apparently just emerging from the soil surrounding the fungi.

Several fruiting bodies of the fungi were removed and brought in to the laboratory for further study.

The BioSCAN project (Brown and Hartop 2016) is an ongoing survey of insects in urban Los Angeles, as part of the investigations of the Natural History Museum of Los Angeles County's Urban Nature Research Center. Data from phase I of the BioSCAN project were reviewed for placing the current finding about *Megaselia marquezii* Hartop et al. (2015) in context. Correlation analysis was done using Microsoft EXCEL on the entire data set for the BioSCAN project.

## Results

Flies on fungi were noted in early April, 2017. Specimens of flies were collected 20 April, and fungi were brought to our laboratory on 24 April, 2017. On fully open fungal sporophores ("mushrooms"), we noted large numbers of eggs laid within the gills (Fig. 2). Two adult female flies were observed feeding on the mushrooms (Fig. 3, Fig. 4). One of the females was exposed when we broke open the cap of a mushroom that was still closed, it appeared from her behavior that she may have just oviposited (Fig. 5). Larvae were found feeding on the lower surface of the mushroom cap, deep within the gills (Fig. 6). One large mushroom (ca. 5 cm in radius) was estimated by subsampling to contain at least 500 larvae. One puparium was found in surrounding soil we collected onsite. The soil and mushrooms were kept for several weeks at room temperature in a plastic bag, and larvae were found to have exited the mushrooms to pupate in the soil within a few days.

We observed hundreds of flies around and on the fungi, on nearby leaf litter and on soil. Among the teneral flies emerging from the soil surrounding the fungi were many females and three males of the "mystery" phorid, identified (from the males) as *Megaselia marquezii* Hartop et al. Because only females are attracted to the fungi, the collection of teneral specimens, including males, was vital to this discovery.

Most of the specimens of this fly collected by the survey Malaise traps (1715 in total) were caught at a single site in April, 2015, when a sample contained 459 individual males ( $459/1715 = 27\%$ ).

## Discussion

Among over 42,000 phorid flies collected by BioSCAN project in Los Angeles (Hartop et al. 2015), *Megaselia marquezii* was the sixth most commonly collected species, occurring in all but two sites in our study. Its occurrence is most highly correlated to that of *M. berndseni* (Schmitz, 1919), a species also known to breed in the fungus *Psathyrella candolleana* in Europe (Table 1). Other phorids known to breed in this fungus are noted in Table 1. The most frequently collected fly in the survey, *M. agarici* (Lintner, 1895) (Fig. 7), is also a fungivore.

Although *Megaselia marquezii* was recorded in many of the BioSCAN trap samples, its ranking as the sixth most common species in the survey is largely based on the huge outburst of specimens found in trap number 5 in April 2015 (Figs 7, 8). Based on our observations of the masses of adults and larvae at the site reported herein, however, this surge of individuals could have been caused by a small group of mushrooms emerging near the trap. Caution is needed when interpreting numbers of phorid flies in trap samples when such scattered resources as mushrooms can lead to spikes in numbers of specimens in the BioSCAN traps.

## Conclusions

So far, in spite of our incomplete knowledge of *Megaselia* life histories, we can say that the urban fauna of Los Angeles is numerically dominated by fungus-feeders (Brown and Hartop 2016), including the newly discovered lifestyle of *M. marquezii*. Of the 42,480 specimens identified, 18,918 (45%) are now known to be of species that are fungivores. Possibly, the widespread irrigation of lawns allows fungal growth that supports an abundant fungivore community, but our ignorance of the fauna of the surrounding natural areas makes such statements highly speculative.

## Acknowledgements

We thank Patsy Carter and Lisa Carter-Davis of "The Inn at 657" for alerting us to the mushrooms that attracted so many phorid flies, and for allowing us to do research on their property. The fungus was identified by Florence Nishida. The BioSCAN project and the Urban Nature Research Center were supported by the Seaver Foundation, the Natural History Museum of Los Angeles County, and donations by Sheila Brennan, Wally and Carol Marks, Susan and Steve Matt, Leslie Mitchner, Karen and Bill Timberlake, Tina and Larry Wiener, and Mary Simun and Amazon Beard.

## Conflicts of interest

## References

- Borgmeier T (1963) Revision of the North American Phorid Flies. Part I. The Phorinae, Aenigmatiinae and Metopininae, except *Megaselia* (Diptera, Phoridae). *Studia Entomologica* 6: 1-256.
- Brown BV, Hartop EA (2016) Big data from tiny flies: patterns revealed from over 42,000 phorid flies (Insecta: Diptera: Phoridae) collected over one year in Los Angeles, California, USA. *Urban Ecosystems* <https://doi.org/10.1007/s11252-016-0612-7>
- Disney RH (1989) Six new species of *Megaselia* reared from fungi in Japan. *Acta Entomologica Bohemoslovaca* 86: 368-380.
- Disney RH, Evans RE (1979) Further records of Phoridae (Dipt.) reared from fungi. *Entomologist's Monthly Magazine* 114: 166-166.
- Disney RH, Evans RE (1990) Some new records of fungus-breeding Phoridae (Diptera). *Dipterist's Digest* 7: 20-23.
- Disney RH, Evans RE (1999) New records of Phoridae (Diptera) reared from fungi. *Entomologist's Record* 111: 235-238.
- Eisfelder I (1956) Die häufigsten Pilzbewohner (Fliegen als Pilzerzehrer). *Zeitschrift für Pilzkunde* 22: 108-117.
- Hartop EA, Brown BV, Disney RHL (2015) Opportunity in our Ignorance: Urban Biodiversity Study Reveals 30 New Species and One New Nearctic Record for *Megaselia* (Diptera: Phoridae) in Los Angeles (California, USA). *Zootaxa* 3941 (4): 451-484. <https://doi.org/10.11646/zootaxa.3941.4.1>
- Hartop EA, Brown BV, Disney RHL (2016a) Flies from L.A., The Sequel: A further twelve new species of *Megaselia* (Diptera: Phoridae) from the BioSCAN Project in Los Angeles (California, USA). *Biodiversity Data journal* 4: e7756. <https://doi.org/10.3897/BDJ.4.e7756>
- Hartop EA, Wong MA, Eiseman CS (2016b) Material from The BioSCAN Project in Los Angeles (California: USA) yields another new species and leads to clarity for two confused type series of *Megaselia* Rondani (Diptera: Phoridae). *Proceedings of the Entomological Society of Washington* 118: 93-100. <https://doi.org/10.4289/0013-8797.118.1.93>



Figure 1.

*Megaselia marquezii* Hartop et al. hovering over *Psathyrella candolleana* (Fr.) Maire.




Figure 2.  
*Megaselia marquezii* Hartop et al. eggs laid in the gills of the *Psathyrella candolleana* (Fr.) Maire.



Figure 3.

A female *Megaselia marquezii* Hartop et al. feeding on *Psathyrella candolleana* (Fr.) Maire. A single larva can be seen feeding in the tissue beneath her.



Figure 4.

A close-up of a female *Megaselia marquezii* Hartop et al. feeding on *Psathyrella candolleana* (Fr.) Maire.





Figure 5.

*Megaselía marquezí* Hartop et al. inside the broken-open cap of a closed *Psathyrella candolleana* (Fr.) Maire. Based on the extended abdomen, she appears to have just oviposited.



Figure 6.

*Megaselia marquezii* Hartop et al. larvae feeding in the gills of the *Psathyrella candolleana* (Fr.) Maire.




Figure 7.

Stacked bar chart of phorid fly catch from one year of sampling at each of 30 sites from BioSCAN project. Each color represents a different species, each bar represents a single site. Black - *Megaselia marquezii* Hartop et al.; red - *Megaselia agarici* (Lintner). Vertical axis- number of specimens; horizontal axis- trapping sites (see Brown and Hartop 2016).




Figure 8.  
Numbers of specimens of male *Megaselia marquezii* Hartop et al. at BioSCAN site 5 throughout 2015.

Table 1.

*Megaselia* species known to be associated with the fungus *Psathyrella candolleana*.

<i>Megaselia</i> species	Region or country	Reference
<i>M. berndseni</i> (Schmitz, 1919)	England	Disney and Evans 1979, Disney and Evans 1990
<i>M. hirtiventris</i> (Wood, 1909)	England	Disney and Evans 1999
<i>M. latior</i> Schmitz. 1936	England	Disney and Evans 1999
<i>M. nigra</i> (Meigen 1830)	England	Disney and Evans 1999
<i>M. okazakii</i> Disney, 2014	Japan	Disney 1989
<i>M. rufipes</i> (Meigen, 1804)	Europe	Eisfelder 1956
<i>M. spinicincta</i> (Wood, 1910)	Europe	Eisfelder 1956