

Spiders (Arachnida: Araneae) in the semideciduous Atlantic Forest: An ecological and morphological trait dataset for functional studies

Ana Munévar^{‡,§}, Pedro Cardoso[§], Yolanda M.G. Piñanez Espejo[‡], Gustavo Andres Zurita[‡]

[‡] Instituto de Biología Subtropical (UNAM-CONICET), Puerto Iguazú, Argentina

[§] Laboratory for Integrative Biodiversity Research (LIBRe), Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland

[‡] Instituto de Biología Subtropical, Universidad Nacional de Misiones–CONICET, Puerto Iguazú, Misiones, Argentina. Facultad de Ciencias Forestales, Universidad Nacional de Misiones, Misiones, Argentina

Corresponding author: Ana Munévar (ana.munevar@helsinki.fi), Pedro Cardoso (pedro.cardoso@helsinki.fi)

Academic editor: Emma McCarroll Shaw

Abstract

Background

The semideciduous Atlantic Forest is one of the most diverse ecosystems in the world with a great diversity of spiders. Most spider-related studies in this ecosystem focused on species richness and composition; however, little is known about their trait diversity (including morphological, ecological and/or physiological traits). Two main datasets were compiled to generate a complete record of spider traits for this ecosystem.

New information

Here, we present two datasets about 259 species of spiders from the semideciduous Atlantic Forest of Argentina. The trait data set compiled information of morphological and ecological traits such as body size, femur length, ocular distance, foraging strategy, prey range, circadian activity and stratum preference; traits were assessed by species considering sexual dimorphism. The second dataset included information about phenology (season when spiders were collected), number of individuals assessed by species and presence/absence of spiders in the different sample sites. This dataset has high potential to help researchers in recording the state of a component of biodiversity (functional) and contributes with the study of ecosystem services and species conservation.

Keywords

Araneae, functional, traits, subtropical, forest, pine, plantations

Introduction

The Atlantic Forest of Argentina, Brazil and Paraguay is one of the most diverse ecosystems in the world; this biome hosts about 7% of the global known species richness and shows high levels of endemism (Oliveira-Filho and Fontes 2000, Eisenlohr et al. 2015). However, 90% of the Atlantic Forest has been replaced by intensive productive systems such as crops, livestock and tree plantations (Di Bitteti et al. 2003). The southern portion of the Atlantic Forest, located in the province of Misiones, Argentina and known as the semideciduous Atlantic Forest, preserves the largest continuous remnants of this ecoregion (Galindo-Leal and Câmara 2005).

Previous studies in the semideciduous Atlantic Forest have reported 550 species of spiders in Argentina (Rubio 2016, Rubio et al. 2018), 448 in Santa Catarina and 183 in Estado de Paraná, Brazil (Gonçalves da Rosa et al. 2019, Brito Pitilin et al. 2019). In contrast to the taxonomic approach, which focuses on species identity, functional diversity is a complementary approach that assumes species are not equal in the context of ecosystem functioning and their response to disturbances (Legras et al. 2018). However, functional studies require that morphological, ecological, physiological and behavioural traits of species are described (Violle et al. 2007).

In the Atlantic Rain Forest (northeast of Brazil), Gonçalves-Souza et al. (2014) described traits for 176 species of spiders; authors compiled four morphological (adhesive structures, eye arrangement/tapetum, body size and compression) and three ecological traits (sheltering behaviour, foraging period and mode) of spiders. In the semideciduous Atlantic forest, Raub et al. (2015) assessed functional diversity of secondary forests based on ecological traits of 220 species/morphospecies of spiders. Despite the species richness included in both studies, neither trait matrix (traits by species) nor details about number of individuals/species and sex considered were available.

Here, we present one of the most complete datasets of morphological and ecological traits for spiders inhabiting both native forests and pine plantations in the semideciduous Atlantic Forest of Argentina.

Project description

Study area description: This study was performed in the semideciduous Atlantic Forest of Argentina, in Misiones province. This ecosystem presents an average annual precipitation of 2000 mm, without a dry season and average temperatures of 15°C in winter (June–August) and 25°C in summer (September–March) (Ligier et al. 1990).

The native forest is composed of a complex and diverse vegetation, with three to five strata: three arboreal strata, a herbaceous stratum (50 cm high) composed of grasses and herbaceous plants and the lowest stratum dominated by mosses, saprophytes and terrestrial orchids (Morellato and Haddad 2000). The highest or emerging arboreal stratum is composed of trees up to 42 m high, usually covered with vines and epiphytes; the medium arboreal stratum reaches 30 m; and the lowest arboreal stratum or understorey consists of small trees, shrubs, woody bamboos and ferns.

Spider samples were collected from two habitats: areas of continuous native forest (Iguazú National Park, Uruguá-i Provincial Park and private reserves) and areas devoted to forestry planted with the exotic conifer *Pinus taeda* (Zurita and Bellocq 2012).

Sampling methods

Description: All individuals used in this study were collected to estimate changes in the taxonomic and the functional diversity of spiders, due to the replacement of the native forest by pine plantations. Changes on taxonomic diversity were previously published in Munévar et al. 2018, whereas functional changes will be published in a future manuscript. Spiders were collected during the summer (Feb-Mar), autumn (May-Jun), winter (Jul-Aug) and spring (Sep-Oct) of 2016 in protected areas of native forest and adjacent areas of pine plantations (*Pinus taeda*); seasonal fieldwork was conducted to include the potential phenology of spider activity (Vamosi et al. 2009). Five collection methods, including pitfall traps, Winkler, entomological vacuum (G-VAC), minor and major beating; methods were used to target different vegetation strata used by spiders (ground, litter layer, herbaceous, shrubs and low arboreal stratum, respectively) (Dias et al. 2010, Azevedo et al. 2014). Minor and major beating refers to collection of spiders by shaking shrubs and low arboreal vegetation; all sampling methods related to spider collection are detailed in Munévar et al. (2018).

All the individuals collected were preserved in alcohol (80%), counted and identified to the species level or morphospecies, using taxonomic literature (i.e. Herbert and Levi 1962, Lopes-Rodrigues and Mendonça 2011, Piacentini and Grismado 2008) and by consulting with specialists from the Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Buenos Aires, Argentina. All specimens were deposited in the spider collection of the “Instituto de Biología Subtropical (IBSI-Ara, G. Rubio) in Misiones, Argentina.

Sampling description: Three morphological and four ecological spider traits were recorded (see below). These traits have been widely used in functional studies of arthropods due to their association with the natural history of the species and habitat use (Brousseau et al. 2018). The number of individuals measured ranged from one to five (both males and females) according to the availability of specimens collected.

Morphological traits

The morphological traits measured were: 1) body size, 2) femur length and 3) ocular distance (Table 1). The measurements were taken from photographs captured using a

stereoscope Leica EZ4 D. All images were analyzed with Image J version 1.46r. This software allows transforming pixels to millimetres and measures distance and areas (Ferreira and Rasband 2012).

Ecological traits

The ecological traits were: 1) foraging strategy, 2) prey range, 3) circadian activity and 4) stratum preference (Table 1). All attributes of the traits were defined at family level using published literature (Cardoso et al. 2011, Dias et al. 2010); the presence or absence of an attribute for a determined trait is denoted by 1 or 0, respectively. Foraging strategy and prey range showed mutually exclusive attributes (e.g. spiders cannot present both, euryphagous and stenophagous diets), while stratum preference and circadian activity present multiple choice attributes (e.g. some species use both, ground and vegetation).

Geographic coverage

Description: The study area is located in northeast Argentina, in Misiones province. Coordinates show a polygon which encloses all sample sites (20 sites in total).

Coordinates: 25°48'44.72" S and 25°48'9.48" S Latitude; 54°18'58.31" O and 54°32'56.39" O Longitude.

Taxonomic coverage

Description: We collected a total of 15838 individuals. Only adults (32%) were identified to species/morphospecies level. We found a total of 368 species/morphospecies distributed in 38 families and 143 genera; the most species richness families were Theridiidae, Araneidae and Salticidae (62, 52 and 39 species, respectively), followed by Anyphaenidae (13 sp.), Corinnidae (12), Thomisidae (12) and Linyphiidae (9) (Fig. 1).

About 26% of the species were collected in the native forest and 40% in pine plantations; 34% of the species were shared between both sites.

From the complete assemblage, 259 species/morphospecies were selected from native forests and/or pine plantations; functional traits, previously mentioned, were assessed in 951 individuals. Species were selected, based on the availability and quality of specimens; the remaining species (109 from the total assemblage) were not in optimal conditions for measurement (in general, only one individual was captured by species).

Taxa included:

Rank	Scientific Name	Common Name
order	Araneae	Spiders

Usage licence

Usage licence: Creative Commons Public Domain Waiver (CC-Zero)

Data resources

Data package title: Semideciduous_Atlantic_Forest_Spiders

Number of data sets: 2

Data set name: Traits_of_Spider

Download URL: [10.6084/m9.figshare.11877783](https://doi.org/10.6084/m9.figshare.11877783)

Data format: Tab delimited file (.csv)

Description: Morphological and ecological traits of spider species in the semideciduous Atlantic Forest of Argentina.

Column label	Column description
Family	The full scientific name of the family in which the taxon is classified.
Scientific Name	The full scientific name.
Sex	The sex of the biological individual(s).
Body Size	Width and length of the prosoma and opisthosoma.
Femur length	Length of femur I & IV.
Ocular distance	Sum of diameters of one side eyes.
Foraging strategy	Tube web, Sheet web weaver, Space web, Orb web, Aerial hunter and Active hunter.
Prey range	Stenophagous, Euryophagous.
Circadian activity	Diurnal, Nocturnal.
Stratum preference	Ground (GR), Trunk (TR), Vegetation (VG).

Data set name:
Ecological_data_of_spiders_communities_present_in_native_forest_and_pine_plantations.

Download URL: [10.6084/m9.figshare.11877777](https://doi.org/10.6084/m9.figshare.11877777)

Data format: Tab delimited file (.csv)

Description: Dataset with presence/absence of species in each habitat type (native forests and pine plantations of different ages), number of individuals assessed,

phenology (season collected) and collection method for spiders inhabiting the semi-deciduous Atlantic Forest of Argentina.

Column label	Column description
InstitutionCode	The name (or acronym) in use by the institution having custody of the object(s) or information referred to in the record.
CollectionID	An identifier for the collection or dataset from which the record was derived.
Scientific Name	The full scientific name, with authorship and date information, if known.
No. Individuals Measured	Number of individuals measured by species.
Presence/Absence Matrix	Presence/absence matrix of species collected in all sample sites (native forest, mature plantations, middle age plantation and young plantation).
Phenology	Seasons of spider collection: 1= winter, 2= autumn, 3= spring, 4= summer.
Collection Method	Collection methods used to capture spiders by stratum. Pitfall trap = ground, Winkler = litter layer, Entomological vacuum (G-VAC) = herbaceous stratum (0-50 cm of height), Minor beating = shrubs (50 cm-2 m of height) and Major beating = low arboreal stratum (2-6 m of height).

Acknowledgements

We are grateful to Gonzalo Rubio, Cristian Grismado, Maria Eugenia Gonzalez, Luis Piacentini, Ivan Magalhaes and Martín Ramírez from the Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Buenos Aires, Argentina, who provided invaluable help during species identification. In addition, The Administration of National Parks (Argentina) for granting the necessary permissions and equipment to develop this project. To all members of the Laboratory for Integrative Biodiversity Research (LIBRe) for your support, and to reviewers for their comments on previous versions that allowed the improvement of this manuscript.

References

- Azevedo GHF, Faleiro BT, Magalhães ILF, Benedetti AR, Ubirajara O, Pena- Barbosa JP, Santos MT, Vilela PF, De Maria M, Santos AJ (2014) Effectiveness of sampling methods and further sampling for accessing spider diversity: a case study in a Brazilian Atlantic rainforest fragment. *Insect Conservation and Diversity* 7 (4): 381-391. <https://doi.org/10.1111/icad.12061>.
- Brito Pitilin R, Prado-Junior J, Brescovit AD, Tunes Buschini ML (2019) Climatic conditions drive the abundance and diversity of spiders community in an Atlantic Forest fragment. *Oecologia Australis* 23 (1): 39-55. <https://doi.org/10.4257/oeco.2019.2301.04>
- Brousseau PM, Gravel D, Handa IT (2018) On the development of a predictive functional trait approach for studying terrestrial arthropods. *Journal of Animal Ecology* 87 (5): 1209-1220. <https://doi.org/10.1111/1365-2656.12834>

- Cardoso P, Pekár S, Jocqué R, Coddington JÁ (2011) Global patterns of guild composition and functional diversity of spiders. *PLOS One* 6 (6): e21710. <https://doi.org/10.1371/journal.pone.0021710>
- Dias SC, Carvalho LS, Bonaldo AB, Brescovitc AD (2010) Refining the establishment of guilds in Neotropical spiders (Arachnida: Araneae). *Journal of Natural History* 44 (3-4): 219-239. <https://doi.org/10.1080/00222930903383503>
- Di Bitteti MS, Placci G, Dietz LA (2003) A biodiversity vision for the upper Paraná Atlantic Forest ecoregion: designing a biodiversity conservation landscape and setting priorities for conservation action. 1a ed. Nancy de Moraes. World Wildlife Fund, Washington, DC.
- Eisenlohr PV, Oliveira-Filho AT, Prado J (2015) The Brazilian Atlantic Forest: new findings, challenges and prospects in a shrinking hotspot. *Biodiversity and Conservation* 4 (9): 2129-2133. <https://doi.org/10.1007/s10531-015-0995-4>
- Ferreira T, Rasband W (2012) ImageJ user guide. ImageJ/Fiji.
- Foelix R (2011) *Biology of spiders*. 3rd Edition. Oxford University Press, USA.
- Galindo-Leal C, Câmara IG (2005) Atlantic Forest hotspot status: an overview. Brief history of conservation in the Paraná Forest. *The Atlantic Forest of South America: Biodiversity Status, Threats, and Outlook*.
- Gonçalves da Rosa M, Brescovit SD, Riviera Duarte Maluche Baretta C, Pires Santos JC, Iuñes de Oliveira Filho LC, Baretta D (2019) Diversity of soil spiders in land use and management systems in Santa Catarina, Brazil. *Biota Neotropica* 19 (2): e20180619. <https://doi.org/10.1590/1676-0611-bn-2018-0619>
- Gonçalves-Souza T, Santos AJ, Romero GQ, Lewinsohn TM (2014) Conservation along a hotspot rim: spiders in Brazilian coastal restingas. *Biodiversity and Conservation* 24: 1131-1146. <https://doi.org/10.1007/s10531-014-0846-8>
- Herbert W, Levi LR (1962) The genera of the spider family Theridiidae. *Bulletin of the Museum of Comparative Zoology. AT HARVARD COLLEGE* 127 (1).
- Legras G, Loiseau N, Gaertner J-C (2018) Functional richness: Overview of indices and underlying concepts. *Acta Oecologica* 87: 34-44. <https://doi.org/10.1016/j.actao.2018.02.007>
- Ligier HD, Matteio HR, Polo HL, Rosso JR (1990) *Provincia de Misiones. Atlas de suelos de la República Argentina*.
- Lopes-Rodrigues EN, Mendoça JM (2011) Araneid orb-weavers (Araneae, Araneidae) associated with riparian forests in southern Brazil: a new species, complementary descriptions and new records. *Zootaxa* 2759: 60-68. <https://doi.org/10.11646/zootaxa.2759.1.3>
- Morellato LC, Haddad CFB (2000) Introduction: The Brazilian Atlantic Forest. *Biotropica* 32: 786-792. <https://doi.org/10.1111/j.1744-7429.2000.tb00618.x>
- Munévar A, Rubio GD, Zurita GA (2018) Changes in spider diversity through the growth cycle of pine plantations in the semi-deciduous Atlantic forest: The role of prey availability and abiotic conditions. *Forest Ecology and Management* 24: 536-544. <https://doi.org/10.1016/j.foreco.2018.03.025>
- Nicholas MF, Baker S, Jordan G (2015) Moving beyond the guild concept: developing a practical functional trait framework for terrestrial beetles. *Ecological Entomology* 40 (1): 1-13. <https://doi.org/10.1111/een.12158>

- Oliveira-Filho AT, Fontes MAL (2000) Patterns of floristic differentiation among Atlantic Forests in southeastern Brazil and the influence of climate. *Biotropica* 32 (4d): 793-810. [https://doi.org/10.1646/0006-3606\(2000\)032\[0793:POFDAA\]2.0.CO;2](https://doi.org/10.1646/0006-3606(2000)032[0793:POFDAA]2.0.CO;2)
- Pekár S, Coddington JA, Blackledge TA (2011) Evolution of stenophagy in spiders (Araneae): Evidence based on the comparative analysis of spider diets. *Evolution* 66 (3): 776-806. <https://doi.org/10.1111/j.1558-5646.2011.01471.x>
- Piacentini LN, Grismado C (2008) Lobizon and Navira, two new genera of wolf spiders from Argentina (Araneae: Lycosidae). *Zootaxa* 2195: 1-33. <https://doi.org/10.11646/zootaxa.2195.1.1>
- Podgaiski LR, Joner F, Lavorel S, Moretti M, Ibanez S, Mendonca MS, Pillar JVD (2013) Spider trait assembly patterns and resilience under fire-induced vegetation change in south Brazilian grasslands. *PLOS One* 8 (3): e60207. <https://doi.org/10.1371/journal.pone.0060207>
- Raub F, Höfer H, Scheuermann L, Miranda de Brites R, Brandl R (2015) Conserving landscape structure – conclusions from partitioning of spider diversity in southern Atlantic forests of Brazil. *Studies on Neotropical Fauna and Environment* 50 (3): 158-174. <https://doi.org/10.1080/01650521.2015.1071959>
- Rubio GD (2016) Using a jumping spider fauna inventory (Araneae: Salticidae) as an indicator of their taxonomic diversity in Misiones, Argentina. *Revista de Biología Tropical* 64 (2): 875-883. <https://doi.org/10.15517/rbt.v64i2.19722>
- Rubio GD, Baigorria JE, Scioscia C (2018) Arañas salticidas de Misiones. Guía para la identificación (tribus basales). 1a ed. Universidad Maimónides: Ediciones Fundación Azara, Ciudad Autónoma de Buenos Aires.
- Vamوسي SM, Heard SB, Vamوسي JC, Webb CO (2009) Emerging patterns in the comparative analysis of phylogenetic community structure. *Molecular Ecology* 18 (4): 572-592. <https://doi.org/10.1111/j.1365-294X.2008.04001.x>
- Vandewalle M, de Bello F, Berg MP, Bolger T, Dolédec S, Dubs F, Feld CK, Harrington R, Harrison PA, Lavorel S, Martins da Silva P, Moretti M, Niemelä J, Santos J, Sattler T, Sousa JP, Sykes MT, Vanbergen AJ, Woodcock BA (2010) Functional traits as indicators of biodiversity response to land use changes across ecosystems and organisms. *Biodiversity and Conservation* 19: 2921-2947. <https://doi.org/10.1007/s10531-010-9798-9>
- Violle C, Navas ML, Vile D, Kazakou E, Fortunel C, Hummel I, Garnier E (2007) Let the concept of trait be functional! *Oikos* 116 (5): 882-892. <https://doi.org/10.1111/j.0030-1299.2007.15559.x>
- Zurita GA, Bellocq MI (2012) Bird assemblages in anthropogenic habitats: identifying a suitability gradient for native species in the Atlantic Forest. *Biotropica* 44 (3): 412-419. <https://doi.org/10.1111/j.1744-7429.2011.00821.x>




Figure 1.

Species richness of spiders by family collected in native forests and pine plantations within the semideciduous Atlantic Forest of Argentina.

Table 1.

Traits description and features

Traits	Description	Measure	Category	Source
Body size	Body size was estimated from four measures: width and length of prosoma and, width and length of opisthosoma.	Prosoma length: Distance between anterior edge of the carapace to the posterior end in dorsal view.	Continuous (mm)	Podgaiski et al. 2013
		Prosoma width : Mayor width of caparace in dorsal view.	Continuous (mm)	Podgaiski et al. 2013
		Opisthosoma width: In the middle of the abdomen, distance from superior to inferior edge in lateral view.	Continuous (mm)	Podgaiski et al. 2013
		Opisthosoma length: Distance between apex base to posterior end of abdomen in lateral view (without spinnerets).	Continuous (mm)	Podgaiski et al. 2013
Femur length	Femur length was estimated considering legs I & IV from one side.	Femur I & IV distance from anterior edge to posterior end, in prolateral view.	Continuous (mm)	Podgaiski et al. 2013
Ocular distance	Sum of diameters of one side of the caparace eyes.	Sum of diameter of four eyes (1 ALE, 1 PLE, 1 PME, 1 AME) from one side of the caparace.	Continuous (mm)	Vandewalle et al. 2010; Nicholas et al. 2015
Foraging strategy	Foraging strategy has six attributes or levels: Tube web, Sheet web weaver, Space web, Orb web, Aerial hunter and Active hunter. All species present just one foraging strategy.	present=1; absent=0	Binary	Foelix 2011

Prey range	Diet could be euryophagus (wide food range also called polyphagous) or stenophagous (restricted food range). Both attributes are mutually exclusive.	present=1; absent=0	Binary	Pekár et al. 2011
Circadian activity	Circadian activity can be diurnal and/or nocturnal. Attributes could be multiple choices (e.g. diurnal and nocturnal).	present=1; absent=0	Multiple choices	Foelix, 2011
Stratum preference	Stratum preference could be ground (GR), trunk (TR) and/or vegetation (VG). Attributes are multiple choices.	present=1; absent=0	Multiple choices	Cardoso et al. 2011, Dias et al. 2010