

Pseudidothea armata sp. n., a new isopod of the genus *Pseudidothea* (Crustacea, Malacostraca, Isopoda) from the Atlantic sector of the Southern Ocean

Nicholas Francesco Noli[‡], Davide Di Franco^{§,I}, Stefano Schiaparelli[‡], Angelika Brandt^{§,I}

[‡] Dipartimento di Scienze della Terra dell'Ambiente e della Vita (DISTAV), Università di Genova, Genova, Italy

[§] Senckenberg Research Institute and Natural History Museum, Frankfurt am Main, Germany

^I Goethe University Frankfurt, Institute for Ecology, Diversity and Evolution, Frankfurt am Main, Germany

Corresponding author: Nicholas Francesco Noli (nicholasfrancesco@hotmail.it)

Academic editor: Luis Ernesto Bezerra

ZooBank: [urn:lsid:zoobank.org:pub:B9BEE8EA-EC40-4DEE-B2BE-6CBBA33857B4](https://doi.org/10.21203/rs.3.rs-1000000/v1)

Abstract

Background

In the framework of the British Antarctic Survey (BAS) Expedition JR 15005 SO-AntEco, held in February-March 2016, the South Orkney Islands seafloor was sampled in order to investigate the distribution and composition of benthic communities around the area.

New information

A new species of the genus *Pseudidothea* Ohlin, 1901 is described from the Burdwood Bank area (South Orkney Islands). It has been collected during the SO-AntEco JR15005 RRS James Clark Ross expedition under the lead of the British Antarctic Survey (BAS). The new species, *Pseudidothea armata* sp. n., is very similar to *P. scutata* (Stephensen, 1947); however, it is characterised by peculiar supra-ocular spines and a different tubercular pattern. The study of the species of the *Pseudidothea* helps to better understand the diversity of the Pseudidotheidae in the Southern Ocean.

Keywords

Southern Ocean, morphology, ecology, distribution

Introduction

The family Pseudidotheidae Ohlin, 1901 is monogeneric with the only genus *Pseudidothea* Ohlin, 1901, comprising four accepted species. The family was erected through the description of its type species *P. bonnieri* Ohlin, 1901, now accepted as *P. miersi* (Studer, 1884) and was recently re-described by Poore and Bardsley (2004). According to WoRMS (World Register of Marine Species) and RAMS (Register of Antarctic Marine Species), only *P. scutata* Stephensen, 1947 (Stephensen 1947) occurs in the Southern Ocean in the area of the Antarctic Peninsula, off Elephant Island and South Shetlands. However, undetermined records of the genus *Pseudidothea* have been reported in several parts of the Antarctic Peninsula and in the Mawson Bank area, Ross Sea (Mills 2020) (Fig. 1).

The biogeographic knowledge of this genus is as follows: in the Southern Hemisphere, most of the records are from Chile (*P. miersi* (Studer 1884, Scarabino et al. 2008)), South Australia (*P. hoplites* (Poore and Bardsley 2004)) and New Zealand (*P. richardsoni* Hurley, 1957 (Hurley 1957)); only one specimen was found in the Southern Ocean in 1927 and has been described by Stephensen (1947) as *Microarcturus scutatus* (Stephensen, 1947). In 1990, it was re-described and placed in the genus *Pseudidothea* as *P. scutata* by Brandt and Wägele (1990).

In order to improve the biogeographic knowledge of the species of the Southern Ocean, *P. armata* sp. n. is described herein. Though only one specimen is available and used for the description, clear characteristics provide sufficient evidence that the new species differs from the closest morphological (and geographical) species *P. scutata*.

Materials and methods

Taxon sampling

The specimen was collected at Burdwood Bank, during the British Antarctic Survey (BAS) Expedition JR 15005 (https://www.bodc.ac.uk/resources/inventories/cruise_inventory/reports/jr15005.pdf) on board the RRS James Clark Ross, by means of a Rauschert dredge at a depth of 852 m (Station 143, 18 April 2016, 60°33.526'S, 41°5.306'W). After the first sorting on board, the specimen was stored and fixed with 96% ethanol in order to preserve it for further genetic analysis.

Photography and laboratory analyses

Only one specimen, the holotype of the new species, was found during the campaign. The holotype was not dissected in order to preserve it for further studies and it was drawn in dorsal and lateral views following standard descriptions (Wilson 2008). Drawings were performed using a camera lucida, followed by digital inking made by combining the stack

photos and the scanned hand-made drawings as layers. The graphic software used was Autodesk SketchBook and digital inking was performed with an XP-PEN Deco 02 graphic tablet. Stacks were obtained by using a Canon EOS 600D and a Leica 125 C, equipped with a Leica DMC 4500 camera. The use of stack photos as the base layer for digital-inking work is not new for crustacean illustrations (see, for example, the paper by Verheye and D'Udekem D'Acoz (2020)).

Additional distribution data

Additional distributional data of *Pseudidothea* (Rdmpage 2016, Registry-Migration.Gbif.Org 2016, Mackay 2018, Appiah-Madson and Distel 2019, Data Manager 2019, Mackay 2019, Mills 2020, Registry-Migration.Gbif.Org 2020, Mackay 2021, Orrell and Informatics Office 2021, Registry-Migration.Gbif.Org 2021, Tablado 2021) were provided through GBIF (Global Biodiversity Information Facility, available from <https://www.gbif.org/>) and OBIS (Ocean Biodiversity Information System <https://obis.org/>).

Quality check and data cleaning were performed using bibliographic research and the rgbif package (<https://CRAN.R-project.org/package=rgbif>) in RStudio software.

Maps were drawn using the QGIS (QGIS.org 2021) package QAntarctica (Matsuoka et al. 2018).

Morphological abbreviations

A = antenna;

P = pereopods;

PL = pleopod;

UR = uropod;

MNA = Italian National Antarctic Museum (Section of Genoa), Genoa;

PNRA = Italian National Antarctic Program;

BAS = British Antarctic Survey

Taxon treatment

***Pseudidothea armata* Noli, Di Franco, Schiaparelli, Brandt 2021, sp. nov.**

- ZooBank [F4392D41-91CF-458B-ACDE-533BAD913573](https://zoobank.org/F4392D41-91CF-458B-ACDE-533BAD913573)

Material

Holotype:

- a. kingdom: Animalia; phylum: Arthropoda; class: Malacostraca; order: Isopoda; family: Pseudidotheidae; genus: *Pseudidothea*; continent: Antarctica; locality: Burdwood Bank; verbatimDepth: 852; decimalLatitude: -60.55876; decimalLongitude: -41.08843; eventID: St143; samplingProtocol: bottom trawl; year: 2016; month: 3; day: 18; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: MNA 10749; identifiedBy: Nicholas Noli; dateIdentified: 2019; type: PhysicalObject; basisOfRecord: PreservedSpecimen; occurrenceID: 6D8BE133-6250-56E1-9B8C-C60147993C49

Description

Measurement. BL = 15 mm; BW = 5 mm.

Body: Entire body surface rough and granular, covered with small hair-like setae (Fig. 2A, B and Fig. 3A, B).

CEPHALOTHORAX. Head with two frontomedial lobes with many small tubercle-like protrusions. Cephalothorax with two large and stout spines, frontally directed (Fig. 2A, B and Fig. 3A, B, Suppl. materials 1, 2), covered with tubercles and with diffused short hair-like setae (not illustrated in drawings, but visible in Fig. 3). Two lateral eyes of medium size, slightly more subtriangular than oval (Fig. 2A B and Fig. 3A, B, Suppl. material 1).

ANTENNA 1. First Antenna (A1) consisting of three peduncular and two flagellar articles. First peduncular article broadest, almost surpassing in width the length of the second peduncular article, with one mediolateral simple bristle; second peduncular article long, nearly 1.5 times the first, the third almost a third of the second, rounded distally. First flagellar article short, ring-like, barely distinguishable; last flagellar article as long as the length of all the other articles of the A1. Distoventrally on this article, six pairs of aesthetascs, another single aesthetasc present in the tip of the A1 together with two simple setae (Fig. 2E and Fig. 3D).

ANTENNA 2. Second Antenna (A2) half as long as body, consisting of five peduncular and three flagellar articles. First peduncular article very small, second about double in length of the first; third peduncular article 3 times longer than second, with a two ventral rows of long simple setae, each group with one long and one short seta. Fourth and fifth peduncular articles longest, nearly subequal in length, with similar setae pattern of the third article. Last peduncular article with a distolateral small feather-like seta. First flagellar article about twice as long as second and third flagellar articles together, bearing short bristles and one longer apical simple seta. Second article smaller and narrower than first, with many simple short bristles. Last flagellar article smallest, claw-like. The whole antenna covered with lots of short and small hairs (Fig. 2D).

PEREONITES. Pereonite 1 fused with cephalothorax, but separated by a ridge. Pereonite 2 only slightly shorter than pereonite 3; pereonite 4 longest. Pereonite 5

slightly shorter than pereonite 2, pereonite 6 and 7 smallest and shortest. Pereonite 1 with one pair of large dorsal spine-like tubercles, anteriorly directed. Pereonites 2-4 with two pairs of tubercles (one spine-like dorsal pair and one shield-like lateral pair). Dorsal pair of tubercles in pereonites 2-4 are large and high, apically flattened, covered with few hair-like small setae and small tubercle-like protrusions; in pereonites 5-7, these are smaller, more slender, apically acute, but with blunt tip and more laterally flattened. Lateral pairs of tubercles are large in pereonites 2-4, smaller in pereonites 5-7. Evident elevations separated by deep grooves characterise the surface of pereonites 2-4, between the dorsal and lateral tubercles. Elevations are also present on pereonites 5-7. However, these are more shallow, barely visible and uniformly covering the surface between the dorsal and lateral tubercles (Fig. 2A, B and Fig. 3A, B).

PLEOTELSON. All pleonites fused to pleotelson, frontolaterally of pleotelson two stout and slightly rounded protrusions, caudally directed (Fig. 2A, B and Fig. 3E). Pleotelson frontally broadest, narrowing caudally. Tip of telson acuminate, but with blunt tip, slightly rounded, slightly bent dorsally; subapical telsonic spines absent, but two pairs of rounded large tubercles in two rows in the pleotelson, dorsally; the pleotelson lacks spines, but is covered entirely by medium-sized rounded protrusions, slightly smaller laterally (Fig. 2A, B and Fig. 3E).

PEREPODS. P1 shorter than P2-7. Basis and propodus longest; basis with few distoventral and distolateral simple setae; a row of small protuberances on the lateral outer-directed side of the article. Carpus trapezoidal, ischium, merus and carpus densely covered with sensory spines, especially on ventral and lateral surfaces. Subchelate propodus broad, oval, dactylus shorter than propodus, with two short smooth claws (the ventral one shorter) and a small spine in between (Fig. 4C). Ventral surface of propodus forming slightly concave "spoon", medial surface with few short simple bristles (Fig. 2C, Fig. 3C and Fig. 4D). P2-4 similar, one strong spine dorsolaterally on basis of P2-P4, distodorsal margins of ischium, carpus and propodus without strong spines, but with rough cuticular surface; merus of P2-4 presents an evident protrusion on the dorsal surface of the distal part; important protrusion is also present dorsomedially on ischium of P2 and P4. Setation in P2-4 is similar, but most prominent in P2. Merus, carpus and propodus with groups of setae arranged in two ventral rows, each group consisting of one long seta and one short seta (Fig. 4A and B). Small hairs present in all pereopods (Fig. 2F, G, H and Fig. 4A, B). P5-7 slightly shorter, but subequal in length to P2-4. P5 with two long setae on basis, P6 and 7 without long setae. Basis of P5-7 presenting one stout rounded spine dorsolaterally, in addition to smaller spines; stout spines are also present on carpus and merus (Fig. 2I, J, L and K). In P5-7, basis longest, ischium and propodus elongated compared to other articles; dactylus presenting a terminal claw with one simple seta in the terminal part. On propodus of P5-7, some feather-like bristles; all pereopods covered with small hairs (Fig. 2I, J and L).

UROPODS. Ventral surface of uropods covered with many simple hairs. Uropod elongated, both rami of uropods about subequal in length (Fig. 3E).

Diagnosis

The new species is characterised by one pair of large blunt anteriorly directed supra-ocular spines on the first pereonite. In dorsal view, these are long, divergent, forming a v-shape, reaching beyond the eyes. Pereonites 2-7 with only two pairs of tubercles: one dorsal pair of spine-like tubercles and one lateral pair of shield-like tubercles. Dorsal tubercles on pereonites 2-4 are large, apically flattened in lateral view, slightly anteriorly directed; on pereonites 5-7, these are smaller, slender, more pointed, but apically blunt and upwards directed. Pleotelson with blunt, strong protrusions.

Etymology

The species is named *armata* for its long supra-ocular and the dorso-apical spine-like tubercles, not simple “shielded” tubercles like *Pseudidothea scutata*, but more like strong blunt “spines”: as a contraposition to *P. scutata* meaning “that bears shield”, *P. armata* sp. n. “bears weapons”.

Distribution

Only known from type locality, the Burdwood Bank (Antarctica), found at 852 m.

Conservation

Specimen is stored and fixed with 96% ethanol in order to preserve it for further genetic analysis.

Remarks

Pseudidothea Ohlin, 1901 is clearly distinguishable from other genera by the oval shape of the body and peculiar conformation of pereonites and first pereopod. *P. bonnieri* was the type species representing the genus, described by Ohlin (1901). Subsequently, it was synonymised with *P. miersi* (Studer 1884) because of the many similarities with the latter species. The genus is only known from the Southern Hemisphere and it is mostly known from the Antarctic Peninsula in the Southern Ocean. A record of *Pseudidothea* Ohlin, 1901 was also recorded in the Ross Sea (Fig. 1), although undetermined to species level (Mackay 2018). To date, four species belong to the genus *Pseudidothea* as referred to above (original names are presented; the type species is marked with an asterisk):

Pseudidothea hoplites - Poore and Bardsley (2004)

**Pseudidothea miersi* - Ohlin 1901

Pseudidothea richardsoni - Hurley 1957

Pseudidothea scutata - Stephensen 1947

Pseudidothea armata sp. n.

Differential diagnosis

Within the genus *Pseudidothea*, *P. scutata* (Stephensen 1947) is most similar to *P. armata* sp. n. in shape and spine pattern. The main differences are the large supra-ocular spines and general body armature, that significantly differs from the apically flattened tubercles in *P. scutata*; similar is also the position of the flattened tubercles dorsally located in every pereonite of *P. scutata*; however, all of these structures are more like blunt spines in *P. armata* sp. n. Another difference can be found in the extreme reduction of the elevations present in between lateral flattened tubercles and dorsal spine-like tubercles. Brandt and Wägele (1990) re-described *P. scutata* and illustrated flattened, irregular elevations on pereonites 5-7, while in *P. armata* sp. n., these are barely present. The pleotelson of *P. scutata* and *P. armata* sp. n. is similar in shape and tubercular pattern; however, it completely lacks pleotelsonic dorsal spines in *P. armata*, only rounded, short and strong tubercles are present in the latter species.

Supra-ocular spines of *P. armata* are long, dorsally pointed and anteriorly directed, reaching beyond the eyes in dorsal view. *P. scutata* presents supraocular tubercles that are shorter, dorsally flattened and do not reach the eyes on dorsal view. In addition, supra-ocular spines in *P. armata* are divergent, forming a v-shape in dorsal view, while in *P. scutata* the supra-ocular tubercles are aligned parallel. The body armature in the two species shows a different pattern of tubercles. In *P. armata*, dorsal tubercles are more similar to blunt spines, apically less flattened than those present in *P. scutata*. In particular, *P. armata*'s spine-like tubercles on pereonites 5-7 are not apically flattened at all, but slender and pointed with a blunt apex, while in *P. armata*, pereonites 5-7 have small apically flat tubercles. *Pseudidothea scutata* presents raised areas in the lateral position, between the dorsal and lateral tubercles; these are marked on pereonites 5-7 and even described as distinct tubercles (Brandt and Wägele 1990, White 1992). *Pseudidothea armata* presents raised areas as well. However, elevations are more evident on pereonites 2-4, separated by deep grooves, making the surface between the dorsal and lateral tubercles look "wrinkly", while elevations on pereonites 5-7 are less visible. Basis of P5 in *P. armata* has two long setae, while there are only simple setae on P5 of *P. scutata*.

The pleotelson of *P. scutata* and *P. armata* is similar in shape and tubercular pattern; however, it completely lacks pleotelsonic dorsal spines in *P. armata*; only rounded, short and strong tubercles are present in the latter species.

Distribution

Only known from type locality.

Identification keys

***Pseudidothea* Ohlin, 1901**

This key to species is based to the *Pseudidothea* key present in the work of Poore and Bardsley "Pseudidotheidae (Crustacea: Isopoda: Valvifera) reviewed with description of a new species, first from Australia" (Poore and Bardsley 2004).

1	Pereonites 2 and 3 with forked dorsolateral spines; all pereonites with lateral rows of blade-like ridges, each with anteriorly and posteriorly directed spines; tergites produced laterally over coxae to form a shield with 3 points	<i>Pseudidothea hoplites</i>
–	Pereonites with low or high flat tubercles; tergites produced laterally as large tubercles or rounded or flattened laterally	2
2	Pereon with large high flat tubercles; pereonite 1 with dorsal pair, pereonites 2–4 with dorsal and lateral pair and pereonites 5–7 with dorsal, dorsolateral and lateral pair; male pleopod 2 with appendix masculina twice as long as rami	<i>Pseudidothea scutata</i>
–	Pereonites with high tubercles, laterally flattened; rounded, short and strong tubercles	3
3	Pereonites with strong laterally flattened tubercles; supra-ocular spines dorsally pointed and anteriorly directed, reaching beyond the eyes, forming a "v" shape in dorsal view; pereonites 5-7 with small apically flat tubercles	<i>Pseudidothea armata</i> sp. n.
–	Pereon with low irregular tubercles; male pleopod 2 with appendix masculina and rami subequal	4
4	Uropodal exopod with a single strong setae, endopod with 3 pappose setae; antenna 2 peduncle with long fine setae on articles 3–5; pereopods without tubercles; male pleopod 1 endopod with 5 lateral spinules proximally, 5 apical plumose setae; exopod with 15 spinules on lateral margin, tapering distally to an obtuse apex	<i>Pseudidothea richardsoni</i>
–	Uropod rami each with single seta; antenna 2 peduncle with short setae on articles 3–5; pereopods with tubercles; male pleopod 1 endopod with plumose setae marginally; exopod with 16–17 spinules laterally, with acute apex bent outwards	<i>Pseudidothea miersii</i>

Acknowledgements

We are indebted with Dr. Huw Griffiths, PI of the Cruise JR15005, part of the Biodiversity, Evolution and Adaptation team of the Environmental Change and Evolution Programme

at the British Antarctic Survey (BAS). Thanks are due to Claudio Ghiglione for sorting and fixation of the Rauschert materials. Compelling and sincere are the thanks to the Senckenberg Institute Crustacea Section of the Marine Zoology Department and to the Italian National Antarctic Museum (MNA, Section of Genoa) and their respective staff for the continuous support and collaboration. This paper was written during the Erasmus+ experience of the first author.

References

- Appiah-Madson H, Distel D (2019) Ocean genome legacy collection. United States Geological Survey. Type: dataset. <https://doi.org/10.15468/JZGQTC>
- Brandt A, Wägele JW (1990) Redescription of *Pseudidothea scutatas* (Stephensen, 1947) (Isopoda, Valvifera) and adaptations to a microphagous nutrition. *Crustaceana* 58 (1): 97-105. <https://doi.org/10.1163/156854090X00813>
- Data Manager (2019) South American Antarctic marine biodiversity literature. SCAR - AntOBIS. Type: dataset. <https://doi.org/10.15468/KXNPUQ>
- Hurley DE (1957) Some Amphipoda, Isopoda and Tanaidacea from Cook Strait. *Zoology Publications Victoria University Collections* 21: 1-20.
- Mackay K (2018) Ross Sea biodiversity survey 2004 (BioRoss). The National Institute of Water and Atmospheric Research (NIWA). Type: dataset. <https://doi.org/10.15468/6XYT8D>
- Mackay K (2019) Biological observations from the Discovery investigations 1925-1952. Southwestern Pacific Ocean Biogeographic Information System (OBIS) Node. Type: dataset. <https://doi.org/10.15468/QOQBU7>
- Mackay K (2021) Marine biological observation data from coastal and offshore surveys around New Zealand. The National Institute of Water and Atmospheric Research (NIWA). Type: dataset. <https://doi.org/10.15468/PZPGOP>
- Matsuoka K, Skoglund A, Roth G (2018) Quantarctica. Norwegian Polar Institute. Type: dataset. <https://doi.org/10.21334/NPOLAR.2018.8516E961>
- Mills S (2020) NIWA invertebrate collection. The National Institute of Water and Atmospheric Research (NIWA). Type: dataset. <https://doi.org/10.15468/6QGSWO>
- Ohlin AG (1901) Isopoda from Tierra Del Fuego and Patagonia, I. Valvifera. *Norstedt & söner*
- Orrell T, Informatics Office (2021) NMNH extant specimen records. National Museum of Natural History, Smithsonian Institution. Type: dataset. <https://doi.org/10.15468/HNHRG3>
- Poore GB, Bardsley T (2004) Pseudidotheidae (Crustacea: Isopoda: Valvifera) reviewed with description of a new species, first from Australia. *Memoirs of Museum Victoria* 61 (1): 75-83. <https://doi.org/10.24199/j.mmv.2004.61.6>
- QGIS.org (2021) QGIS Geographic Information System. <http://www.qgis.org>
- RdmPAGE (2016) International Barcode of Life project (iBOL). The International Barcode of Life Consortium. Type: dataset. <https://doi.org/10.15468/INYGC6>
- Registry-Migration.Gbif.Org (2016) Southern Ocean Isopoda. British Antarctic Survey. Type: dataset. <https://doi.org/10.15468/LAITOD>
- Registry-Migration.Gbif.Org (2020) Collection Crustacea - ZMB. Senckenberg. Type: dataset. <https://doi.org/10.15468/FWGHFF>

- Registry-Migration.Gbif.Org (2021) Australian Museum provider for OZCAM. Australian Museum. Type: dataset. <https://doi.org/10.15468/E7SUSI>
- Scarabino F, Roccatagliata D, Doti B (2008) Range Extension of *Pseudidothea Miersii* (Studer, 1884) (Isopoda, Valvifera) and New Evidence of Its Likely Commensal Relationship with a Sea Urchin. *Crustaceana* 81 (7): 883-888. <https://doi.org/10.1163/156854008784771694>
- Stephensen K (1947) Tanaidacea, Isopoda, Amphipoda and Pycnogonida. Scientific results of the Norwegian Antarctic Expeditions 1927-28 et sqq. 2 (27): 1-90.
- Stephensen KH (1947) Tanaidacea, Isopoda, Amphipoda, and Pycnogonida. Norwegian Antarctic Expeditions (1927-1928 et sqq.).
- Studer T (1884) Isopoden, gesammelt während der reise SMS Gazelle um die erde 1874-76. Kurfürstlich Akademie der Wissenschaften, Berlin, 27 pp.
- Tablado A (2021) Museo Argentino de Ciencias Naturales "Bernardino Rivadavia" (MACN). Invertebrates National Collection (MACNIn). Museo Argentino de Ciencias Naturales. Type: dataset. <https://doi.org/10.15468/UUZ636>
- Verheye ML, D'Udekem D'Acoz C (2020) Integrative taxonomy of giant crested *Eusirus* in the Southern Ocean, including the description of a new species (Crustacea: Amphipoda: Eusiridae). *Zoological Journal of the Linnean Society* XX: 1-47. [In english]. <https://doi.org/10.1093/zoolinnean/zlaa141>
- White M (1992) Antarctic Isopoda Valvifera J. W. Wägele Koeltz Scientific Books, Königstein (1991) 213 pages. 130DM. ISBN 3 87429 324 6. *Antarctic Science* 4 (1): 121-121. <https://doi.org/10.1017/S0954102092240189>
- Wilson GF (2008) A review of taxonomic concepts in the Nannoniscidae (Isopoda, Asellota), with a key to the genera and a description of *Nannoniscus oblongus* Sars. *Zootaxa* 1680 (1). <https://doi.org/10.11646/zootaxa.1680.1.1>




Figure 1.

Distribution records of *Pseudidothea* Ohlin, 1901 in the Southern Ocean. Light blue square represents the location of *Pseudidothea armata* sp. n, recorded during the SO-AntEco JR15005 RRS James Clark Ross expedition, in Burdwood Bank area (South Orkneys) in the framework of the British Antarctic Survey (BAS), 2016.




Figure 2.

Pseudidothea armata sp. n. male holotype (MNA 10749). **A** dorsal view; **B** lateral view; **C** P1. Scale refers to 0.5 mm length. **D** A2; **E** A1; **F** P2; **G** P3; **H** P4; **I** P5; **J** P6; **K** Dactylus of P6; **L** P7. Scale refers to 1 mm length for **D-L**.




Figure 3.

Pseudidothea armata sp. n. male holotype (MNA 10749). **A** stack photo of lateral view; **B** stack photo of dorsal view; **C** stack photo of P1, scale bar represents 0.5 mm; **D** stack photo of A1, scale bar represents 1 mm; **E** stack photo of pleotelson. Scale bar represents 1 mm length.




Figure 4.

Pseudidothea armata sp. n. male holotype (MNA 10749). **A** left P2; **B** right P2; **C** right P1, dorsal part; **D** right P1, focus on inner part. Scale refers to 1 mm length.

Supplementary materials

Suppl. material 1: MNA 10749 LATERAL

Authors: Nicholas Noli

Data type: Digital inking drawing

Brief description: *Pseudidothea armata* sp. n. male holotype (MNA 10749) lateral view.

[Download file](#) (22.43 MB)

Suppl. material 2: MNA 10749 DORSAL

Authors: Nicholas Noli

Data type: Digital inking drawing

Brief description: *Pseudidothea armata* sp. n. male holotype (MNA 10749) dorsal view.

[Download file](#) (20.65 MB)