

The extremely high diversity of Collembola in relict forests of Primorskii Krai of Russia

Nataliya Kuznetsova[‡], Anna Bokova[‡], Alexander Kuprin[§], Mikhail Potapov[‡], Yulia Shveenкова[‡], Natalya Ivanova[¶]

[‡] Moscow State Pedagogical University, Moscow, Russia

[§] Federal Scientific Center of the East Asia Terrestrial Biodiversity Far Eastern Branch of the Russian Academy of Sciences, Vladivostok, Russia

[‡] Privolzhskaya Lesostep Nature Reserve, Penza, Russia

[¶] Institute of Mathematical Problems of Biology RAS – the Branch of Keldysh Institute of Applied Mathematics of Russian Academy of Sciences, Pushchino, Russia

Corresponding author: Nataliya Kuznetsova (mpnk@yandex.ru), Natalya Ivanova (natalya.dryomys@gmail.com)

Academic editor: Nina Filippova

Abstract

Background

The diversity of soil animals of relict forest ecosystems in East Asia continues to be insufficiently studied and almost not represented in international databases, including GBIF. This article is based on 7550 records of 175 species which were collected in Ussuriiskii, Sikhote-Alinskii and Kedrovaya Pad' natural reserves of Russian Far East in 2016–2017. A multi-scale sampling design allowed us to estimate population densities and local species richness of Collembola at areas of different sizes. The work continues [the digitization of the collections of the Moscow Pedagogical State University \(MPSU\)](#) and their publication through [GBIF.org](#), which began in 2019. This article is based on original data including 2377 specimens of springtails from eight forests and 648 soil cores.

New information

Within the framework of modern taxonomy, this work represents the first publication of lists of Collembola species of forests of Primorsky Krai. The work focused on the relict protected cedar-deciduous forests. Nine species new to science were described and data on the fauna of the region were significantly revised. Considerable contribution was made to the biogeography of Collembola of East Asia. The design of the sampling allowed us to draw conclusions about the species saturation of springtails at various spatial scales within the habitat: from a few square cm to 100 sq. m. Number of species reached record high values reflecting the benchmark state of Collembola communities of undisturbed old temperate forest ecosystems.

Keywords

sampling event, springtails, soil fauna, mesofauna, microarthropods, broad-leaf – cedar pine forests, natural reserves, multi-scale sampling design, species richness, population density, Ussuriiskii natural reserve, Sikhote-Alinskii natural reserve, Kedrovaya Pad' natural reserve

Introduction

Collembola, or springtails, is one of the most abundant and diverse groups of soil microarthropods that play an important role in the processes of destruction of organic residues (Petersen and Luxton 1982, Hopkin 1997). We aimed to estimate the upper limit of the local species diversity of Collembola in a region known for its high species richness of various groups of organisms (Latham and Ricklefs 1993). Part of this region is located in the south of the Russian Far East, where broad-leaf – cedar pine forest ecosystems are distributed. The highest diversity of insects in the temperate zone is described for these forests (Chernov et al. 2011). UNESCO included this area in the World Heritage List ([Central Sikhote-Alin](#)) noting that “the combination of glacial history, climate and relief has allowed the development of the richest and most unusual temperate forests in the world”.

Regular research of Collembola in the south of the Primorskii Krai of Russia began about half a century ago showing a great originality of the fauna (e.g. Kutyreva 1979, Kutyreva 1984, Martynova 1988). During the last decades, new approaches to taxonomy have led to the taxonomic revision of many genera of springtails (Pomorski and Sveenkova 2006, Deharveng et al. 2011, Jie et al. 2011, Jordana et al. 2011, Huang and Potapov 2012, Smolis et al. 2012, Smolis and Deharveng 2015). However, these works did not focus on the issues of local species richness.

We collected springtails in the region in the years 2016–2017. The data from 2016 was published (Kuznetsova et al. 2019). However, the species list was limited to abundant species only. The present study includes the full species list, based on samples from 2016 and 2017 taking into account the recently described new species for science. A specific multi-scale sampling design allowed us to estimate a local species richness of Collembola at areas of different sizes.

General description

Purpose: The purpose of the paper is to present information on species composition and abundances of Collembola in relict broad-leaf – cedar pine forest ecosystems of the Russian Far East. Diversity and abundance are presented in the most detailed form of sampling-evidence.

Sampling methods

Description: The dataset (Kuznetsova et al. 2021) provides information on the number of individuals of springtail species collected in broad-leaf – cedar pine forests of three natural reserves in July 2016 and August 2017 (7550 occurrences). Three forests were examined in Ussuriiskii natural reserve, 2 – in Kedrovaya Pad', 2 – in Sikhote-Alinskii natural reserve and 1 – in Chuguevsky District. The sampling plots “Fauri” and “Grabovaya” were in mountain forest, plots “Turova” and “Kedrovaya” were on a slope, plots “Pikhtovaya”, “Kema”, “Chuguev” and “Anikin” were in the river valleys. The material includes about 24 thousands individuals of 175 species from eight sampling series and 648 soil cores. They were collected by Natalia Kuznetsova, Mikhail Potapov, Anna Geraskina, Alexander Kuprin, Anastasia Korotkevich and were identified by Natalia Kuznetsova, Mikhail Potapov, Babenko Anatoly, Shveenкова Yulia.

Sampling description: The sampling was based on a multi-scale design to study the structure of biodiversity at different spatial scales (Lande 1996, Azovsky et al. 2000). A fractal arrangement of cores allows us to reduce sample effort because the same core is used for the analysis at different scales (Marsh and Ewers 2012). A few cases of applying the approach in soil zoology include testate amoebae (Tsyganov et al. 2014), oribatid mites (Bolger et al. 2014) and springtails (Kuznetsova and Saraeva 2018). We used the small size of the corer (8 cm² in section) to implement special attention on the diversity and spatial structure of the population at the micro level. Soil was investigated down to 20 cm. A total of 81 cores were taken in each sampling plot. Cores were placed in the corners of different-scale equilateral triangles inscribed in squares with sides 10 cm, 25 cm, 1 m and 10 m. The different-scale triangles were designed following the principles of fractal geometry. The sample design is described in detail by Saraeva et al. (2015).

Extraction of Collembola from cores: Plastic containers were used for storage and transportation of individual cores (Fig. 1). Each core was placed in a 0.3 litre plastic container with a vent hole covered with a gas cloth (Fig. 2). The containers were transported to MPGU (Moscow). Extraction of Collembola was performed in the laboratory using Tullgren's funnels at approximately 25°C. Extraction into 70% alcohol was continued for 4-5 days until the cores were completely dry.

Laboratory processing: All the specimens were mounted on slides in Phoea liquid according to a standard procedure (Ghilarov 1975, Potapov and Kuznetsova 2011).

Sampling plots: Short descriptions of some sampling plots (“Turova”, “Grabovaya”, “Kedrovaya”, “Pikhtovaya”) were published (Kuznetsova et al. 2019). The descriptions of the other plots are given for the first time. All the plots were located on brown soil.

The sampling plot “Turova” (Figs 3, 4) is in the Ussuriiskii natural reserve; it is a cedar pine–deciduous forest on a flat slope (*Pinus koraiensis* Siebold & Zucc., *Acer mandshuricum* Maxim., *Quercus mongolica* Fisch. etc.); in the undergrowth *Lonicera maackii* (Rupr.) Maxim., *Eleutherococcus senticosus* (Rupr. & Maxim.) Maxim. etc.; in the

above-soil cover [*Oxalis acetosella* L.](#), [*Carex* L.](#), [*Adiantum pedatum* L.](#) etc. The dead cover was ~10 cm.

The sampling plot “Grabovaya” (Fig. 5) is in the Ussuriiskii natural reserve; it is on Mount Grabovaya, in fir–hornbeam forest ([*Abies holophylla* Maxim.](#), [*Carpinus cordata* Blume](#), [*Pinus koraiensis*](#), [*Betula costata* Trautv.](#) etc.); in the undergrowth are [*Caprinus cordata*](#), [*Ac er tegmentosum* Maxim.](#), [*Acer barbinerve* Maxim. ex Miq.](#), [*Eleutherococcus senticosus*](#) etc.; in the above-soil cover are [*Oxalis acetosella*](#), [*Leptorumohra amurensis* \(Milde\) Tzvelev](#) etc. The dead cover is > 80% and the thickness of the litter is ~ 4 cm.

The sampling plot “Kedrovaya” is in the Kedrovaya Pad' natural reserve; it is a cedar pine–fir broadleaved forest on a slope ([*Pinus koraiensis*](#), [*Abies holophylla*](#), [*Tilia mandshurica* Rupr. & Maxim.](#), [*Carpinus cordata*](#) etc.); in the undergrowth are five maple species, including [*Acer tegmentosum*](#) and [*Acer barbinerve*](#); in the above-soil cover are [*Leptorumohra amurensis*](#), [*Dryopteris crassirhizoma* Nakai](#), [*Maianthemum dilatatum* \(Alph.Wood\) A.Nelson & J.F.Macbr.](#), [*Oxalis acetosella*](#) etc. The dead cover ~ 50% and the litter thickness is 4 cm.

The sampling plot “Pikhtovaya” is in the Kedrovaya Pad' natural reserve; it is a valley fir and deciduous forest ([*Abies holophylla*](#), [*Juglans mandshurica* Maxim.](#), [*Pinus koraiensis*](#) etc.); in the undergrowth are [*Juglans mandshurica*](#), [*Carpinus cordata*](#), [*Acer mono* Maxim.](#), [*Acer tegmentosum*](#), [*Acer mandshuricum*](#) etc.; in the above-soil cover are [*Leptorumohra amurensis*](#), [*Dryopteris crassirhizoma*](#) etc. The dead cover is ~ 50% and the thickness of the litter is 3–4 cm.

The sampling plot “Fauri” (Figs 6, 7) is in the Sikhote-Alinskii natural reserve, Kabani station, at 932 m alt.; it is a coniferous wood ([*Abies nephrolepis* \(Trautv. ex Maxim.\) Maxim.](#), [*Betula platyphylla* Sukaczev](#), [*Pinus koraiensis*](#)); in the undergrowth are [*Rhododendrom fauriei* Franch.](#), [*Acer ukurunduense* Trautv. & C.A.Mey.](#), [*Picea jezoensis* \(Siebold & Zucc.\) Carrière](#); in the above-soil cover are [*Leptorumohra amurensis*](#), [*Oxalis acetosella*](#), [*Maianthemum bifolium* \(L.\) F.W.Schmidt](#) etc. The dead cover is 50–100% and the thickness of the litter is 3–7 cm.

The sampling plot “Kema” is nearby the Sikhote-Alinskii natural reserve, in the valley of Brusnichnaya River (tributary of the Kema); it is mixed forest ([*Pinus koraiensis*](#), [*Populus maximowiczii* A.Henry](#), [*Picea jezoensis*](#), [*Ulmus glabra* Huds.](#), [*Abies nephrolepis*](#)); in the undergrowth are [*Acer mono*](#), [*Acer barbinerve*](#), [*Acer tegmentosum*](#), [*Philadelphus tenuifolius* Rupr. & Maxim.](#), [*Eleutherococcus senticosus*](#) etc.; in the above-soil cover are [*Leptorumohra amurensis*](#), [*Oxalis acetosella*](#), [*Maianthemum bifolium*](#), [*Carex siderosticta* Hance](#), [*Cardamine leucantha* \(Tausch\) O.E.Schulz](#), [*Cacalia hastata* L.](#) etc. The dead cover is 5–65% and the thickness of the litter is 7–10 cm.

The sampling plot “Chuguev” (Figs 8, 9) is in the Chuguevski District near Verkhneussuriyski Station of the Federal Scientific Center of the East Asia Terrestrial Biodiversity; it is a valley mixed forest ([*Abies nephrolepis*](#), [*Populus maximowiczii*](#), [*Fraxinus mandshurica* Rupr.](#), [*Betula costata*](#), [*Pinus koraiensis*](#) etc.); in the above-soil cover are [*Abies nephrolepis*](#), [*Acer*](#)

mono, *Acer tegmentosum*, *Lonicera maackii*, *Philadelphus tenuifolius* etc.; in the above-soil cover are *Oxalis acetosella*, *Leptorumohra amurensis*, [Carex campylorhina V.I.Krecz.](#), *Cardamine leucantha*, [Athyrium rubripes \(Kom.\) Kom.](#) etc.). The dead cover is 20–85% and the thickness of the litter is 6–9 cm.

The sampling plot “Anikin” is in the Ussuriyskii natural reserve, Suvorovskoye forest district, Anikinsky station, valley of Anikin River; it is valley broadleaf forest (*Juglans mandshurica*, *Populus maximowiczii*, *Fraxinus mandshurica*) with *Pinus koraiensis*; in the above-soil cover are *Carex* L., *Leptorumohra amurensis* etc. The dead cover is 50–90% and the thickness of the litter is 3–4 cm.

Quality control: We used both modern taxonomic papers and keys (Martynova 1988, Babenko et al. 1994, Potapov 2001) for the taxonomic determination of Collembola. The material was checked by leading experts in taxonomy of Collembola. Scientific names were checked using the [GBIF species matching tool](#).

Step description: Data on species were digitised, standardised according to the Darwin Core (Wieczorek et al. 2012), the quality of the data was checked and errors were corrected and then published through [GBIF.org](#) (Kuznetsova et al. 2021).

Geographic coverage

Description: Primorskii Krai of the Russian Far East (Fig. 10).

Coordinates: 43.115 and 45.648 Latitude; 131.487 and 137.01 Longitude.

Taxonomic coverage

Description: So far, the taxonomical knowledge of different families and genera of Collembola is highly irregular in the area under study. Our identification of particular groups of Collembola, therefore, considerably depended on taxa. The species of Neelidae, Symphypleona, Lepidocyrtinae and Entomobryini were identified, based on the appearance (body size, colour pattern, length of limbs and other easily recognisable features), other taxa - on modern taxonomy, family Tomoceridae - on traditional characters. Families Hypogastruridae, Onychiuridae and Isotomidae were identified down mostly to species level, while the genera *Isotoma* and *Desoria* still are less certain and were differentiated as morpho-species (sp. 1, sp. 2 etc). Family Odontellidae is less understood in the area and so it was mostly represented by the "sp." in the list. Some species were described as new to science in the material: *Anurida* - 6 spp. n. (Babenko et al. 2019), *Oligaphorura* – 2 spp. n. (Xin et al. 2019); *Pseudachorutes* – 1 sp. n. and re-description of three species (Babenko et al. 2021).

Taxa included:

Rank	Scientific Name
------	-----------------

phylum	Arthropoda
class	Collembola

Temporal coverage

Data range: 2016-7-23 - 2016-7-29; 2017-8-06 - 2017-8-13.

Usage licence

Usage licence: Other

IP rights notes: [Creative Commons Attribution \(CC-BY\) 4.0 License](#)

Data resources

Data package title: Collembola of the relict forests of the Russian Far East.

Resource link: <https://www.gbif.org/dataset/321e6294-7e96-44c2-ac5d-6b009ef17618>

Number of data sets: 1

Data set name: Collembola of the relict forests of the Russian Far East.

Character set: UTF-8

Data format: Darwin Core Archive

Column label	Column description
eventID	An identifier for the event https://dwc.tdwg.org/terms/#dwc:eventID
samplingProtocol	Sampling protocol (Tullgren funnels) https://dwc.tdwg.org/terms/#dwc:samplingProtocol See details in the Sampling methods section.
sampleSizeValue	Size of the sampling core (8 cm ²). https://dwc.tdwg.org/terms/#dwc:sampleSizeValue See details in the Sampling methods section.
sampleSizeUnit	The unit of measurement of the size sampling core (cm ²) https://dwc.tdwg.org/terms/#dwc:sampleSizeUnit See details in the Sampling methods section.
decimalLatitude	The geographic latitude in decimal degrees of the geographic centre of the data sampling place. https://dwc.tdwg.org/terms/#dwc:decimalLatitude
decimalLongitude	The geographic longitude in decimal degrees of the geographic centre of the data sampling place. https://dwc.tdwg.org/terms/#dwc:decimalLongitude

geodeticDatum	Spatial reference system (SRS) upon which the geographic coordinates given in decimalLatitude and decimalLongitude are based. https://dwc.tdwg.org/terms/#dwc:geodeticDatum
coordinateUncertaintyInMetres	The maximum uncertainty distance in metres. https://dwc.tdwg.org/terms/#dwc:coordinateUncertaintyInMeters
coordinatePrecision	The fraction of a degree corresponding to the number of significant digits in the source coordinates. https://dwc.tdwg.org/terms/#dwc:coordinatePrecision
country	Country name (Russian Federation). https://dwc.tdwg.org/terms/#dwc:country
countryCode	The standard code for the Russian Federation according to ISO 3166-1-alpha-2 (RU). https://dwc.tdwg.org/terms/#dwc:countryCode
stateProvince	Region name. The first level administrative division. https://dwc.tdwg.org/terms/#dwc:stateProvince
locality	The specific description of the place. https://dwc.tdwg.org/terms/#dwc:locality
locationID	An identifier for the set of location information https://dwc.tdwg.org/terms/#dwc:locationID We used this term to link cores (events) in the sampling plot.
habitat	A description of the habitat in which the Event occurred https://dwc.tdwg.org/terms/#dwciri:habitat We indicated habitat type as a landscape position (valley or mountain) and tree species dominates in the canopy.
verbatimEventDate	The verbatim original date of the Event occurred. https://dwc.tdwg.org/terms/#dwc:verbatimEventDate
year	The four-digit year of the Event occurred. https://dwc.tdwg.org/terms/#dwc:year
month	The integer month of the Event occurred. https://dwc.tdwg.org/terms/#dwc:month
day	The integer day of the month of the Event occurred. https://dwc.tdwg.org/terms/#dwc:day
eventDate	Field data collection date (YYYY-MM-DD). https://dwc.tdwg.org/terms/#dwc:eventDate
institutionCode	The acronym of the Institute. https://dwc.tdwg.org/terms/#dwc:institutionCode
institutionID	An identifier for the institution having custody of the object(s) or information referred to in the record. https://dwc.tdwg.org/terms/#dwc:institutionID
basisOfRecord	Basis of the record (PreservedSpecimen). https://dwc.tdwg.org/terms/#dwc:basisOfRecord
occurrenceID	An identifier for the record. https://dwc.tdwg.org/terms/#dwc:occurrenceID
identificationRemarks	Original identification. The dwc:verbatimIdentification was not used because it is currently not supported on the IPT. https://dwc.tdwg.org/terms/#dwc:identificationRemarks
scientificName	Scientific name. https://dwc.tdwg.org/terms/#dwc:scientificName

identificationQualifier	A brief phrase or a standard term ("cf.", "aff.") to express the determiner's doubts about the Identification. https://dwc.tdwg.org/terms/#dwc:identificationQualifier
taxonRank	The taxonomic rank. https://dwc.tdwg.org/terms/#dwc:taxonRank
kingdom	The full scientific name of the kingdom (Animalia). https://dwc.tdwg.org/terms/#dwc:kingdom
phylum	The full scientific name of the phylum. https://dwc.tdwg.org/terms/#dwc:phylum
class	The full scientific name of the class. https://dwc.tdwg.org/terms/#dwc:class
identifiedBy	List of persons, who identified collected Collembola. https://dwc.tdwg.org/terms/#dwc:identifiedBy
identificationReferences	DOI of references used in the identification. Used for taxa, which did not match the GBIF Backbone. https://dwc.tdwg.org/terms/#dwc:identificationReferences
lifeStage	The life stage of individuals. Here it is used for juvenile individuals indicated. https://dwc.tdwg.org/terms/#dwc:lifeStage
individualCount	The number of individuals represented in the core. https://dwc.tdwg.org/terms/#dwc:individualCount
occurrenceStatus	A statement about the presence or absence of a Taxon at a Location. https://dwc.tdwg.org/terms/#dwc:occurrenceStatus
language	A language of the resource (EN). https://dwc.tdwg.org/terms/#dwc:language

Additional information

In total, in the relict forests of the Far East, we found the highest diversity of Collembola that has ever been observed in the ecosystems of the temperate zone and possibly the world: up to 90 species per area 10 x 10 m (sampling plot). The species saturation reaches 30 species on an area of 8 square centimetres (one core) and can exceed 60 species on 1 square metre (Fig. 11).

Acknowledgements

The study was funded by the Russian Fund for Basic Research, project 16-04-01228-a. The authors are grateful to Anatoly Babenko for the taxonomical analysis of the part of the material, to Svetlana Bondarchuk for the botanical consultations and to Anna Geraskina and Anastasia Korotkevich who participated in the sampling of the material. We are also thankful to the administration and employees of Ussuriiskii Nature Reserve, Sikhote-Alinskii natural reserve and the national park Zemlya Leoparda (which includes Kedrovaya Pad' nature reserve) for help in conducting field research and to Valeria Makeeva and Maria Antipova for assistance in laboratory processing of samples. We are grateful to reviewers Yuri Mazei and Anastasia Taskaeva for their useful comments. The work of Natalya Ivanova was done within the framework of the Basic Scientific Research

theme of the Institute of Mathematical Problems of Biology RAS – the Branch of Keldysh Institute of Applied Mathematics of the Russian Academy of Sciences.

The authors declare that they have no conflict of interest.

References

- Azovsky AI, Chertoproud MV, Kucheruk NV, Rybnikov PV, Sapozhnikov FV (2000) Fractal properties of spatial distribution of intertidal benthic communities. *Marine Biology* 136 (3): 581-590. <https://doi.org/10.1007/s002270050718>
- Babenko A, Potapov M, Stebaeva S, Chernova N (1994) Keys of springtails of the fauna of Russia and adjacent countries: family Hypogastruridae. Nauka, Moscow, 336 pp. [In Russian].
- Babenko A, Kuznetsova N, Nakamori T, Shveenкова Y (2021) A review of *Pseudachorutes* Tullberg, 1871 (Collembola, Neanuridae) from the East Asia, with description of six new species. *Zootaxa* 4938 (4): 351-391. <https://doi.org/10.11646/zootaxa.4938.4.1>
- Babenko AB, Kuznetsova NA, Shveenкова YB (2019) New species of the genus *Anurida* (Collembola, Neanuridae) from the Far East of Russia. *Zoologicheskii Zhurnal* 98 (2): 130-148. <https://doi.org/10.1134/S004451341902003X>
- Bolger T, Arroyo J, Kenny J, Caplice M (2014) Hierarchical analysis of mite community structures in Irish forests—a study of the relative contribution of location, forest type and microhabitat. *Applied Soil Ecology* 83: 39-43. <https://doi.org/10.1016/j.apsoil.2013.06.004>
- Chernov YI, Lelei AS, Storozhenko SY (2011) Taxonomic diversity of insects in the Russian Far East. In: Lelei A (Ed.) Identification guide to the Russian Far East insects. Additional volume. Analysis of the fauna and the general index of names. Dal'nauka, Vladivostok, 552 pp. [In Russian]. [ISBN 978–5–8044–1106–1].
- Deharveng L, Bedos A, Weiner W (2011) Two new species of the genus *Leenurina* Najt & Weiner, 1992 (Collembola, Neanuridae, Capitanuriniinae) from Primorskii Krai (Russia). *ZooKeys* 115: 39-52. <https://doi.org/10.3897/zookeys.115.1464>
- Ghilarov M (1975) Sampling of small arthropods (microfauna) and nematodes. In: Ghilarov M (Ed.) Methods of soil zoological studies. Nauka, Moscow, 125 pp. [In Russian].
- Hopkin SP (1997) Biology of the springtails (Insecta: Collembola). Oxford University Press, Oxford, 330 pp.
- Huang C, Potapov M (2012) Taxonomy of the *Proisotoma* complex. IV. Notes on chaetotaxy of femur and description of new species of *Scutisotoma* and *Weberacantha* from Asia. *Zootaxa* 3333 (1). <https://doi.org/10.11646/zootaxa.3333.1.3>
- Jie D, Potapov M, Sokolova E (2011) Further study on the labial palp in the Isotomidae (Collembola) with reference to the genus *Heteroisotoma* Stach. *Zootaxa* 2991 (1). <https://doi.org/10.11646/zootaxa.2991.1.7>
- Jordana R, Potapov M, Baquero E (2011) New species of *Entomobryini* from Russia and Armenia (Collembola, Entomobryomorpha). *Soil Organisms* 83 (2): 221-248.

- Kutyreva LT (1979) Fauna and population of Collembola in soils of spruce–deciduous forests of Southern Primorye. In: Dolin V (Ed.) Fauna and ecology of invertebrates. Moscow Pedagogical State Institute, Moscow. [In Russian].
- Kutyreva LT (1984) Springtails in secondary oak forests of Primorye. In: Ghilarov M (Ed.) Fauna and ecology of Collembola. Nauka, Moscow. [In Russian].
- Kuznetsova NA, Saraeva AK (2018) Beta-diversity partitioning approach in soil zoology: A case of Collembola in pine forests. *Geoderma* 332: 142-152. <https://doi.org/10.1016/j.geoderma.2017.09.030>
- Kuznetsova NA, Bokova AI, Saraeva AK, Shveenikova YB (2019) Communities of Collembola in the forests of Southern Primorye as a benchmark of high diversity and organization complexity. *Biology Bulletin* 46 (5): 483-491. <https://doi.org/10.1134/s1062359019050066>
- Kuznetsova NA, Potapov MB, Shveenikova YB, Ivanova N (2021) Collembola of the relict forests of the Russian Far East. Sampling event dataset. 1.4. Moscow Pedagogical State University (MPGU). Release date: 2021-9-13. URL: <https://doi.org/10.15468/dyadwn>
- Lande R (1996) Statistics and partitioning of species diversity, and similarity among multiple communities. *Oikos* 76 (1). <https://doi.org/10.2307/3545743>
- Latham RE, Ricklefs RE (1993) Global patterns of tree species richness in moist forests: energy-diversity theory does not account for variation in species richness. *Oikos* 67: 325-333. <https://doi.org/10.2307/3545479>
- Marsh C, Ewers R (2012) A fractal-based sampling design for ecological surveys quantifying β -diversity. *Methods in Ecology and Evolution* 4 (1): 63-72. <https://doi.org/10.1111/j.2041-210x.2012.00256.x>
- Martynova E (1988) The order Collembola - springtails. In *Keys of insects of the Far East of the USSR. V. I. Primary-winged, ancient-winged, with incomplete metamorphosis*. Nauka, Leningrad, 452 pp. [In Russian].
- Petersen H, Luxton M (1982) A comparative analysis of soil fauna populations and their role in decomposition processes. *Oikos* 39 (3). <https://doi.org/10.2307/3544689>
- Pomorski RJ, Sveenkova YB (2006) New genus with three new species of *Thalassaphorurini* (Collembola: Onychiuridae) from Russian Far East. *Insect Systematics & Evolution* 37: 191-196. <https://doi.org/10.1163/187631206788831092>
- Potapov M (2001) Synopses on Palaearctic Collembola. V.3. Isotomidae. Senckenberg Museum of Natural History Görlitz, 603 pp.
- Potapov M, Kuznetsova N (2011) Methods of research of microarthropod communities: a manual for students and postgraduates. *Tovarishchestvo Nauchnykh Izdaniy KMK*, Moscow, 77 pp. [In Russian].
- Saraeva AK, Potapov MB, Kuznetsova NA (2015) Different-scale distribution of Collembola in homogenous ground vegetation: sphagnum moss. *Entomological Review* 95 (5): 557-577. <https://doi.org/10.1134/s0013873815050012>
- Smolis A, Deharveng L, Bedos A (2012) Description of two new species of the genus *Micranurida* Börner, 1901 (Collembola: Neanuridae, Pseudachorutinae) from Russia, with notes on the genus *Lanzhotia* Rusek, 1985. *Entomologica Fennica* 23: 149-154. <https://doi.org/10.33338/ef.84592>
- Smolis A, Deharveng L (2015) Diversity of *Paranura* Axelson, 1902 (Collembola: Neanuridae: Neanurinae) in Pacific region of Russia and United States. *Zootaxa* 4033 (2). <https://doi.org/10.11646/zootaxa.4033.2.2>

- Tsyganov AN, Komarov AA, Mitchell EA, Shimano S, Smirnova OV, Aleynikov AA, Mazei YA (2014) Additive partitioning of testate amoeba species diversity across habitat hierarchy within the pristine southern taiga landscape (Pechora-Ilych Biosphere Reserve, Russia). *European Journal of Protistology* 51: 42-54. <https://doi.org/10.1016/j.ejop.2014.11.003>
- Wieczorek J, Bloom D, Guralnick R, Blum S, Döring M, Giovanni R, Robertson T, Vieglais D (2012) Darwin Core: an evolving community-developed biodiversity data standard. *PLOS One* 7 (1). <https://doi.org/10.1371/journal.pone.0029715>
- Xin S, Shveenkova YB, Xie Z, Babenko AB (2019) New *Oligaphorura* species (Collembola: Onychiuridae) from the forests of East Asia. *Zootaxa* 4661 (2): 256-270. <https://doi.org/10.11646/zootaxa.4661.2.2>



Figure 1.
Plastic containers



Figure 2.
One plastic container with a core



Figure 3.

Above-soil cover in broad-leaf – cedar pine forest in Ussuriiskii natural reserve (plot “Turova”), photo by N. Kuznetsova.



Figure 4.

Above-soil cover in broad-leaf – cedar pine forest in Ussuriiskii natural reserve (plot “Turova”), photo by N. Kuznetsova.



Figure 5.

Mountain fir–hornbeam forest in Ussuriiskii natural reserve (plot “Grabovaya”), photo by N. Kuznetsova.



Figure 6.

Above-soil cover in mountain coniferous forest with *Rhododendrom fauriei* (plot "Fauri") in Sikhote-Alinskii natural reserve, photo by A. Geras'kina.



Figure 7.

Mountain coniferous forest with *Rhododendrom fauriei* (plot "Fauri") in Sikhote-Alinskii natural reserve, photo by R. Naumenko.



Figure 8.
Valley mixed forest (plot “Chuguev”), photo by A. Geras'kina.



Figure 9.
Litter in plot “Chuguev”, photo by A. Geras'kina.



Figure 10.

Geographic coverage. Study areas in the Primorsky Krai (Kuznetsova et al. 2021, doi.org/10.15468/dyadwn).




Figure 11.
Number of Collembola species in a core, on one square metre, in one sampling plot.