

Distribution of vascular plants north of Lake Baikal: a new, open access dataset

Denis V. Sandanov[‡], Elena P. Brianskaia[‡], Eduard A. Batotsyrenov[§]

[‡] Institute of General and Experimental Biology SB RAS, Ulan-Ude, Russia

[§] Baikal Institute of Nature Management SB RAS, Ulan-Ude, Russia

Corresponding author: Denis V. Sandanov (denis.sandanov@gmail.com)

Academic editor: Alexander Sennikov

Abstract

Background

The area north of Lake Baikal has been poorly studied. Moreover, most of the studies conducted in this region were focused on mountain ridges or river valleys. This region includes a part of Baikal-Amur Mainline (BAM), a broad-gauge railway in the centre of Siberia, Russia. The railway is an alternative route of the Trans-Siberian Railway; BAM starts in southern Siberia (Taishet station of Irkutsk Oblast), passes through the northern part of Lake Baikal and finishes in the Russian Far East (Sovetskaya Gavan station of Khabarovsk Krai). BAM has four connections with the Trans-Siberian Railway and is the centre of economic development for many regions of Russia. Maya Ivanova and Alexandr Chepurnov summarised the existing floristic information for this region in detailed species distribution maps which they published in the book “Flora of the western part of developing regions of Baikal-Amur Mainline (BAM)” (1983). After publishing this book, very few floristic studies have been performed in the study region. All available botanical information is still accumulated in a number of printed papers or books with limited circulation, which are not widely known to the international scientific community.

New information

We have digitised the point distribution maps from the book of Ivanova and Chepurnov and georeferenced all occurrence and sampling localities. The resulting dataset includes 9972 occurrences for 770 vascular plant species and subspecies from the area north of Lake Baikal. Additionally, the dataset includes information on the distribution of 43 rare and endangered species with 366 occurrences. From our point of view, the dataset makes a contribution to the global biodiversity data mobilisation, providing plant species distribution data for such a remote mountainous area.

Keywords

floristic studies, species occurrences, printed maps, Baikal-Amur Mainline, Russia, Siberia

Introduction

Lake Baikal and its surrounding terrestrial ecosystems have recently undergone diverse climate change processes (Moore et al. 2009). The surface air temperature has warmed by 1.2°C during the last century; temperature increases have been observed in all seasons, but are greatest in winter and spring (Shimaraev et al. 2002). These changes are reflected in shifts in the phenology of vascular plants in the Barguzinsky Nature Reserve over the last 40 years with significant advances of spring events and delays of those associated with senescence in autumn (Rosbakh et al. 2021). The purpose of our work is to make available essential baseline data for analysing the manner in which the flora and vegetation around the northern end of Lake Baikal (Fig. 1) is responding to the global climate change. The northern part of this region includes a section of the Baikal-Amur Mainline (BAM), a broad-gauge railway line that goes through the centre of Siberia, which has led to the economic development along its route.

The mountains and river valleys around the northern part of Lake Baikal have been covered by a few botanical studies (Malyshev 1972, Tyulina 1976, Tyulina 1981, Ivanova and Chepurnov 1983). Ivanova and Chepurnov (1983) summarised previous floristic surveys and herbarium collections from the western part of BAM. They listed 1352 species and subspecies from 428 genera and 97 families occurring in the region. Species distribution maps for a larger area were published previously in "Alpine flora of Stanovoye Nagorye Upland" (Malyshev 1972) and "Flora of Central Siberia" (Malyshev and Peshkova 1979a, Malyshev and Peshkova 1979b).

Maps from these monographs have been critically analysed (Ivanova and Chepurnov 1983). In some cases, herbarium specimens were verified for clarification of species localities. New records of vascular species in the study area were added, based on specimens collected by N.S. Vodopyanova, M.M. Ivanova, Yu.N. Petrochenko, A.A. Chepurnov, M.G. Azovsky, V.V. Telyatiev and other researchers who worked at BAM (Table 1). These botanical studies were summarised in a book on the flora of the Baikal Siberian Region and its genesis (Malyshev and Peshkova 1984).

Recent botanical studies do not cover the whole area to the north of Lake Baikal. Even the publications that cover some parts of it (e.g. Upper Chara Depression (Garaschenko 1993); Kodar and Udokan Ranges (Shvetsova 2000); around Lake Baikal (Ivanova 2003); Vitimsky Nature Reserve (Czeczjotkin 1986, Czeczjotkin 1989, Chechyotkina 1993, Bardunov 2005, Chechyotkina 2010); and Baikalo-Lensky Nature Reserve (Stepantsova 2009, Stepantsova 2010, Stepantsova and Zheleznaya 2016) do not include distribution maps, being focused primarily on species lists or new species records in particular. Some

recent publications do, however, provide geographical coordinates of species occurrences from satellite navigators.

The eastern part of BAM, the Chara floristic region, has been affected by large-scale human activities: copper mining in the Udokan Range, gold mining in the Olekma-Chara highland and proposals for extensions of BAM.

The development of portable satellite trackers has made incorporating georeference information into collection and observation records common. Our purpose for digitising the maps published in Ivanova & Chepurnov (1983) and freely sharing the resulting species occurrences is to provide the baseline data that will aid all those interested in the BAM's flora and in mapping its changes over time.

General description

Purpose: Digitising the vascular plant species distribution maps covering the western part of Baikal-Amur Mainline, which are published in Ivanova & Chepurnov (1983). This source contains crucial information on species distributions in the northern part of Lake Baikal, which is a less studied area of the Baikal Siberian Region. Other distribution maps currently available for this territory have a larger scale and many plant species are represented only by a few occurrences there.

Project description

Title: №121030900138-8 «Biota of terrestrial ecosystems of Baikal Region: composition, structure, eco-geographic patterns»

Personnel: Denis Sandanov, Elena Brianskaia

Study area description: Baikal Region, Russia

Design description: The project is designed to benefit many different areas of study, such as: plant taxonomy, floristics, vegetation science, plant biology and population ecology, fauna and ecology of insects, ecology and geography of vertebrates.

Funding: Russian Federal Budget

Sampling methods

Description: The study area is situated on the northern edges of three regions of Russia: Irkutsk Oblast, Republic of Buryatia and Zabaikalsky Krai. Some of the species occurrences at the north-western part of Lake Baikal, including Baikal Range, are now included in the Baikalo-Lensky Nature Reserve. The eastern part of the study area is legally protected in the Barguzinsky and Dzherginsky Nature Reserves, Zabaikalsky

National Park and Frolikhinsky Sanctuary. The north-eastern part of Irkutsk Oblast includes the Vitimsky Nature Reserve (Fig. 1).

Sampling description: In total, 770 maps were scanned from the book. Using the position of Lake Baikal and neighbouring rivers, we defined the projection of the maps (Fig. 2). We used a similar technique as employed for our previous dataset describing the distributions of endemic alpine species of northern Asia (Brianskaia et al. 2021). All the maps were adjusted to the same size and horizontal position in order to obtain standardised images of the maps. Digitalisation was performed in QGIS 3.10 software with the help of its georeferencing tools. The most accurate projected coordinate system was Asia North Albers Equal Area Conic. The water bodies shapefile was downloaded from the open source (https://vsegei.ru/ru/info/ggk_1000ns/) in scale 1:1 Mio. The river drainage shapefile fits very well with the original paper maps, but there were problems with the shape of Lake Baikal, especially in its northern part. In such cases, species distribution maps were georeferenced by snapping control points to the destination vector shapefile, which was the contour of Lake Baikal. We used control points (usually 5-8) to link maps to the destination shapefile, which resulted in the transformation of the maps according to the spatial projection of the destination features (WGS 1984). Subsequently, species distribution locations were digitised from each map. Coordinates of each location were calculated in the attribute table (Sandanov and Brianskaia 2021).

Quality control: We performed the final examination of the digitised species distribution maps in QGIS 3.10. For each species, we compared the output digitised occurrences with the original maps in order to check missing distribution records. The majority of occurrences (98%) matched consistently with the printed maps. Other 187 distribution records were manually adjusted for better matching with their habitats. These records mostly belong to the psammophytes occurring along the shoreline of Lake Baikal, especially at its northern part (Fig. 3). The diameter of points denoting the species occurrences is equal to 16 km. In this process, digitised localities of the psammophyte plants were moved closer to the shoreline. Taking this procedure into account, we estimate the coordinate uncertainty as 20 km for all the species in this study, taken as a matter of precaution.

Geographic coverage

Description: The study area includes the western part of BAM from Ust-Kut Town in the west and the Chara Depression in the east. It is a mountainous region involving several ranges of Stanovoy Highlands (Upper Angara, North Muya, South Muya, Kodar, Udokan), Baikal and Barguzin Ranges (Fig. 1). The main river of the study area is the Lena River. One of its southern tributaries is the Vitim River, which flows to the Lena from the north-east of Lake Baikal. The Vitim has tributaries draining the area, the Muya, Mamakan and Mama tributaries from the west and the Kalar and Kalakan tributaries from the east. The two major rivers on the west side of the region originate in the western part of the Stanovoy Highlands: the Upper Angara River that flows into the northern end of Lake Baikal and the Chaya River that flows into the Lena River.

The territory is divided into several floristic regions (Ivanova and Chepurnov 1983) (Fig. 1):

1. Lena River Region. Basins of Rivers Lena and Kirenga within Ust-Kutskii and Kazachinsko-Lenskii Districts of Irkutsk Oblast.

2. Baikal Region. This includes the north-western and north-eastern parts of Lake Baikal and is surrounded by the Baikal and Barguzin Ranges. It belongs to Severobaikalskii District of the Republic of Buryatia and partially to Kazachinsko-Lenskii District of Irkutsk Oblast.

3. Upper Angara Depression with Upper Angara Range. It is included in Severobaikalskii District of the Republic of Buryatia.

4. Muya-Kuanda Depression, which is bordered by the North Muya and South Muya Ranges. Most of this region is situated in Severobaikalskii and Bauntovskii Districts of the Republic of Buryatia, but the lowlands bordered on their right by the Vitim River belong to Kalarskii District of Zabaikalsky Krai.

5. Chara Depression with the Kodar, Udokan and Kalarsky Ranges. This is the eastern part of the Stanovoy Highlands. It does not include the Kuanda River, which is considered part of the Muya-Kuanda Depression floristic region. This region is located at the north of Kalarskii District of Zabaikalsky Krai.

We mapped all localities recorded in the original printed maps. Most of these localities were included within the study area, but a few lay outside the digitised floristic regions (Fig. 4) due to the presence of general distribution data in the original maps.

Coordinates: 53.48 and 58.05 Latitude; 121.29 and 104.23 Longitude.

Taxonomic coverage

Description: The dataset includes 770 species and subspecies of vascular plants with 9972 occurrences from 81 families and 266 genera. The whole list of the flora of this region includes 1352 species and subspecies. Therefore, the dataset contains more than a half of the flora (57%) because the distribution maps were provided for the most common species only. In reporting the data, we retained the family attributions used in the source to facilitate comparisons. The top 10 families include 58.9% of the taxa and 56.9% of the occurrences (Table 2). In the original floristic analysis, *Scrophulariaceae* appeared in the top 10 families (Ivanova and Chepurnov 1983), but it is replaced by *Apiaceae* in our dataset. *Scrophulariaceae* in the current circumscription is represented in the dataset by only one species (Sandanov and Brianskaia 2021), *Scrophularia incisa* (Fig. 3). Comparisons of percentages for all other families reveal further similarities between the complete floristic checklist and the species included in the dataset (Table 2). The comparisons testify that the dataset is representative for some part of the flora including families with high numbers of species. The digitised data can also be used for studies of the distribution patterns of key vascular plant species in the study region.

Our comparisons revealed that the list of top 10 genera was the same in the book and the dataset (Table 3). *Carex* and *Salix* are the leading genera in both lists. Standing next in the floristic list, *Potentilla* and *Artemisia* do not have distribution maps for the widely distributed species and that is why their position within the dataset is not so high. Other genera have similar positions as in the whole floristic checklist of the region.

The dataset contains information on the distribution of vascular plants species which are included in regional Red Data Books of the Baikal Siberian Region (Pronin 2013, Polyakov 2017, Trofimova 2020) (Table 4). These data are complementary to the recently-published dataset with occurrences of rare and endangered species of the Transbaikalia (Sandanov et al. 2021) and will be helpful in planning and implementing future conservation activities.

Temporal coverage

Notes: Dates of the specimen records used to prepare the printed maps ranged from 1912 to 1979 (Table 1).

Usage licence

Usage licence: Creative Commons Public Domain Waiver (CC-Zero)

IP rights notes: This work is licensed under a Creative Commons Attribution (CC-BY) 4.0 Licence.

Data resources

Data package title: Occurrences of vascular plants in the western part of Baikal-Amur Mainline.

Resource link: <https://www.gbif.org/dataset/1e7b25d0-ec44-4fa8-8338-6e38e7a11214>

Alternative identifiers: 1e7b25d0-ec44-4fa8-8338-6e38e7a11214; http://gbif.ru:8080/ipt/resource?r=vasc_plants_north_baikal

Number of data sets: 1

Data set name: Occurrences of vascular plants in the western part of Baikal-Amur Mainline.

Character set: UTF-8

Download URL: <https://doi.org/10.15468/a8c783>

Data format: Darwin Core Archive format.

Data format version: 1.15

Description: The northern part of Lake Baikal has been sparsely covered by botanical studies which were usually concentrated on mountain ridges or river valleys. The floristic information for this region with point distribution maps of vascular plant species is summarised in the book by M.M. Ivanova and A.A. Chepurinov “Flora of the western part of developing regions of Baikal-Amur Mainline (BAM)” (Ivanova and Chepurinov 1983). All available maps from this book have been digitised and occurrences of vascular plants were organised in a dedicated dataset. The dataset includes 9972 occurrences for 770 vascular plant species and subspecies occurring around the northern part of Lake Baikal (the western part of Baikal-Amur Mainline), which is a hard-to-access mountainous region.

Column label	Column description
occurrenceID	An identifier for the record, unique within this dataset. An abbreviation in the identifier' number (IVBAM).
basisOfRecord	The specific nature of the data record in standard label of the Darwin Core classes: HumanObservation.
scientificName	The full scientific name of the species as recorded in the book by M.M. Ivanova and A.A. Chepurinov (1983) “Flora of the western part of developing regions of Baikal-Amur Mainline (BAM)”.
genus	The full scientific name of the genus in which the taxon is classified.
specificEpithet	The name of the species epithet as recorded in the book by M.M. Ivanova and A.A. Chepurinov (1983) “Flora of the western part of developing regions of Baikal-Amur Mainline (BAM)”.
infraspecificEpithet	The name of the lowest or terminal infraspecific epithet as recorded in the book by M.M. Ivanova and A.A. Chepurinov (1983) “Flora of the western part of developing regions of Baikal-Amur Mainline (BAM)”.
taxonRank	The taxonomic rank of the most specific name in the scientificName.
acceptedNameUsage	The full name, with authorship and date information, if known, of accepted taxon.
kingdom	The full scientific name of the kingdom in which the taxon is classified.
phylum	The full scientific name of the phylum or division in which the taxon is classified
class	The full scientific name of the class in which the taxon is classified.
order	The full scientific name of the order in which the taxon is classified.
family	The full scientific name of the family in which the taxon is classified.
decimalLatitude	The geographic latitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic centre of a Location
decimalLongitude	The geographic longitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic centre of a Location.

georeferencedBy	A list of persons who determined the georeference (spatial representation) for the Location.
geodeticDatum	The ellipsoid, geodetic datum or spatial reference system (SRS) upon which the geographic coordinates given in decimalLatitude and decimalLongitude are based.
eventDate	The date-time or interval during which an Event occurred. This is the publication date of the book by M.M. Ivanova and A.A. Chepurnov (1983) "Flora of the western part of developing regions of Baikal-Amur Mainline (BAM)".
coordinateUncertaintyInMetres	The horizontal distance (in metres) from the given decimalLatitude and decimalLongitude describing the smallest circle containing the whole of the Location.
verbatimCoordinateSystem	The coordinate format for the verbatimLatitude and verbatimLongitude or the verbatimCoordinates of the Location.
higherGeography	A list of geographic names less specific than the information captured in the locality term.
continent	The name of the continent in which the Location occurs
country	The name of the country or major administrative unit in which the Location occurs.
countryCode	The standard code for the country in which the Location occurs.
type	The nature or genre of the resource.
language	A language of the resource.
licence	A legal document giving official permission to do something with the resource.
associatedReferences	A list (concatenated and separated) of identifiers (publication, bibliographic reference, global unique identifier, URI) of literature associated with the Occurrence.
taxonRemarks	Comments or notes about the taxon or name. Usually contains notes about definition of the taxon "sensu lato" or "sensu stricto" as recorded in the book by M.M. Ivanova and A.A. Chepurnov (1983) "Flora of the western part of developing regions of Baikal-Amur Mainline (BAM)".

Acknowledgements

The authors are very thankful to Dr. Mary Barkworth (Utah State Univeristy, Department of Biology) who kindly helped in proofreading and editing the English in the manuscript.

The research of Denis Sandanov and Elena Brianskaia was funded by the Russian Federal Budget (project number 121030900138-8) and partially by RFBR (project number 20-45-380009). The work of Eduard Batotsyrenov was funded within the budget topic for Baikal Institute of Nature Management SB RAS (project number AAAA-A21-121011990023-1).

References

- Bardunov LV (Ed.) (2005) Биота Витимского заповедника. Флора. [Biota of Vitim Nature Reserve. Flora.]. Academic Publishing House “Geo”, Novosibirsk, 207 pp. [In Russian]. [ISBN 5-9747-0012-0]
- Brianskaia E, Sandanov D, Li Y, Wang Z (2021) Distribution of alpine endemic plants of northern Asia: a dataset. Biodiversity Data Journal 9 <https://doi.org/10.3897/bdj.9.e75348>
- Chechyotkina LG (1993) Floristic findings in Stanovoye Nagorye (East Siberia). Botanical Journal 78 (2): 125-126. [In Russian].
- Chechyotkina LG (2010) Geographical analysis of the flora of vascular plants of State Nature Reserve “Vitimskiy” (Stanovoye and Patomskoye highlands). Problems of studying and preserving the plant world of Eurasia, Irkutsk, 13-16 September 2010. Institute of Geography SB RAS, Irkutsk, 205-208 pp. [In Russian]. [ISBN 978-5-94797-152-1].
- Czeczotkin EV (1986) A supplement to the flora of Stanovoye Nagorye Mountains. Botanical Journal 71 (11): 1562-1564. [In Russian].
- Czeczotkin EV (1989) On some plant species new to and rare in the northern Baikal Region. Botanical Journal 74 (7): 1051-1054. [In Russian].
- Garaschenko AV (1993) Флора и растительность Верхнечарской котловины (Северное Забайкалье). [Flora and vegetation of Upper Chara Depression (Northern Transbaikalia)]. Nauka, Siberian Branch, Novosibirsk, 280 pp. [In Russian]. [ISBN 5-02-03546-4]
- Ivanova MM, Chernunov AA (1983) Флора западного участка районов освоения БАМ. [Flora of the western part of developing regions of Baikal-Amur Mainline (BAM)]. Nauka, Siberian Branch, Novosibirsk, 223 pp. [In Russian].
- Ivanova MM (2003) Floristic findings in the vicinities of Lake Baikal and neighboring territories. Turczaninowia 6 (2): 51-78. [In Russian].
- Malyshev LI (Ed.) (1972) Высокогорная флора Станового Наргорья. [Alpine flora of the Stanovoye Nagorye uplands]. Nauka, Siberian Branch, Novosibirsk, 272 pp. [In Russian].
- Malyshev LI, Peshkova GA (Eds) (1979a) Флора Центральной Сибири. [Flora of Central Siberia]. 1. Nauka, Siberian Branch, Novosibirsk, 536 pp. [In Russian].
- Malyshev LI, Peshkova GA (Eds) (1979b) Флора Центральной Сибири. [Flora of Central Siberia]. 2. Nauka, Siberian Branch, Novosibirsk, 1048 pp. [In Russian].
- Malyshev LI, Peshkova GA (1984) Особенности и генезис флоры Сибири (Предбайкалье и Забайкалье). [Peculiarities and genesis of the Siberian flora (Cis-Baikal and Trans-Baikal Regions)]. Nauka, Siberian Branch, Novosibirsk, 265 pp. [In Russian].
- Moore MV, Hampton SE, Izmet'eva LR, Silow EA, Peshkova EV, Pavlov BK (2009) Climate change and the world's “Sacred Sea”— Lake Baikal, Siberia. BioScience 59 (5): 405-417. <https://doi.org/10.1525/bio.2009.59.5.8>
- Polyakov OA (Ed.) (2017) Красная книга Забайкальского края. Растения. [Red Data Book of Zabaikalsky Krai. Plants]. ООО Dom Mira, Novosibirsk, 384 pp. [In Russian]. [ISBN 978-5-4364-0042-6]

- Pronin NM (Ed.) (2013) Красная книга Республики Бурятия: Редкие и находящиеся под угрозой исчезновения виды животных, растений и грибов. [Red Data Book of Republic of Buryatia: Rare and endangered species of animals, plants, and fungi]. Buryat Scientific Center Publishing House, Ulan-Ude, 688 pp. [In Russian]. [ISBN 978-5-7925-0400-4]
- Rosbakh S, Hartig F, Sandanov DV, Bukharova EV, Miller TK, Primack RB (2021) Siberian plants shift their phenology in response to climate change. *Global Change Biology* 27 (18): 4435-4448. <https://doi.org/10.1111/gcb.15744>
- Sandanov D, Brianskaia E (2021) Occurrences of vascular plants in the western part of Baikal-Amur Mainline. 1.15. GBIF. Release date: 2021-12-07. URL: <https://doi.org/10.15468/a8c783>
- Sandanov D, Brianskaia E, Dugarova A (2021) Rare vascular plant species of Transbaikalia (Republic of Buryatia and Zabaikalsky Krai), Russia. 1.7. GBIF. Release date: 2021-11-25. URL: <https://doi.org/10.15468/rh8s6n>
- Shimaraev MN, Kuimova LN, Sinyukovich VN, Tsekhanovskii VV (2002) Manifestation of global climatic changes in Lake Baikal during the 20th century. *Doklady Earth Sciences* 383A: 288-291.
- Shvetsova VN (2000) Сравнительный анализ флоры сосудистых растений Кодаро-Удоканского горного района. [Comparative analysis of the vascular plants flora of Kodar-Udokan mountain region]. Institute of Plant and Animal Ecology, Ekaterinburg, 194 pp. [In Russian].
- Stepantsova NV (2009) Флора сосудистых растений Байкало-Ленского государственного природного заповедника. [Flora of the vascular plants of the Baikal-Lena State Nature Reserve]. Central Siberian Botanical Garden SB RAS, Novosibirsk, 365 pp. [In Russian].
- Stepantsova NV (2010) Additions to the "Flora of Siberia" in Lena-Katanga floristic area of Irkutsk region. *Botanical Journal* 95 (7): 992-100. [In Russian].
- Stepantsova NV, Zheleznaya EL (2016) New and rare plant species of the Baikal-Lena Reserve flora (Irkutsk Region). *Turczaninowia* 19 (2): 70-76. [In Russian]. <https://doi.org/10.14258/turczaninowia.19.2.9>
- Trofimova SM (Ed.) (2020) Красная книга Иркутской области. [Red Data Book of Irkutsk Oblast]. Republic Publishing House, Ulan-Ude, 552 pp. [In Russian]. [ISBN 978-5-91407-216-9]
- Tyulina LN (1976) Влажный прибайкальский тип поясности растительности. [Humid Baikal type of vegetation belts]. Nauka, Siberian Branch, Novosibirsk, 319 pp. [In Russian].
- Tyulina LN (1981) Растительность южной части Баргузинского хребта. [Vegetation of the southern part of the Barguzin Range]. Nauka, Siberian Branch, Novosibirsk, 85 pp. [In Russian].




Figure 1.

General map of the study area.

The information on region topography, water bodies, floristic regions and protected areas are combined in one map. Floristic regions (according to Ivanova and Chepurinov 1983) are marked by colours.




Figure 2.

A georeferenced map (the distribution of *Matteuccia struthiopteris* (L.) Tod.) showing the study area.




Figure 3.

Georeferencing the distribution map of *Scrophularia incisa* in QGIS 3.10.

Overlay is the GIS shapefiles (denoted by colours), background is the original printed map (black and white).




Figure 4.
Location of vascular plants occurrences within the study area.

Table 1.

History of botanical studies in the northern part of Lake Baikal (from 1912 till 1979).

Territory	Botanists	Years of study
Ust-Kutskii District	Nomokonov L.I., Reshikov M.A.	1950
	Popov M.G. and co-authors	1951
Kazachinsko-Lenskii District	Alexandrov P.	1912
	Belov A.V., Garashenko A.V.	1967
	Azovsky M.G.	1979
Baikal Range and north-western part of Lake Baikal	Popov M.G., Malyshev L.I.	1955
	Tyulina L.N., Ivanova M.M.	1958
	Molozhnikov V.N., Granina G.T.	1966
	Petrochenko Yu.N.	1966-1967
	Malyshev L.I.	1967
	Makryi T.V. Telyatiev V.V.	1974 1979
Barguzin Range and north-eastern part of Lake Baikal	Tyulina L.N.	1939-1961
	Popov M.G. and coauthors	1954
	Petrochenko Yu.N.	1963, 1966
	Ivanova M.M.	1965
	Malyshev L.I.	1966-1968
Upper Angara Depression	Sukachev V.N., Poplavskaya G.I., Shipchinskii N.V.	1912 1955
	Malyshev L.I.	1963, 1966
	Petrochenko Yu.N.	1975
	Molozhnikov V.N.	1976
	Ivanova M.M.	1977
	Azovsky M.G.	
Upper Angara Range	Petrochenko Yu.N.	1963, 1966
	Malyshev L.I.	1966
	Azovsky M.G.	1977, 1979
Muya-Kuanda Depression	Malyshev L.I., Petrochenko Yu.N.	1965
	Ivanova M.M.	1967
	Petrochenko Yu.N.	1976
	Chepurnov A.A.	1977
	Azovsky M.G.	1978
North Muya Range	Petrochenko Yu.N.	1963
	Malyshev L.I., Petrochenko Yu.N.	1965
	Ivanova M.M.	1967
	Malyshev L.I., Petrochenko Yu.N.	1968
South Muya Range	Malyshev L.I., Petrochenko Yu.N.	1966
	Ivanova M.M., Andralaitis S.Yu.	1966-1967
	Petrochenko Yu.N.	1968
Chara Depression	Mikheev V.S.	1963-1964
	Garashenko A.V.	1964,
	Malyshev L.I., Petrochenko Yu.N.	1975-1976
	Vodopyanova N.S.	1964
	Chepurnov A.A.	1967
		1978

Kodar Range	Malyshev L.I., Petrochenko Yu.N. Vodopyanova N.S. Chepurnov A.A.	1964 1967 1978
Udokan Range	Vodopyanova N.S. Ivanova M.M., Bardunov L.V. Chepurnov A.A.	1964-1967 1969 1978

Table 2.

Taxonomic distribution of vascular plants in the northern part of Lake Baikal. Families are listed in descending order of the number of species and subspecies.

No.	Family	No. of species and subspecies	No. of records	Percent of species and subspecies (dataset)	Percent of species and subspecies (Ivanova and Chepurinov 1983)
1	Poaceae	95	1104	12.3	9.5
2	Cyperaceae	84	1075	10.9	9.3
3	Asteraceae	64	661	8.3	10.6
4	Caryophyllaceae	38	495	4.9	4.2
5	Ranunculaceae	33	411	4.3	5.2
6	Rosaceae	32	414	4.2	5.5
7	Salicaceae	31	728	4	3
8	Brassicaceae	29	272	3.8	4.1
9	Fabaceae	27	269	3.5	4.6
10	Apiaceae	20	249	2.6	2.5
11	Juncaceae	17	256	2.2	1.6
12	Orchidaceae	17	116	2.2	1.7
13	Orobanchaceae	17	216	2.2	0.2
14	Saxifragaceae	17	424	2.2	2.4
15	Ericaceae	16	299	2.1	1.3
16	Boraginaceae	14	125	1.8	1.4
17	Polygonaceae	12	210	1.6	1.9
18	Lamiaceae	11	98	1.4	2.2
19	Potamogetonaceae	11	76	1.4	1.1
20	Violaceae	9	125	1.2	1.1
21	Lycopodiaceae	8	190	1	0.6
22	Betulaceae	7	183	0.9	1.2
23	Gentianaceae	7	102	0.9	1.5
24	Caprifoliaceae	6	116	0.8	0.4
25	Cystopteridaceae	6	120	0.8	0.4
26	Papaveraceae	6	45	0.8	0.9
27	Plantaginaceae	6	41	0.8	0.3
28	Typhaceae	6	45	0.8	0.1
29	Amaryllidaceae	5	39	0.7	0.4
30	Campanulaceae	5	117	0.7	0.4
31	Crassulaceae	5	76	0.7	0.6

32	Geraniaceae	5	52	0.7	0.7
33	Liliaceae	5	54	0.7	2.1
34	Polemoniaceae	5	77	0.7	0.4
35	Primulaceae	5	49	0.7	1
36	Dryopteridaceae	4	58	0.5	0.4
37	Grossulariaceae	4	73	0.5	0.3
38	Lentibulariaceae	4	34	0.5	0.5
39	Araceae	3	12	0.4	0.1
40	Athyriaceae	3	52	0.4	0.7
41	Iridaceae	3	14	0.4	0.2
42	Melanthiaceae	3	51	0.4	0.2
43	Montiaceae	3	29	0.4	0.2
44	Nymphaeaceae	3	20	0.4	0.3
45	Onagraceae	3	54	0.4	0.4
46	Pinaceae	3	70	0.4	0.5
47	Rubiaceae	3	17	0.4	0.6
48	Selaginellaceae	3	45	0.4	0.3
49	Tofieldiaceae	3	51	0.4	0.2
50	Adoxaceae	2	21	0.3	0.1
51	Alismataceae	2	20	0.3	0.2
52	Amaranthaceae	2	10	0.3	0.1
53	Aspleniaceae	2	13	0.3	0.2
54	Cupressaceae	2	39	0.3	0.1
55	Juncaginaceae	2	18	0.3	0.2
56	Ophioglossaceae	2	19	0.3	0.2
57	Polygalaceae	2	13	0.3	0.1
58	Pteridaceae	2	39	0.3	0.3
59	Thelypteridaceae	2	18	0.3	0.1
60	Woodsiaceae	2	36	0.3	0.1
61	Acoraceae	1	4	0.1	0.1
62	Asparagaceae	1	28	0.1	0.1
63	Ceratophyllaceae	1	6	0.1	0.1
64	Dennstaedtiaceae	1	6	0.1	0.1
65	Diapensiaceae	1	29	0.1	0.1
66	Droseraceae	1	6	0.1	0.1
67	Ephedraceae	1	12	0.1	0.1
68	Equisetaceae	1	29	0.1	0.6

69	Euphorbiaceae	1	32	0.1	0.1
70	Haloragaceae	1	6	0.1	0.1
71	Hydrocharitaceae	1	5	0.1	0.1
72	Isoetaceae	1	6	0.1	0.1
73	Linaceae	1	6	0.1	0.1
74	Lythraceae	1	2	0.1	0.1
75	Menyanthaceae	1	5	0.1	0.1
76	Onocleaceae	1	10	0.1	0.1
77	Polypodiaceae	1	11	0.1	0.1
78	Scheuchzeriaceae	1	3	0.1	0.1
79	Scrophulariaceae	1	3	0.1	3
80	Tamaricaceae	1	4	0.1	0.1
81	Urticaceae	1	4	0.1	0.4
Total		770	9972	-	-

Table 3.

Top 10 genera within the study area. Genera are listed in descending order of the number of species.

№	Genera	No. of species and subspecies (dataset)	No. of records (dataset)	Percent of species and subspecies (dataset)	Percent of species and subspecies (Ivanova and Chepurmov 1983)
1	<i>Carex</i>	66	852	24.8	23.8
2	<i>Salix</i>	30	708	11.3	8.6
3	<i>Poa</i>	16	251	5.6	4.9
4	<i>Saxifraga</i>	14	382	5.3	4.7
5	<i>Potentilla</i>	13	175	4.9	7.0
6	<i>Pedicularis</i>	10	140	3.8	4.2
7	<i>Astragalus</i>	9	127	3.4	4.0
8	<i>Oxytropis</i>	9	55	3.4	4.2
9	<i>Artemisia</i>	7	93	2.6	6.3
10	<i>Polygonum</i>	6	114	2.3	4.2

Table 4.

The list of vascular plant species included in regional Red Data Books of the Baikal Siberian Region.

Species	Number of records	Region, where the species is considered rare and endangered
<i>Arctous alpina</i>	8	Zabaikalsky Krai
<i>Atragene ochotensis</i>	9	Zabaikalsky Krai
<i>Borodinia macrophylla</i>	9	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Calypso bulbosa</i>	4	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Caragana jubata</i>	6	Republic of Buryatia, Zabaikalsky Krai
<i>Carex malyshchevii</i>	6	Zabaikalsky Krai
<i>Carex sabulosa</i>	3	Zabaikalsky Krai
<i>Cotoneaster neo-popovii</i>	4	Irkutsk Oblast, Republic of Buryatia
<i>Cotoneaster tjulinae</i>	5	Republic of Buryatia
<i>Craniospermum subvillosum</i>	9	Irkutsk Oblast, Republic of Buryatia
<i>Cypripedium calceolus</i>	5	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Cypripedium guttatum</i>	11	Zabaikalsky Krai
<i>Cypripedium macranthon</i>	4	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Deschampsia turczaninowii</i>	8	Irkutsk Oblast, Republic of Buryatia
<i>Epipactis helleborine</i>	2	Irkutsk Oblast, Republic of Buryatia
<i>Epipogium aphyllum</i>	6	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Gastrolychnis popovii</i>	4	Republic of Buryatia
<i>Isoetes setacea</i>	6	Republic of Buryatia, Zabaikalsky Krai
<i>Lilium pilosiusculum</i>	8	Zabaikalsky Krai
<i>Lilium pensylvanicum</i>	7	Zabaikalsky Krai
<i>Lilium pumilum</i>	3	Zabaikalsky Krai
<i>Listera cordata</i>	3	Republic of Buryatia
<i>Lycopodium juniperoideum</i>	6	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Mertensia serrulata</i>	6	Republic of Buryatia
<i>Neottia camtschatea</i>	3	Irkutsk Oblast, Republic of Buryatia
<i>Neottianthe cucullata</i>	2	Republic of Buryatia
<i>Nymphaea candida</i>	2	Irkutsk Oblast, Republic of Buryatia
<i>Nymphaea tetragona</i>	12	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Oxytropis kodarensis</i>	4	Irkutsk Oblast, Zabaikalsky Krai
<i>Phlojodicarpus villosus</i>	10	Zabaikalsky Krai
<i>Phyllodoce coerulea</i>	2	Zabaikalsky Krai
<i>Potentilla adenotricha</i>	5	Zabaikalsky Krai
<i>Pulsatilla ajanensis</i>	5	Republic of Buryatia, Zabaikalsky Krai

<i>Rhodiola quadrifida</i>	28	Zabaikalsky Krai
<i>Rhodiola rosea</i>	35	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Rhododendron adamsii</i>	23	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Rhododendron aureum</i>	45	Zabaikalsky Krai
<i>Rhododendron redowskianum</i>	35	Zabaikalsky Krai
<i>Rhynchospora alba</i>	3	Irkutsk Oblast, Republic of Buryatia, Zabaikalsky Krai
<i>Ribes dikuscha</i>	4	Irkutsk Oblast, Republic of Buryatia
<i>Tillaea aquatica</i>	2	Irkutsk Oblast, Republic of Buryatia
<i>Trapa natans</i>	2	Zabaikalsky Krai
<i>Zannichellia repens</i>	2	Irkutsk Oblast, Republic of Buryatia
Total (43 species)	366	-