

New records, detailed distribution and abundance of rove-beetles (Insecta, Coleoptera, Staphylinidae) collected between 1990 and 2015 in Azores (Portugal) with an updated checklist

Paulo A. V. Borges[‡], Lucas Lamelas-Lopez[‡], Volker Assing^{†,§}, Michael Schülke[‡]

[‡] cE3c – Centre for Ecology, Evolution and Environmental Changes / Azorean Biodiversity Group and Universidade dos Açores, Rua Capitão João d'Ávila, São Pedro, 9700-042, Angra do Heroísmo, Azores, Portugal

[§] Gabelsbergerstraße 2, 30163, Hannover, Germany

[‡] Blankenfelder Straße 99, D-13127, Berlin, Germany

[†] Deceased author

Corresponding author: Paulo A. V. Borges (paulo.av.borges@uac.pt)

Academic editor: Pedro Cardoso

Abstract

Background

The dataset we present consists of an inventory compiling all records and knowledge about Staphylinidae (Insecta, Coleoptera) in the Azores and is part of a long-term monitoring performed between 1990 and 2015 in different habitat types of eight islands of the Azores Archipelago. Most samples come from the BALA project (Biodiversity of Arthropods from the Laurisilva of Azores) that sampled native forests in the Azores. Additional sampled habitats include exotic forests, intensive and semi-natural pasturelands, orchards, caves and lava flows. Most of the records (about 96.7%) were collected in standardised sampling campaigns, which included pitfall traps and beating transect protocols. Non-standardised records are based on hand-collecting and sifting, as well as cave, colour and malaise traps.

New information

We provide a long-term inventory of Staphylinidae (Insecta, Coleoptera) collected in the course of several standardised sampling campaigns and recorded with non-standardised methods. We collected a total of 10744 specimens belonging to 69 identified species of Staphylinidae, which represents 51% of the species known from the Azores Archipelago. Four endemic species were sampled, representing 40% of the known Azorean endemic species. From this dataset, seven species are new for the Azores: *Aleochara funebris*

Wollaston, 1864; *Amischa forcipata* Mulsant & Rey, 1873; *Bledius unicornis* (Germar, 1825); *Carpelimus troglodytes* (Erichson, 1840); *Cypha seminulum* (Erichson, 1839); *Paraphloeostiba gayndahensis* (MacLeay, 1871); *Tachyporus caucasicus* Kolenati, 1846. We also registered a total of 66 new island records for eight Azorean islands. This contribution continues a series of publications on the distribution and abundance of Azorean arthropods. We also provide an updated list of Azorean rove-beetles (Staphylinidae) that now includes 136 species, ten of them considered Azorean endemics.

Keywords

arthropods, Staphylinidae, Coleoptera, Azorean native forest, biodiversity, dataset, inventory, long term monitoring

Introduction

According to the latest list of rove-beetles from the Azores (Assing 2010), a total of 115 species and subspecies of Staphylinidae is known from the Azores, which is 21% of the known beetle fauna of the Archipelago (total number of beetle species in the Azores: 536 species and subspecies; see Oromí et al. 2010). Being one of the most diverse families of beetles, this group of mostly predatory insects is also known for its taxonomic complexity and frequent erroneous identifications (Barratt et al. 2003, Shaw and Solodovnikov 2016).

Despite the high species richness, very few endemic species have been recorded from the Archipelago; only ten species are confirmed as endemic in this paper (see Suppl. material 1). Several endemic staphylinids were discovered during the first scientific expeditions to the Azores (Crotch 1867, Fauvel 1900, Bernhauer 1936, Bernhauer 1940), others were found only recently (Israelson 1985, Pace 2004, Assing 2013). In addition to these works, the main contributors to the study of Azorean Staphylinidae were: Fauvel (1897), Fauvel (1898), Fauvel (1902), Méquignon (1942), Jarrige (1953), Smetana (1970), Brinck (1977), Serrano (1982), Israelson (1984), Gillerfors (1986), Gillerfors (1988), Serrano and Borges (1987), Borges and Serrano (1989) and Borges (1990), who added new records and provided taxonomic information.

However, in contrast to other Macaronesian archipelagos (see, for example, Assing and Schülke 2006), a recent revision of the Staphylinidae fauna of the Azores has not been attempted.

The main objective of this contribution is to provide an inventory of the Staphylinidae (Coleoptera, Insecta), based on a compilation of records from long-term monitoring sampling campaigns performed in eight islands of Azores between 1990 and 2015. This contribution is part of a series of publications on the distribution and abundance of Azorean arthropods (Borges et al. 2016, Borges et al. 2017).

General description

Purpose: The main objective is to provide an inventory of the Staphylinidae (Coleoptera, Insecta), based on a compilation of records from long-term monitoring sampling campaigns performed in eight islands of Azores between 1990 and 2015. In addition, an updated checklist of Azorean Staphylinidae is provided.

Additional information: This contribution is part of a series of publications on the distribution and abundance of Azorean arthropods (Borges et al. 2016, Borges et al. 2017) and updates the latest checklist of Azorean Staphylinidae (see Assing 2010).

Most data come from the following studies (see also Suppl. material 1): “1” – Confirmed occurrence, (see Assing 2010 list); BALA - records based on BALA protocol (see Borges et al. 2016; samples part of current study); LAND-USE (see Cardoso et al. 2009, samples part of current study); MACDIV (see Malumbres-Olarte et al. 2019); ASSING – Non-standardised samples of one of us (Volker Assing) (part of current study); Marcelino - Marcelino et al. (2021); INTERF (data from project INTERFRUTA part of current study); WET – from Borges et al. (2018); AGRO – see (Borges et al. 2021b); SLAM - Long term monitoring of Azorean forests using SLAM traps for Terceira Island (see Matthews et al. 2018, Borges et al. 2020, Tsafack et al. 2021); SLAM for other islands (unpublished data); ASSING 2013 (new species described in Assing 2013).

In this same Checklist of Azorean Staphylinidae (Suppl. material 1), we provide some relevant information related to taxonomic or nomenclature changes, splitting it into four columns “synonym”, “different combination”, “misidentification” and “emendation/misspelling”.

Project description

Title: Inventory of the Azorean rove-beetles (Insecta, Coleoptera, Staphylinidae)

Personnel: Leader: Paulo A. V. Borges,

Fieldwork BALA project (Gaspar et al. 2008, Borges et al. 2016): Ana Santos, Álvaro Vitorino, Ana Rodrigues, Anabela Arraiol, Annabella Borges, Artur Serrano, Carla Rego, Carlos Aguiar, Catarina Melo, Clara Gaspar, Emanuel Barcelos, Fernando Pereira, Francisco Dinis, François Rigal, Genage André, Hugo Mas, Isabel R. Amorim, João Amaral, João Moniz, Joaquín Hortal, Kostas Triantis, Lara Dinis, Luís Vieira, Paula Gonçalves, Pedro Cardoso, Sandra Jarroca, Sérgio Ribeiro.

Fieldwork for SLAM - Long term monitoring of Azorean forests using SLAM traps; see list at Costa and Borges (2021).

Fieldwork for LAND-USE project: see list at Cardoso et al. (2009).

Fieldwork for MACDIV: see list at Malumbres-Olarte et al. (2019).

Taxonomists: Michael Schülke, Paulo A. V. Borges, Volker Assing.

Study area description: The Azores Archipelago extends for 615 km and is located in the North Atlantic Ocean (37-40°N, 25-31°W), about 1600 km from Europe and 2200 km from North America (Fig. 1). The Archipelago is formed by nine main islands and some small islets, all of them of volcanic origin. The islands are divided into three main groups: the western group (Corvo and Flores), the central group (Faial, Pico, Graciosa, São Jorge and Terceira) and the eastern group (São Miguel and Santa Maria). The climate is temperate oceanic, with regular and abundant rainfall, high levels of relative humidity, above 95% on average in native forests and persistent winds, mainly during the winter and autumn seasons. Since Portuguese colonisation in the 15th century, the landscape of the Azores has been dramatically altered by replacing native forests with forest tree plantations, crops, pastures and urban areas. Currently, the native laurel forest comprises about 5% of the total surface of the Archipelago and has remained only at higher elevations and in inaccessible areas of the islands (Gaspar et al. 2008, Borges et al. 2020).

Design description: This inventory of Staphylinidae includes records of several standardised sampling campaigns and non-standardised observations, performed between 1990 and 2015 (Gaspar et al. 2008, Borges et al. 2016). Most of the records (about 96.7%) were collected in standardised sampling campaigns, which included pitfall and beating protocols. Non-standardised records are based on hand-collecting and sifting, as well as cave, colour and malaise traps. Additional non-standard records are based on cave, colour and malaise traps and collecting trips conducted by Volker Assing in Terceira, São Miguel and Santa Maria islands in 2013 and by Andreas Kleeberg in Pico and São Miguel in 2015. Collected samples were sorted and subsequently identified by an expert taxonomist in laboratory.

Funding: Species collecting was funded mostly by four projects:

“Reservas Florestais dos Açores: Cartografia e Inventariação dos Artrópodes Endémicos dos Açores” (BALA) (Direcção Regional dos Recursos Florestais, project 17.01-080203) (1999-2003);

“Agriculture, habitat fragmentation, indicator species and conservation of endemic fauna and flora in the Azores – the 2010 Target” (Direcção Regional da Ciência e Tecnologia, DRCT - Postdoc M112/F/014/2007) (2007-2009);

“Predicting extinctions on islands: a multi-scale assessment” (Fundação para a Ciência e Tecnologia- FCT- PTDC/BIA-BEC/100182/2008) (2010-2013);

“Understanding biodiversity dynamics in tropical and subtropical islands as an aid to science based conservation action” (ISLANDBIODIV) (Fundação para a Ciência e Tecnologia , FCT/NETBIOME/0003).

The database management was funded by FEDER (85%) and by Azorean Public funds (15%) through Operational Programme Azores 2020, under the project AZORESBIOPORTAL –PORBIOTA (ACORES-01-0145-FEDER-000072).

Sampling methods

Description: The study was conducted in several habitats of eight islands of the Azores Archipelago (Fig. 1): Flores, São Jorge, Pico, Faial, Terceira, Graciosa, São Miguel and Santa Maria Islands. The sampled habitats include exotic, mixed and native forests, intensive and semi-natural pastures, orchards, caves and lava flows (see details in Event Table at Borges et al. 2021a).

Sampling description: This inventory of Staphylinidae (Coleoptera) includes records obtained in several standardised sampling campaigns and by non-standardised methods performed between 1990 and 2015. Most of the records (96.7%) are based on standardised sampling campaigns, which included pitfall and beating protocols. The used methodology is in accordance with BALA protocol - Biodiversity of Arthropods of the Laurissilva of Azores (Gaspar et al. 2008, Borges et al. 2016) implemented during BALA I (1999-2004) and BALA II (2010-2011) projects. Non-standardised records are based on hand-collecting and sifting, as well as cave, colour and malaise traps.

Standardised sampling campaigns included pitfall traps and beating transects. Pitfall trap transect protocols were conducted with 33 cl plastic cups, partially filled with propylene glycol, in the soil (cup rim at surface level) every 5 m. Traps were protected from rain using a plastic plate, placed about 5 cm above surface level and fixed to the ground with wire. The pitfall traps remained active in the field for 14 days. Beating transects were performed by beating the canopy of woody vegetation, using a beating tray. The protocol was conducted when the vegetation was dry. A 5 m wide square was established every 15 m (total of 10 squares per transect). Two woody plant specimens of the most abundant species (up to three species when available) were sampled in each square. For each selected plant, a branch was chosen at random and a beating tray placed beneath it. The tray consisted of a 1 m wide and 60 cm deep cloth inverted pyramid, with a plastic bag at the vertex. Five beatings were made using a stick for each plant individual sampled.

Quality control: All sorted specimens were identified by a taxonomical expert. Taxonomic nomenclature followed: Schülke and Smetana (2015), Brunke et al. (2021), Yoo et al. (2021).

In terms of species colonisation status, we followed two classifications:

1) For the GBIF database that incorporates occurrence data, we followed the information in the last checklist of Azorean Staphylinidae (Assing 2010). This information is analysed through the main text of this manuscript.

2) However, species classified as "native" by Volker Assing (Assing 2010) have a dubious colonisation origin and were probably inadvertently introduced from the Western

Palearctic. It is highly probable that this happened following human colonisation of the islands and prior to the first reports on Azorean beetles in the 19th century. Exceptions are *Atheta pasadenae* and *Hypomedon debilicornis*, for which the native distribution is unknown.

For that reason, in the current checklist of Azorean Staphylinidae (Suppl. material 1), we now add a column in which we create a new possible tentative categorisation for the colonisation status of the species as follows:

- endemic: species for which we have some evidence that they are true endemics, occurring mostly in native habitats.
- Doubtfully endemic: species whose status is doubtful, based on our current knowledge on the distribution of congeneric species.
- Non-endemic: these include all the previous named as “native” and “introduced” since, in most cases, we currently have no confidence on their status.
- Non-endemic cosmopolitan: these include species with cosmopolitan distribution.

Geographic coverage

Description: Azores, Portugal (Flores, São Jorge, Pico, Faial, Terceira, Graciosa, São Miguel and Santa Maria).

Coordinates: 36.862 and 39.623 Latitude; -31.399 and -24.895 Longitude.

Taxonomic coverage

Description: This publication covers the Staphylinidae family (Insecta, Coleoptera).

Taxa included:

Rank	Scientific Name	Common Name
family	Staphylinidae	Rove-beetles

Traits coverage

Not available

Temporal coverage

Data range: 1990-6-01 - 2015-9-30.

Collection data

Collection name: Entomoteca Dalberto Teixeira Pombo

Collection identifier: DTP

Specimen preservation method: Ethanol 96% for posterior DNA analyses

Curatorial unit: Curator: Paulo A. V. Borges

Usage licence

Usage licence: Creative Commons Public Domain Waiver (CC-Zero)

Data resources

Data package title: Inventory of the Azorean rove-beetles (Insecta, Coleoptera, Staphylinidae)

Resource link: <https://www.gbif.org/dataset/81df7e48-1f76-4125-901b-140bd96dfa49>

Alternative identifiers: http://ipt.gbif.pt/ipt/resource?r=staphylinidae_occurrences_azores

Number of data sets: 2

Data set name: Event Table

Character set: UTF-8

Download URL: http://ipt.gbif.pt/ipt/resource?r=staphylinidae_occurrences_azores

Data format: Darwin Core Archive format

Data format version: Version 1.5

Description: The dataset was published in Global Biodiversity Information Facility platform, GBIF (Borges et al. 2021a). The following data table includes all the records for which a taxonomic identification of the species was possible. The dataset submitted to GBIF is structured as a sample event dataset that has been published as a Darwin Core Archive (DwCA), which is a standardised format for sharing biodiversity data as a set of one or more data tables. The core data file contains 2731 records (eventID). This IPT (Integrated Publishing Toolkit) archives the data and thus serves as the data repository. The data and resource metadata are available for download in the Portuguese GBIF Portal IPT (Borges et al. 2021a).

Column label	Column description
--------------	--------------------

eventID	Identifier of the events, unique for the dataset.
stateProvince	Name of the region of the sampling site.
islandGroup	Name of archipelago.
island	Name of the island.
country	Country of the sampling site.
countryCode	ISO code of the country of the sampling site.
municipality	Municipality of the sampling site.
decimalLongitude	Approximate centre point decimal longitude of the field site in GPS coordinates.
decimalLatitude	Approximate centre point decimal latitude of the field site in GPS coordinates.
geodeticDatum	The ellipsoid, geodetic datum or spatial reference system (SRS) upon which the geographic coordinates given in decimalLatitude and decimalLongitude are based.
coordinateUncertaintyInMetres	Uncertainty of the coordinates of the centre of the sampling plot in metres.
coordinatePrecision	A decimal representation of the precision of the coordinates given in the decimalLatitude and decimalLongitude.
georeferenceSources	A list (concatenated and separated) of maps, gazetteers or other resources used to georeference the Location, described specifically enough to allow anyone in the future to use the same resources.
locationID	Identifier of the location.
fieldNumber	Code of the sample.
locationRemarks	Comments or notes about the Location.
locality	Name of the locality.
habitat	The habitat of the sample.
year	Year of the event.
month	Month of the event.
eventDate	Date or date range the record was collected.
samplingProtocol	The sampling protocol used to capture the species.

Data set name: Occurrence Table

Download URL: http://ipt.gbif.pt/ipt/resource?r=staphylinidae_occurrences_azores

Data format: Darwin Core Archive

Data format version: version 1.5

Description: The dataset was published in Global Biodiversity Information Facility platform, GBIF (Borges et al. 2021a). The following data table includes all the records

for which a taxonomic identification of the species was possible. The dataset submitted to GBIF is structured as a occurrence table that has been published as a Darwin Core Archive (DwCA), which is a standardised format for sharing biodiversity data as a set of one or more data tables. The core data file contains 3903 records (occurrenceID). This IPT (Integrated Publishing Toolkit) archives the data and thus serves as the data repository. The data and resource metadata are available for download in the Portuguese GBIF Portal IPT (Borges et al. 2021a).

Column label	Column description
eventID	Identifier of the events, unique for the dataset.
type	Type of the record, as defined by the Public Core standard.
licence	Reference to the licence under which the record is published.
institutionID	The identity of the institution publishing the data.
institutionCode	The code of the institution publishing the data.
collectionID	The identity of the collection publishing the data.
collectionCode	The code of the collection where the specimens are conserved.
datasetName	Name of the dataset.
basisOfRecord	The nature of the data record.
occurrenceID	Identifier of the record, coded as a global unique identifier.
recordedBy	A list (concatenated and separated) of names of people, groups or organisations who performed the sampling in the field.
identifiedBy	A list (concatenated and separated) of names of people, groups or organisations who assigned the Taxon to the subject.
dateIdentified	The date on which the subject was determined as representing the Taxon.
organismQuantity	A number or enumeration value for the quantity of organisms.
organismQuantityType	The type of quantification system used for the quantity of organisms.
lifeStage	The life stage of the organisms captured.
scientificName	Complete scientific name including author and year.
scientificNameAuthorship	Name of the author of the lowest taxon rank included in the record.
kingdom	Kingdom name.
phylum	Phylum name.
class	Class name.
order	Order name.
family	Family name.
genus	Genus name.

specificEpithet	Specific epithet.
infraspecificEpithet	Infrapecific epithet.
taxonRank	Lowest taxonomic rank of the record.
establishmentMeans	The process of establishment of the species in the location, using a controlled vocabulary: in the GBIF database, we used the Borges et al. (2010) original data: 'native', 'introduced', 'endemic'.
identificationRemarks	Information about morphospecies identification (code in Dalberto Teixeira Pombo Collection).

Additional information

We collected a total of 10,744 specimens belonging to 69 species of Staphylinidae, which represent 51% of the Staphylinidae species currently known from the Azores Archipelago (see Table 1, Suppl. material 1). Of the collected species, a total of 40 species is considered introduced (n = 5392 specimens), 25 native non-endemic (n = 5109 specimens) and four endemic (n = 188 specimens). A total of 58 individuals were identified only at genus level representing nine morphospecies and can be seen at the GBIF IPT (Borges et al. 2021a).

At archipelago level, the most abundant Staphylinidae species were the introduced *Anotylus nitidifrons* (n = 1247 specimens) and *Atheta fungi* (n = 1139) and the native non-endemic *Ocypus aethiops* (n = 1219). At island level, the introduced *Anotylus nitidifrons* was also the most abundant species in Flores (n = 399) and Faial (n = 171) Islands. The introduced *Atheta fungi* was the most abundant species on Graciosa (n = 303) and Santa Maria Islands (n = 649). The native *Ocypus olens* was the most abundant species on Pico Island (n = 71). The native *Ocypus aethiops* was the most abundant species on São Jorge and Terceira Islands; and the native *Quedius curtipennis* was the most abundant species recorded on São Miguel Island (n = 327). The most abundant endemic Staphylinidae was *Notothecta dryochares* (n = 119), collected in six islands (Table 1).

From this dataset, seven species are new for the Azores (Table 1; Suppl. material 1): *Aleochara funebris* Wollaston, 1864; *Amischa forcipata* Mulsant & Rey, 1873; *Bledius unicornis* (Germar, 1825); *Carpelimus troglodytes troglodytes* (Erichson, 1840); *Cypha seminulum* (Erichson, 1839); *Paraphloeostiba gayndahensis* (MacLeay, 1871) and *Tachyporus caucasicus* Kolenati, 1846. We also registered a total of 66 new island records (Table 1) distributed as follows across the individual islands: four for Flores, nine for Faial, six for Pico, 14 for Graciosa, nine for São Jorge, 11 for Terceira, one for São Miguel and 11 for Santa Maria.

Updated Checklist of Azorean Staphylinidae

The latest list of Azorean rove-beetles from the Azores (Assing 2010) included a total of 115 species and subspecies of Staphylinidae. Since then, a new endemic species was

recorded from Azores, *Medon varamontis* Assing, 2013, known only from São Miguel (see Assing 2013) and the previous family Scydmaenidae (including three Azorean species) was included in the Staphylinidae as a subfamily. In addition, seven species were added in the current study (see above and Table 1). Other additions are explained below.

The catalogue of Palearctic Staphylinidae (Schülke and Smetana 2015) lists several species as recorded from the Azores that are not in the latest list of Azores species (Assing 2010). Only four of these species are now added to our updated list (see Suppl. material 1): *Aleochara verna* Say, 1833, *Atheta aeneicollis* (Sharp, 1869), *Sepedophilus littoreus* (Linnaeus, 1758) and *Sepedophilus testaceus* (Fabricius, 1793). The decision to add these species is based on the following reasoning:

- *Aleochara verna*: this species was mentioned previously to Azores by Maus (1998), but without recording the island. Now we record the species for Terceira and Santa Maria (see Table 1).

- *Atheta aeneicollis*: we found no literature record. The species was added in the 2nd edition of the Staphylinidae catalogue by the Editors (Schülke and Smetana 2015), but the record is not traceable backwards. We have now found many specimens in several islands (see Table 1).

- *Sepedophilus littoreus*: Schatzmayr (1945) Ponta Delgada (São Miguel), cited by Horion (1967), Hammond (1973) and Herman (2001), also listed in Borges (1990); the presence of this species in the Azores would be plausible.

- *Sepedophilus testaceus*: old records by Fauvel (1902) and Bernhauer (1940), citing the Crotch (1867) record of *S. sericeus*, and Méquignon (1942). All these records are doubtful and may possibly be based on misidentification, most probably *S. lusitanicus*. However, Hammond (1973), who revised the British species and described *S. lusitanicus*, explicitly wrote that he had seen *S. testaceus* from the Azores. The species was also listed by Borges (1990).

For 14 species listed in the Palearctic Catalogue as present in the Azores (Schülke and Smetana 2015), we found no clear support for their addition to the current list of Azorean rove-beetles:

- *Atheta maderensis*: recorded by Brinck (1977). The record is based on confusion with *Atheta pasadenae* (see Gusarov 2016).

- *Atheta zosterae*: recorded by Bernhauer (1940), Méquignon (1942) and Israelson (1984) (misidentification). Later identified as *Atheta nigra* (Kraatz).

- *Astenus gracilis*: record most likely based on Crotch 1867 and Méquignon (1942); almost certainly misidentification of *A. lyonesius*.

- *Cafius sericeus*: recorded by Israelson (1984) as *Remus sericeus* from Santa Maria. Israelson (1990) stated that this record was based on a misidentification of *Remus pruinus*.

- *Euplectus afer* was recorded by Méquignon (1942) from São Miguel; this record is based on misidentification of *E. infirmus*.

- *Gyrohypnus punctulatus*: old records of Crotch (1867) and Méquignon (1942); almost certainly belong to *G. fracticornis*.

- *Leptacinus batychrus*: recorded by Fauvel (1897), Fauvel (1902) and Bernhauer (1940) from São Miguel. This is a misidentification of *Leptacinus pusillus* (Stephens, 1833).

- *Philonthus rufipes*: was recorded as *P. immundus* by Bernhauer (1940) from São Miguel and subsequently listed by Borges (1990). *P. immundus* is a synonym of *P. ventralis* (Gravenhorst, 1802).

- *Sepedophilus marshami*: this record is based on Crotch (1867), who recorded *Conosomus sericeus* Latreille from Flores. This species was later synonymised with *S. marshami* (Hammond 1973). The record is doubtful and may refer to *S. testaceus* or *S. lusitanicus*; most likely it refers to *S. lusitanicus*.

- For five additional species, no primary records from the Azores were found. Consequently, the respective record from the Azores in the Palearctic Catalogue (Schülke and Smetana 2015) should be deleted: *Deleaster dichrous* Gravenhorst, 1802, *Placusa tachyporoides* (Waltl, 1838), *Scydmaenus rufus* Müller, P.W.J. & Kunze, 1822, *Tomoglossa laeta* Eppelsheim, 1884 and *Tomoglossa luteicornis* (Erichson, 1837).

Based on Suppl. material 1, in addition to the old records, several sources of data have contributed to the new records to Azores and individual islands (Fig. 2). Corvo and São Jorge Islands are possibly still not well studied (Fig. 2).

The current list of Azorean rove-beetles has now 136 species (Suppl. material 1). This list includes ten endemic species, but we are confident that only five are really true endemics, the other five being possibly species from other origins, but not yet recorded in the mainland. The five “true endemics” (*Atheta floresensis* Pace, 2004; *Euconnus azoricus* Franz, 1969; *Medon varamontis* Assing, 2013; *Notothecta dryochares* (Israelson, 1985) and *Phloeostiba azorica* (Fauvel, 1900)) are commonly found in native forests of Azores and, in particular, *N. dryochares* is very abundant in the canopies of Azorean endemic trees.

Most of the species assigned to native non-endemic and introduced status in Assing (2010) are mostly of Palearctic origin and we decided to create an alternative colonisation status categorisation, assigning this species as “non-endemic” and “non-endemic cosmopolitan”. At least 16 species have a worldwide distribution and were classified as “non-endemic cosmopolitan”.

The study of Azorean Staphylinidae is far from complete. We are conducting additional surveys in the Azores (e.g. Costa and Borges 2021, Tsafack et al. 2021) and new records will be soon available covering several habitats.

Acknowledgements

We are grateful to all researchers who collaborated in the field and lab work (see field team list above). The Forest Services and Natural Parks provided logistic support on each island.

Species collecting was possible, based on the funding from the following projects: “Reservas Florestais dos Açores: Cartografia e Inventariação dos Artrópodes Endémicos dos Açores” (BALA) (Direcção Regional dos Recursos Florestais, project 17.01-080203) (1999-2003); “Agriculture, habitat fragmentation, indicator species and conservation of endemic fauna and flora in the Azores – the 2010 Target” (Direcção Regional da Ciência e Tecnologia, DRCT - Postdoc M112/F/014/2007) (2007-2009); “Predicting extinctions on islands: a multi-scale assessment” (Fundação para a Ciência e Tecnologia- FCT- PTDC/ BIA-BEC/100182/2008) (2010-2013); “Understanding biodiversity dynamics in tropical and subtropical islands as an aid to science based conservation action” (ISLANDBIODIV) (Fundação para a Ciência e Tecnologia , FCT/NETBIOME/0003).

The database management was funded by FEDER (85%) and by Azorean Public funds (15%) through Operational Programme Azores 2020, under the project AZORESBIOPORTAL –PORBIOTA (ACORES-01-0145-FEDER-000072).

Author contributions

PAVB led the project and performed research; LLL and PAVB prepared the databases; VA and MS identified species and led taxonomic work. PAVB led the writing with substantial input from the other authors.

References

- Assing V, Schülke M (2006) Systematic catalogue of the entomofauna of the Madeira Archipelago and Selvagens Islands - Staphylinoidea, Staphylinidae (Coleoptera). Vol II, Part 2. Boletim do Museu Municipal do Funchal 1 (Supl.): 1-167. URL: https://islandlab.uac.pt/fotos/publicacoes/publicacoes_Assing06_SystematicCatalogueOfTheEntomofaunaOfTheMadeiraArchipelagoAndSelvagensIslands.pdf
- Assing V (2010) Coleoptera, Staphylinidae. In: Borges PAV, Costa A, Cunha R, Gabriel R, et al. (Eds) A list of the terrestrial and marine biota from the Azores. Princípio, Cascais, 230-231 pp. URL: https://islandlab.uac.pt/fotos/publicacoes/publicacoes_Listagem_ml.pdf [ISBN 978-989-8131-75-1].
- Assing V (2013) A revision of Palaearctic Medon IX. New species, new synonymies, a new combination, and additional records (Coleoptera: Staphylinidae: Paederinae). Entomologische Blätter und Coleoptera 109: 233-270. URL: <http://bionames.org/references/e1ffe485e2261a32049a4cca22c47c32>

- Barratt BIP, Derraiak JGB, Rufaut CG, Goodman AJ, Dickinson KJM (2003) Morphospecies as a substitute for Coleoptera species identification, and the value of experience in improving accuracy. *Journal of the Royal Society of New Zealand* 33 (2): 583-590. <https://doi.org/10.1080/03014223.2003.9517746>
- Bernhauer M (1936) Neuheiten der Palaearktischen Staphyliniden-fauna I, II. [vide pp. 315-316: *Atheta (Hummerliella) azorica* n. sp.]. Pubblicazioni del Museo Entomologico Pietro Rossi 1: 315-316.
- Bernhauer M (1940) Staphyliniden von den Azoren und Madeira. *Societas Scientiarum Fennica. Commentationes Biologicae* 8 (2): 1-10.
- Borges PAV, Rigal F, Ros-Prieto A, Cardoso P (2020) Increase of insular exotic arthropod diversity is a fundamental dimension of the current biodiversity crisis. *Insect Conservation and Diversity* 13 (5): 508-518. <https://doi.org/10.1111/icad.12431>
- Borges PAV, Serrano ARM (1989) New records of the coleopterous fauna (Insecta, Coleoptera) from the Azores. *Boletim do Museu Municipal do Funchal* 41 (209): 5-24. URL: <https://repositorio.uac.pt/handle/10400.3/1887>
- Borges PAV (1990) A checklist of the Coleoptera from the Azores with some systematic and biogeographic comments. *Boletim do Museu Municipal do Funchal* 42 (220): 87-136. URL: <https://repositorio.uac.pt/handle/10400.3/1904>
- Borges PAV, Gaspar C, Crespo LC, Rigal F, Cardoso P, Pereira F, Rego C, Amorim IR, Melo C, Aguiar C, André G, Mendonça E, Ribeiro S, Hortal J, Santos AMC, Barcelos L, Enghoff H, Mahnert V, Pita M, Ribes J, Baz A, Sousa AB, Vieira V, Wunderlich J, Parmakelis A, Whittaker RJ, Quartau JA, Serrano ARM, Triantis KA (2016) New records and detailed distribution and abundance of selected arthropod species collected between 1999 and 2011 in Azorean native forests. *Biodiversity Data Journal* 4 <https://doi.org/10.3897/bdj.4.e10948>
- Borges PAV, Amorim IR, Terzopoulou S, Rigal F, Emerson BC, Serrano ARM (2017) Cryptic diversity in the Azorean beetle genus *Tarphius* Erichson, 1845 (Coleoptera: Zopheridae): An integrative taxonomic approach with description of four new species. *Zootaxa* 4236 (3): 401-449. <https://doi.org/10.11646/zootaxa.4236.3.1>
- Borges PAV, Gabriel R, Pimentel C, Brito M, Serrano ARM, Crespo LC, Assing V, Stüben P, Fattorini S, Soares AO, Mendonça E, Nogueira E (2018) Biota from the coastal wetlands of Praia da Vitória (Terceira, Azores, Portugal): Part 1 - Arthropods. *Biodiversity Data Journal* 6 (e27194). <https://doi.org/10.3897/bdj.6.e27194>
- Borges PAV, Santos AMC, Elias RB, Gabriel R (2020) The Azores archipelago: biodiversity erosion and conservation biogeography. *Encyclopedia of the World's Biomes -Earth Systems and Environmental Sciences. Reference Module in Earth Systems and Environmental Sciences*. Elsevier, Amsterdam, Netherlands, 18 pp. <https://doi.org/10.1016/B978-0-12-409548-9.11949-9>
- Borges PAV, Lamelas-López L, Assing V, Schülke M (2021a) Inventory of the Azorean rove-beetles (Insecta, Coleoptera, Staphylinidae). v1.5. Universidade dos Açores. <https://doi.org/10.15468/fh6qar>
- Borges PAV, Nunes R, Lamelas-López L, Pereira E, Costa R, Monjardino P, Lopes DH, Soares AO, Gil A, Rigal F, Lövei GL, Ferrante M (2021b) Monitoring Arthropods in Azorean Agroecosystems: the project AGRO-ECOSERVICES. *Biodiversity Data Journal* 9 (e77548). <https://doi.org/10.3897/BDJ.9.e77548>

- Brinck P (1977) Staphylinidae (Coleoptera): Atheta from the Azores and Madeira. Boletim do Museu Municipal do Funchal 31 (136): 84-86. URL: <https://publications.cm-funchal.pt/jspui/handle/100/1219>
- Brunke AJ, Pentinsaari M, Klimaszewski J (2021) Integrative taxonomy of Nearctic and Palaearctic Aleocharinae: new species, synonymies, and records (Coleoptera, Staphylinidae). ZooKeys 1041: 27-29. <https://doi.org/10.3897/zookeys.1041.64460>
- Cardoso P, Aranda SC, Lobo JM, Dinis F, Gaspar C, Borges PAV (2009) A spatial scale assessment of habitat effects on arthropod communities of an oceanic island. Acta Oecologica 35 (5): 590-597. <https://doi.org/10.1016/j.actao.2009.05.005>
- Costa R, Borges PAV (2021) SLAM Project - Long term ecological study of the impacts of climate change in the natural forest of Azores: I - the spiders from native forests of Terceira and Pico Islands (2012-2019). Biodiversity Data Journal 9 (e69924). <https://doi.org/10.3897/BDJ.9.e69924>
- Crotch GR (1867) On the Coleoptera of the Azores. Proceedings of the Zoological Society of London 359-391. URL: https://islandlab.uac.pt/fotos/publicacoes/publicacoes_CROTC67_OntheColeopteraoftheAzores_A00127.pdf
- Fauvel A (1897) Catalogue des Staphylinides de la Barbarie et des îles des Açores, Madère, Salvages et Canaries. Revue de Entomologie 16: 237-369.
- Fauvel A (1898) Catalogue des Staphylinides de la Barbarie et des îles des Açores, Madère, Salvages et Canaries. Notices Entomologiques 10: 237-371.
- Fauvel A (1900) Staphylinides Paléarctiques nouveaux. Revue de Entomologie 19: 219-220.
- Fauvel A (1902) Catalogue des Staphylinides de la Barbarie et des îles des Açores, Madère, Salvages et Canaries. Revue de Entomologie 21: 45-189.
- Gaspar C, Borges PAV, Gaston KJ (2008) Diversity and distribution of arthropods in native forests of the Azores archipelago. Arquipélago Life and Marine Sciences 25: 1-30. URL: http://www.oceanos.uac.pt/storage/2008/10/pp_1_30_Gaspar_et al_25.pdf
- Gillerfors G (1986) Contribution to the coleopterous fauna of the Azores. Boletim do Museu Municipal do Funchal 38 (172): 16-27. URL: <https://publications.cm-funchal.pt/jspui/bitstream/100/1169/1/Bolmmf-1986-art172.pdf>
- Gillerfors G (1988) Second contribution to the coleopterous fauna of the Azores. Bocagiana 118: 1-6. URL: <https://publications.cm-funchal.pt/jspui/bitstream/100/1433/1/Boc118-1988.pdf>
- Hammond PM (1973) Notes on British Staphylinidae 3. The British species of *Sepedophilus* Gistel (*Conosomus* Auctt.). Entomologist's Monthly Magazine 108: 130-165.
- Herman LH (2001) Catalog of the Staphylinidae (Insecta: Coleoptera). 1758 to the end of the second millennium. II. Tachyporine group. Bulletin of the American Museum of Natural History 265: 651-1061. <https://doi.org/10.1206/0003-0090.265.1.2>
- Horion AD (1967) Faunistik der Mitteleuropäischen Käfer. Band XI: Staphylinidae. 3 Teil: Habrocerinae bis Aleocharinae (ohne Subtribus Athetae). [Faunistics of the Central European Beetles. Volume XI: Staphylinidae. 3 Part: Habrocerinae to Aleocharinae (without subtribe Athetae)]. 11. Kommisionsverlag Buchdruckerei Aug. Feyel, Überlingen-Bodensee, 419 pp. [In German].
- Israelson G (1984) Coleoptera from the Azores. Boletim do Museu Municipal do Funchal 36 (161): 142-161. URL: <https://publications.cm-funchal.pt/jspui/bitstream/100/1193/1/Bolmmf-1984-art161.pdf>

- Israelson G (1985) Notes on the coleopterous fauna of the Azores, with description of new species of *Atheta* Thomson (Coleoptera). Boletim do Museu Municipal do Funchal 37 (165): 5-19. URL: <https://publications.cm-funchal.pt/jspui/bitstream/100/1181/1/Bolmmf-1985-art165.pdf>
- Israelson G (1990) Further notes on the coleopterous fauna of the Azores, with speculations on its origin. Bocagina 138: 1-8. URL: <https://publications.cm-funchal.pt/jspui/bitstream/100/1419/1/Boc138-1990.pdf>
- Jarrige J (1953) Les Staphylinus des îles Atlantides (Col. Staphylinidae). Bulletin de la Société Entomologique de France 58: 160-164. <https://doi.org/10.3406/bsef.1953.18676>
- Malumbres-Olarte J, Cardoso P, Crespo LC, Gabriel R, Pereira F, Carvalho R, Rego C, Nunes R, Ferreira M, Amorim IR, Rigal F, Borges PAV (2019) Standardised inventories of spiders (Arachnida, Araneae) of Macaronesia I: The native forests of the Azores (Pico and Terceira islands). Biodiversity Data Journal 7 (e32625). <https://doi.org/10.3897/bdj.7.e32625>
- Marcelino J, Borges PAV, Borges I, Pereira E, Santos V, Soares AO (2021) Standardised arthropod (Arthropoda) inventory across natural and anthropogenic impacted habitats in the Azores archipelago. Biodiversity Data Journal 9 (e62157). <https://doi.org/10.3897/bdj.9.e62157>
- Matthews TJ, Sadler J, Carvalho R, Nunes R, Borges PAV (2018) Differential temporal beta-diversity patterns of native and non-native arthropod species in a fragmented native forest landscape. Ecography 42 (1): 45-54. <https://doi.org/10.1111/ecog.03812>
- Maus CH (1998) Taxonomical contributions to the subgenus *Coprochara* Mulsant & Rey, 1874 of the genus *Aleochara* Gravenhorst, 1802. Koleopterologische Rundschau 68: 81-100. URL: https://www.zobodat.at/pdf/KOR_68_1998_0081-0100.pdf
- Méquignon A (1942) Voyage de MM. L. Chopard et A. Méquignon aux Açores (Août-Septembre 1930). XIV. Catalogue des Coléoptères Açoréens. Annales de la Société Entomologique de France 111: 1-66.
- Oromí P, Serrano ARM, Borges PAV (2010) Coleoptera (Coordination). In: Borges PAV, Costa R, Cunha R, Gabriel R, et al. (Eds) A list of the terrestrial and marine biota from the Azores. Principia, Cascais, 222-232 pp. URL: https://islandlab.uac.pt/fotos/publicacoes/publicacoes_Listagem_ml.pdf [ISBN 978-989-8131-75-1].
- Pace R (2004) Hygronomini e Athetini delle Cina con note sinonimiche (Coleoptera, Staphylinidae). Revue Suisse de Zoologie 111 (3): 457-523. <https://doi.org/10.5962/bhl.part.80248>
- Schatzmayr A (1945) Appunti Coleopterologici: XVI. Natura, Milano 36: 44-48.
- Schülke M, Smetana A (2015) Staphylinidae. In: Löbl I, Löbl D (Eds) Catalogue of Palaearctic Coleoptera. New, updated Edition. 2. Brill, Leiden, 304–1134 pp.
- Serrano ARM (1982) Contribuição para o conhecimento do povoamento, distribuição e origem dos Coleópteros do arquipélago dos Açores (Insecta, Coleoptera). Boletim do Museu Municipal do Funchal 34 (147): 67-104. URL: <https://publications.cm-funchal.pt/jspui/bitstream/100/1208/1/Bolmmf-1982-art147.pdf>
- Serrano ARM, Borges PAV (1987) A further contribution to the knowledge of the Coleoptera (Insecta) from the Azores. Boletim do Museu Municipal do Funchal 39 (187): 51-69. URL: <https://publications.cm-funchal.pt/jspui/bitstream/100/1159/1/Bolmmf-1987-art187.pdf>
- Shaw JJ, Solodovnikov A (2016) Systematic and biogeographic review of the Staphylinini rove beetles of Lord Howe Island with description of new species and

taxonomic changes (Coleoptera, Staphylinidae). ZooKeys 638: 1-25. <https://doi.org/10.3897/zookeys.638.10883>

- Smetana A (1970) Staphylinidae (Coleoptera) from the Azores and Madeira. Boletim do Museu Municipal do Funchal 25 (110): 53-67. URL: <https://publications.cm-funchal.pt/jspui/bitstream/100/1219/1/Bolmmf-1977-art136.pdf>
- Tsafack N, Fattorini S, Boieiro M, Rigal F, Ros-Prieto A, Ferreira MT, Borges PAV (2021) The Role of Small Lowland Patches of Exotic Forests as Refuges of Rare Endemic Azorean Arthropods. Diversity 13 (9): 443. <https://doi.org/10.3390/d13090443>
- Yoo I, Song J, Ahn K (2021) Total-evidence phylogeny of the coastal *Cafius* complex (Coleoptera: Staphylinidae) and its new generic concepts. Zoologica Scripta 50 (6): 734-751. <https://doi.org/10.1111/zsc.12506>




Figure 1.

The Azores Archipelago with its location in the middle Atlantic (left panel) and the nine Azorean islands (right panel): the western group (COR - Corvo and FLO - Flores), the central group (FAI - Faial, PIC - Pico, GRA - Graciosa, SJO - São Jorge and TER - Terceira) and the eastern group (SMG - São Miguel and SMA - Santa Maria).




Figure 2.

The number of rove-beetle species per island with an indication of species listed by 2010 (Old records; see Assing 2010), the species added in literature after 2010 and the new records based in the current work. COR - Corvo; FLO - Flores; FAI - Faial; PIC - Pico; SJG - São Jorge; GRA- Graciosa; TER - Terceira; SMG - São Miguel; SMR - Santa Maria.

Table 1.

Inventory of the Staphylinidae species collected in eight islands of Azores, from 1990 to 2015. The list includes individuals identified at species-level. Scientific name, colonisation status (CS) following the classification in Assing (2010) (int - introduced; nat - native non-endemic; end - endemic) and abundance per island are provided. Bold scientific names constitute new records for Azores and bold numbers new records for a given island. FLO - Flores; FAI - Faial; PIC - Pico; SJG - São Jorge; GRA- Graciosa; TER - Terceira; SMG - São Miguel; SMR - Santa Maria.

Scientific Name	CS	FLO	FAI	PIC	GRA	SJG	TER	SMG	SMR	Total
<i>Aleochara bipustulata</i> (Linnaeus, 1760)	int	5	1	0	3	1	63	0	11	84
<i>Aleochara clavicornis</i> L. Redtenbacher, 1849	int	1	0	0	0	0	0	0	0	1
<i>Aleochara funebris</i> Wollaston, 1864	nat	0	0	0	1	0	0	0	0	1
<i>Aleochara puberula</i> Klug, 1833	int	0	0	11	0	0	0	0	0	11
<i>Aleochara verna</i> Say, 1833	int	0	0	0	0	0	1	0	6	7
<i>Aloconota sulcifrons</i> (Stephens, 1832)	nat	2	23	0	1	0	186	11	3	226
<i>Amischa analis</i> (Gravenhorst, 1802)	int	13	31	1	0	0	244	1	21	311
<i>Amischa forcipata</i> Mulsant & Rey, 1873	int	0	0	0	0	0	1	0	18	19
<i>Anotylus complanatus</i> (Erichson, 1839)	int	0	0	1	0	0	0	0	0	1
<i>Anotylus nitidifrons</i> (Wollaston, 1871)	int	399	171	28	168	1	246	234	0	1247
<i>Anotylus nitidulus</i> (Gravenhorst, 1802)	int	0	1	0	0	0	2	0	5	8
<i>Astenus lyonessius</i> (Joy, 1908)	nat	4	4	0	2	0	9	0	6	25
<i>Atheta aeneicollis</i> (Sharp, 1869)	int	0	2	10	4	0	123	2	24	165
<i>Atheta atramentaria</i> (Gyllenhal, 1810)	int	0	0	0	0	0	10	0	0	10
<i>Atheta fungi</i> (Gravenhorst, 1806)	int	2	24	13	303	5	136	7	649	1139
<i>Atheta nigra</i> (Kraatz, 1856)	int	0	0	0	0	0	1	0	0	1
<i>Atheta palustris</i> (Kiesenwetter, 1844)	int	0	0	0	1	1	1	1	0	4
<i>Atheta pasadenae</i> Bernhauer, 1906	nat	0	0	0	0	0	338	0	0	338
<i>Bledius unicoloris</i> (Germar, 1825)	int	0	0	0	0	0	1	0	0	1
<i>Carpelimus bilineatus</i> (Stephens, 1834)	int	0	0	0	0	0	0	0	130	130
<i>Carpelimus corticinus</i> (Gravenhorst, 1806)	nat	20	43	2	2	2	32	3	3	107
<i>Carpelimus pusillus</i> (Gravenhorst, 1802)	int	0	0	0	0	0	0	0	183	183
<i>Carpelimus troglodytes troglodytes</i> (Erichson, 1840)	int	0	0	0	0	0	0	1	0	1
<i>Carpelimus zealandicus</i> (Sharp, 1900)	int	0	0	0	0	0	0	2	3	5

<i>Coproporus pulchellus</i> (Erichson, 1839)	int	0	0	2	3	0	210	0	3	218
<i>Cordalia obscura</i> (Gravenhorst, 1802)	int	54	41	0	38	0	435	0	51	619
<i>Cypha seminulum</i> (Erichson, 1839)	int	0	0	0	1	0	4	0	0	5
<i>Euplectus infirmus</i> Raffray, 1910	int	0	1	0	1	0	2	0	0	4
<i>Gabrius nigrutilus</i> (Gravenhorst, 1802)	int	0	0	0	4	0	36	1	6	47
<i>Gyrophypnus fracticornis</i> (Müller, 1776)	int	0	0	4	0	0	37	0	0	41
<i>Habrocerus capillaricornis</i> (Gravenhorst, 1806)	nat	0	0	3	0	4	3	0	0	10
<i>Hydrosnecta longula</i> (Heer, 1839)	nat	0	0	0	0	0	0	9	0	9
<i>Hypomedon debilicornis</i> (Wollaston, 1857)	nat	0	0	0	0	0	0	0	1	1
<i>Lithocharis nigriceps</i> Kraatz, 1859	int	0	0	0	0	0	0	0	23	23
<i>Lithocharis ochracea</i> (Gravenhorst, 1802)	int	0	0	0	0	0	0	0	7	7
<i>Medon apicalis</i> (Kraatz, 1857)	nat	0	1	0	0	0	0	0	0	1
<i>Medon varamontis</i> Assing, 2013	end	0	0	0	0	0	0	5	0	5
<i>Nacaeus impressicollis</i> (Motschulsky, 1858)	int	0	0	0	0	0	2	0	0	2
<i>Notothecta caprariensis</i> (Israelson, 1985)	end	0	0	0	0	0	0	1	0	1
<i>Notothecta dryochares</i> (Israelson, 1985)	end	0	0	5	1	1	48	38	26	119
<i>Ocypus aethiops</i> (Waltl, 1835)	nat	0	0	0	15	204	1000	0	0	1219
<i>Ocypus olens</i> (Müller, 1764)	nat	34	50	71	1	202	229	35	5	627
<i>Oligota parva</i> Kraatz, 1862	int	0	0	0	0	0	0	0	5	5
<i>Oligota pumilio</i> Kiesenwetter, 1858	nat	0	7	0	1	0	29	0	5	42
<i>Oxytelus sculptus</i> Gravenhorst, 1806	int	0	0	0	0	0	0	0	26	26
<i>Paraphloeostiba gayndahensis</i> (MacLeay, 1871)	int	0	3	0	0	0	200	0	0	203
<i>Phacophallus parumpunctatus</i> (Gyllenhal, 1827)	int	0	0	0	0	0	0	0	9	9
<i>Philonthus discoideus</i> (Gravenhorst, 1802)	int	0	0	0	0	0	0	0	2	2
<i>Philonthus quisquiliarius quisquiliarius</i> (Gyllenhal, 1810)	int	0	0	0	0	0	0	48	0	48
<i>Philonthus umbratilis</i> (Gravenhorst, 1802)	int	0	0	0	0	0	0	2	0	2
<i>Philonthus ventralis</i> (Gravenhorst, 1802)	nat	0	0	0	0	0	0	0	2	2
<i>Phloeonomus punctipennis</i> Thomson, 1867	nat	0	2	1	2	1	8	0	7	21
<i>Phloeostiba azorica</i> (Fauvel, 1900)	end	0	0	3	17	1	6	36	0	63
<i>Platystethus nitens</i> (Sahlberg, 1832)	nat	2	0	3	0	0	0	0	11	16
<i>Proteinus atomarius</i> Erichson, 1840	nat	0	24	14	0	3	52	9	17	119

<i>Pseudoplectus perplexus</i> (Jacquelin du Val, 1854)	nat	2	0	0	0	0	11	0	0	13
<i>Quedius curtipennis</i> Bernhauer, 1908	nat	0	1	0	0	3	73	327	270	674
<i>Quedius simplicifrons</i> Fairmaire, 1862	nat	0	8	14	0	59	16	3	1	101
<i>Rugilus orbiculatus</i> (Paykull, 1789)	nat	104	25	1	72	2	344	2	217	767
<i>Sepedophilus lusitanicus</i> Hammond, 1973	nat	0	1	19	19	35	16	93	11	194
<i>Stenomastax madeirae</i> Assing, 2003	nat	0	0	0	0	0	579	0	0	579
<i>Stenus guttula guttula</i> Müller, 1821	nat	0	4	0	0	0	2	0	0	6
<i>Sunius propinquus</i> (Brisout de Barneville, 1867)	nat	0	0	0	1	0	1	0	3	5
<i>Tachyporus caucasicus</i> Kolenati, 1846	int	0	0	0	0	0	0	0	1	1
<i>Tachyporus chrysomelinus</i> (Linnaeus, 1758)	int	379	6	0	12	0	4	2	32	435
<i>Tachyporus nitidulus</i> (Fabricius, 1781)	int	8	2	0	4	0	9	1	270	294
<i>Trichiusa robustula</i> Casey, 1893	int	1	0	1	0	3	4	0	7	16
<i>Trichophya pilicornis</i> (Gyllenhal, 1810)	nat	0	0	0	0	0	3	0	0	3
<i>Xantholinus longiventris</i> Heer, 1839	int	0	0	2	5	0	50	0	0	57

Supplementary material

Suppl. material 1: Checklist of Azorean Staphylinidae

Authors: Paulo A. V. Borges, Volker Assing & Michael Schülke

Data type: Occurrences

Brief description: Detailed distribution of Azorean Staphylinidae in the nine Azorean islands (AZ - Azores without reference to a given island; COR - Corvo FLO - Flores; FAI - Faial; PIC - Pico; SJG - São Jorge; GRA- Graciosa; TER - Terceira; SMG - São Miguel; SMR - Santa Maria). New records per island are marked. We add also the known taxonomic or nomenclature changes in Azores in four categories ("synonym", "different combination", "misidentification" and "emendation/ misspelling").

"1" – Confirmed occurrence, based on Assing (2010) list; BALA - records based on BALA protocol (see Borges et al. 2016; samples part of current study); LAND-USE (see Cardoso et al. 2009, samples part of current study); MACDIV (see Malumbres-Olarte et al. 2019); ASSING - Non-standardised samples of one of us (Volker Assing) (part of current study); Marcelino - Marcelino et al. (2021); INTERF (data from project INTERFRUTA part of current study); WET – from Borges et al. (2018); AGRO – from Borges et al. (2021b); SLAM - Long term monitoring of Azorean forests using SLAM traps for Terceira Island (see Matthews et al. 2018; Borges et al. 2020; Tsafack et al. 2021); SLAM for other islands (unpublished data); ASSING 2013 (new species described in Assing 2013).

[Download file](#) (14.57 kb)