

# Fish diversity of a spring field in Hopong Town, Taunggyi District, Shan State, Myanmar (the Salween River Basin), with genetic comparisons to some “species endemic to Inle Lake”

Yuichi Kano<sup>‡,§</sup>, Yusuke Fuke<sup>‡</sup>, Prachya Musikasinthorn<sup>¶</sup>, Akihisa Iwata<sup>‡</sup>, Tin Mya Soe<sup>#</sup>, Sein Tun<sup>#,¶</sup>, LKC Yun<sup>#,«</sup>, Seint Seint Win<sup>»^</sup>, Shoko Matsui<sup>‡,^</sup>, Ryoichi Tabata<sup>‡,‡</sup>, Katsutoshi Watanabe<sup>‡</sup>

<sup>‡</sup> Kyushu University, Fukuoka, Japan

<sup>§</sup> Kyushu Open University, Fukuoka, Japan

<sup>‡</sup> Kyoto University, Kyoto, Japan

<sup>¶</sup> Kasetsart University, Bangkok, Thailand

<sup>#</sup> Inlay Lake Wildlife Sanctuary, Nyaung Shwe, Myanmar

<sup>¶</sup> Natma Taung National Park, Kanpetlet, Myanmar

<sup>«</sup> Hkakaborazi National Park, Putao, Myanmar

<sup>»</sup> Taunggyi University, Taunggyi, Myanmar

<sup>^</sup> Kyaing Tong University, Keng Tung, Myanmar

<sup>^</sup> Osaka Museum of Natural History, Osaka, Japan

<sup>‡</sup> Lake Biwa Museum, Kusatsu, Japan

Corresponding author: Katsutoshi Watanabe ([watanak@terra.zool.kyoto-u.ac.jp](mailto:watanak@terra.zool.kyoto-u.ac.jp))

Academic editor: Yahui Zhao

## Abstract

Hopong, a small town in the Salween (Thanlwin) River Basin, Myanmar, is located 35 km northeast of Inle Lake, a famous ancient lake with numerous endemic fish species. We surveyed the fish fauna of a spring pond in Hopong in 2016, 2019 and 2020 and identified 25 species. Of these, seven, including *Inlecypris auropurpureus* and *Sawbwa resplendens*, had been considered endemic to Inle Lake and at least three species were genetically unique. Eight were suspected or definite introduced species, including *Oreochromis niloticus* and *Gambusia affinis*. We were unable to identify a nemacheilid species of the genus *Petruichthys*, which would need a taxonomic examination. The Hopong area is being developed rapidly and, hence, it is crucial to conserve its native fish species and the freshwater ecosystems.

## Keywords

Burma, *Inlecypris auropurpureus*, *Microrasbora rubescens*, mitochondrial DNA (mtDNA), *Physoschistura brunneana*, *Sawbwa resplendens*

## Introduction

Hopong is a small town in the Salween (Thanlwin) River Basin, situated 15 km east of Taunggyi, the capital of Shan State, Myanmar (Fig. 1). The town has several natural wetlands and springs. *Danio* (*Celestichthys*) *margaritatus*, a minute cyprinid fish, is a symbolic fish species from this area (Roberts 2007, Conway et al. 2008). However, the fields of Hopong have been developed rapidly with extensive road construction and urbanisation (Roberts 2007).

Hopong is about 35 km northeast of Inle Lake, a famous ancient lake with many endemic fish species (Annandale 1918, Kano et al. 2016, Win 2018). The Hopong area and Inle Lake are both in the Salween River Basin, so are close together geographically; however, they belong to different drainages separated by highlands and are ca. 400 km apart by river (Fig. 1). The endemism of fishes in Inle Lake has not been validated, because studies of the fish fauna of the middle–upper Salween Basin are insufficient. Thus, the ichthyofauna of the Hopong area is of biogeographical interest, especially in terms of comparison with that of Inle Lake, including the validation of species endemic to Inle Lake.

Generally, studies of freshwater fish biodiversity in Myanmar are insufficient; most are fragmentary reports on fauna of specific regions or new species. Annandale (1918) and Hora and Mukerji (1934) reported the fish fauna of Inle Lake and Shan State, respectively. Musikasinthorn (1998a) first reported *Channa panaw* from the Irrawaddy and Sittang River Basins, Myanmar. Britz (2003) described a new species of miniature fish, *Danionella mirifica*, from northern Myanmar. Roberts (2012) reported a new species of arowana, *Scleropages inscriptus*, from the Malay Peninsula, Myanmar. Recently, Vidthayanon et al. (2005) provided a rough list of inland fish species of Myanmar. Data on the freshwater fish biodiversity of Myanmar are generally limited and more information is required.

Here, we report the results of a freshwater fish inventory for a spring field in Hopong conducted in 2016, 2019 and 2020. The list contains 25 species, including seven “species endemic to Inle Lake”. We provide DNA barcoding (mitochondrial COI sequences) data for the species, which clarify their genetic uniqueness and should promote biogeographical research in this area.

## Methods

Sampling was conducted nine times in Hopong from March 2016 to March 2020 (Fig. 2). Fish were collected with hand-nets, throwing nets and fish traps. Except for several individuals that were only photographed or recorded on-site and released, the collected specimens were photographed fresh (Kano and Nakajima 2014), fixed in 10% formalin and then transferred to 70% ethanol. The specimens and their tissue samples were catalogued and deposited at the Research Laboratory of Ichthyology, Department of Fishery Biology, Faculty of Fisheries, Kasetsart University, Bangkok, Thailand (RLIKU) and Kyoto

University, Kyoto, Japan (tissue samples). All specimens were assigned IDs associated with the records of location (latitude, longitude and region name), collection date, DNA sequence accession numbers etc. The data were registered in the integrated Monsoon Asia “ffish.asia” online database of freshwater organism biodiversity (Watanabe et al. 2009, Kano et al. 2013) and can be retrieved at <https://ffish.asia/Hopong2020>.

To obtain DNA barcoding data (partial mitochondrial COI sequences) for the species collected from Hopong (102 specimens from 19 species including all native species), total DNA was extracted using a Genomic DNA Purification kit (Promega) or Monarch Genomic DNA Purification Kit (New England Biolabs). The COI gene was amplified by PCR using the primer pair FishF1 (5'-TCA ACC AAC CAC AAA GAC ATT GGC AC-3') and FishR1 (5'-TAG ACT TCT GGG TGG CCA AAG AAT CA-3') (Ward et al. 2005). The amplification consisted of an initial denaturation step (94°C, 2 min); 30 cycles of 94°C for 30 s, 56–60°C for 30 s and 72°C for 60 s; and a final extension (72°C, 7 min). The PCR products were purified with Illustra ExoStar (GE Healthcare) and sequenced on an ABI 3130xl Genetic Analyzer (Applied Biosystems) using an amplification primer and the BigDye Terminator Cycle sequencing FS Ready Reaction kit ver. 3.1 (Applied Biosystems). The DNA sequences (640 bp) were deposited at DDBJ/EMBL/GenBank (accession numbers: LC190268–LC190330, LC190383, LC190395–LC190405, LC645163–LC645189).

The obtained sequences were aligned using MAFFT (Katoh and Standley 2013) at Unipro UGENE (Okonechnikov et al. 2012). Sequence data for the same or related species from Inle Lake and the surrounding rivers, as reported by Kano et al. (2016), were included in the analysis for comparison with the Hopong populations. To visualise the relationships amongst populations, haplotype networks, based on the TCS algorithm (statistical parsimony; Clement et al. 2000), were constructed for five genera endemic to this region—*Inlecypis*, *Microrasbora*, *Sawbwa*, *Petruichthys* and *Physoschistura*—using POPART (Leigh and Bryant 2015). The mean uncorrected sequence differences amongst different populations were calculated using MEGA 7 (Kumar et al. 2016).

## Results

The survey uncovered 25 species (Table 1). We collected *Danio margaritatus* in a shallow wetland habitat, as described by Roberts (2007). We also obtained *Devario sondhii*, which was reported by Hora and Mukerji (1934) with a specimen obtained in the Hopong area. We also found a likely undescribed species of *Petruichthys*, which we tentatively labelled “*Petruichthys* sp. (Hopong)”. Of the 25 species, seven had been treated as “species endemic to Inle Lake”: *Danio erythromicron*, *Inlecypis auropurpureus*, *Microrasbora rubescens*, *Sawbwa resplendens*, *Petruichthys brevis*, *Channa harcourtbutleri* and *Mastacembelus caudiocellatus* (Annandale 1918, Kano et al. 2016). Eight were suspected or definite introduced species: *Cyprinus rubrofasciatus*, *Esomus danrica*, *Parambassis lala*, *Oreochromis niloticus*, *Trichogaster labiosa*, *Oryzias uwai*, *Gambusia affinis* and *Poecilia reticulata* (Annandale 1918, Hora and Mukerji 1934, Musikasinthorn 1998b, Kano et al. 2016). Three of these (i.e. *E. danrica*, *T. labiosa*, *O. uwai*) are distributed widely in

Southeast Asia; hence the possibility that these species are native cannot be completely ruled out.

Haplotype networks of the populations of three species from the Hopong area (*Inlecypis auropurpureus* [Fig. 3A], *Microrasbora rubescens* [Fig. 3B] and *Physoschistura rivulicola* [Fig. 3C]) showed genetic uniqueness compared to Inle Lake populations, even within species. *Inlecypis auropurpureus*, *Microrasbora rubescens* and *Physoschistura rivulicola* showed clear genetic divergence (4.1%, 3.2% and 2.1%, respectively, in mean uncorrected sequence differences). By contrast, *Sawbwa resplendens* (Fig. 3D) and *Petruichthys brevis* (Fig. 3E) showed no clear genetic divergence amongst local populations (0.3% and < 0.1%, respectively). *Petruichthys* sp. (Hopong) showed an obviously distinct genetic profile from the sympatric *P. brevis* (12.6% mean uncorrected sequence difference; Fig. 3E).

## Discussion

A total of eleven freshwater fish species have been reported from Hopong in previous papers (Hora and Mukerji 1934, Roberts 2007, Kullander et al. 2017). All of them, except for *Barbus hexastichus* (valid as *Neolissochilus hexastichus*), are judged to be included in the 25 species reported in this study. Hora and Mukerji (1934) reported *Barbus hexastichus* from Hopong as having two pairs of barbels, a light brown body colour and a black round spot at the base of the caudal fin. However, this record should be reconsidered since its main distribution range is in India (Talwar and Jhingran 1991). In fact, *Neolissochilus nigrovittatus*, which is distributed around Inle Lake (Talwar and Jhingran 1991, Kano et al. 2016), also fits the description of the fish by Hora and Mukerji (1934). Further survey and examination are necessary for the *Neolissochilus* species in this area.

Our results indicated that at least seven “species endemic to Inle Lake” (Annandale 1918, Kano et al. 2016, Win 2018) have wider distributions than previously thought, beyond the Inle Lake Basin (Table 1). Artificial introduction from Inle Lake cannot explain their wide distributions because of the genetic divergence of the Hopong populations from the Inle Lake populations, except for *Sawbwa resplendens* (only one DNA sample and details unknown) and *Petruichthys brevis* (Fig. 3D and E). Therefore, Hopong shares fish fauna with the Inle Lake Basin, but the populations of several fish species in Hopong are genetically unique and require detailed taxonomic examinations.

*Danio margaritatus* is symbolic of Hopong (Roberts 2007). Conway et al. (2008) reported that *D. margaritatus* is distributed not only in Hopong (the type locality of the species; Roberts 2007), but also in wider areas of Myanmar and Thailand following Clarke (2007) and Hary (2007). However, these sources are not formal scientific reports (website and local report, respectively) and, in fact, we could not access them. Thus, at present, it cannot be determined whether *D. margaritatus* is distributed only in the Hopong area or more widely. If the former is the case, conservation of *D. margaritatus* and its habitat, i.e. shallow ponds/wetlands with plants and clear water from springs (Fig. 2B; Roberts 2007), is very important. *Devario sondhii* might also be endemic to the Hopong area; more information on its distribution is necessary to determine its conservation status.

*Petruichthys* sp. (Hopong) is genetically distinguishable from the sympatric *P. brevis* (Fig. 3 E). Morphologically, its body is smaller and the male is redder than that of *P. brevis*. Taxonomic examination of this fish is required.

The Hopong area is being developed rapidly (Roberts 2007). In addition, several invasive alien fish have become established, including *Oreochromis niloticus* and *Gambusia affinis*. This strongly suggests that the native freshwater fishes and other native freshwater organisms in Hopong are now severely threatened. As explained above, Hopong has a unique fish fauna that includes endemic species and species shared with Inle Lake, an unique ancient lake in Southeast Asia. Therefore, wetland and biodiversity conservation in Hopong are a high priority.

## Acknowledgements

We deeply appreciate the kind assistance from Nyi Nyi Kyaw (Forest Department, Ministry of Natural Resources and Environmental Conservation, Myanmar), Win Naing Thaw and Naing Zaw Htun (Nature and Wildlife Conservation Division), the late Than Htay (Inle Lake Wildlife Sanctuary), late Ohn, Than Nwai and Sein Thet (Forest Resource Environment Development and Conservation Association, Myanmar; FREDa) and Nyi Nyi Lwin (Green Leaf, Myanmar). This work was supported by the Sumitomo Foundation (the Grant for Environmental Research Project, 193271), JSPS KAKENHI (Grant Number JP26304007, JP19J23130, JP20HP8020) and the research grant of Kurita Water and Environment Foundation (18K013, 19K008).

## Conflicts of interest

## References

- Annandale N (1918) Fish and fisheries of the Inlé Lake. Records of the Indian Museum 14: 33-64. <https://doi.org/10.5962/bhl.part.18603>
- Britz R (2003) *Danionella mirifica*, a new species of miniature fish from Upper Myanmar (Ostariophysi: Cyprinidae). Ichthyological Exploration of Freshwaters 14 (3): 217-222. URL: <http://zoobank.org/9107eb6f-4aec-49ac-8e91-3e729ab07e20>
- Clarke M (2007) New population of *Celestichthys* discovered. Practical Fishkeeping, Emap Active Limited, Peterborough, UK, 1 p..
- Clement M, Posada D, Crandall KA (2000) TCS: a computer program to estimate gene genealogies. Molecular Ecology 9 (10): 1657-1659. <https://doi.org/10.1046/j.1365-294x.2000.01020.x>
- Conway K, Chen W, Mayden R (2008) The “Celestial Pearl danio” is a miniature *Danio* (s.s) (Ostariophysi: Cyprinidae): evidence from morphology and molecules. Zootaxa 1686: 1-28. <https://doi.org/10.11646/zootaxa.1686.1.1>
- Hary C (2007) *Celestichthys margaritatus*: Perlhuhnbärblinge gibt es nicht nur in Burma. Die Aquarien- und Terrarienzeitschrift 60: 6.

- Hora SL, Mukerji DD (1934) Notes on fishes in the Indian Museum. XXII. On the collection of fish from the S. Shan States and the Pegu yomas, Burma. Records of the Indian Museum 36 (1): 123-138. URL: <http://zoobank.org/8d402a24-1057-48fb-8831-50d7ae635b88>
- Kano Y, Adnan MS, Grudpan C, Grudpan J, Magtoon W, Musikasinthorn P, Natori Y, Ottomanski S, Praxaysonbath B, Phongsa K, Rangsiruji A, Shibukawa K, Shimatani Y, So N, Suvannaraksha A, Thach P, Thanh PN, Tran DD, Utsugi K, Yamashita T (2013) An online database on freshwater fish diversity and distribution in mainland Southeast Asia. Ichthyological Research 60 (3): 293-295. <https://doi.org/10.1007/s10228-013-0349-8>
- Kano Y, Nakajima J (2014) Non-killing simple photography techniques of small–middle freshwater fishes in the field. Japanese Journal of Ichthyology 61: 123-125. [In Japanese].
- Kano Y, Musikasinthorn P, Iwata A, Tun S, Yun L, Win S, Matsui S, Tabata R, Yamasaki T, Watanabe K (2016) A dataset of fishes in and around Inle Lake, an ancient lake of Myanmar, with DNA barcoding, photo images and CT/3D models. Biodiversity Data Journal 4: e10539. <https://doi.org/10.3897/bdj.4.e10539>
- Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. Molecular Biology and Evolution 30 (4): 772-780. <https://doi.org/10.1093/molbev/mst010>
- Kullander S, Rahman MM, Norén M, Mollah AR (2017) *Devario* in Bangladesh: Species diversity, sibling species, and introgression within danionin cyprinids (Teleostei: Cyprinidae: Danioninae). PLOS One 12 (11). <https://doi.org/10.1371/journal.pone.0186895>
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. Molecular Biology and Evolution 33 (7): 1870-1874. <https://doi.org/10.1093/molbev/msw054>
- Leigh J, Bryant D (2015) POPART: full-feature software for haplotype network construction. Methods in Ecology and Evolution 6 (9): 1110-1116. <https://doi.org/10.1111/2041-210x.12410>
- Musikasinthorn P (1998a) *Channa panaw*, a new channid fish from the Irrawaddy and Sittang River Basins, Myanmar. Ichthyological Research 45 (4): 355-362. <https://doi.org/10.1007/bf02725188>
- Musikasinthorn P (1998b) First record of *Parambassis lala* (Pisces: Ambassidae) from Inle Lake, the Salween River basin, Myanmar. Natural History Bulletin of the Siam Society 46 (2): 203-205. URL: <http://zoobank.org/7024605f-bf0a-47ec-8659-9a69d00ca4a3>
- Okonechnikov K, Golosova O, Fursov M, the UGENE team (2012) Unipro UGENE: a unified bioinformatics toolkit. Bioinformatics 28 (8): 1166-1167. <https://doi.org/10.1093/bioinformatics/bts091>
- Roberts T (2007) The 'celestial pearl danio', a new genus and species of colourful minute cyprinid fish from Myanmar (Pisces: Cypriniformes). Raffles Bulletin of Zoology 55: 131-140.
- Roberts TR (2012) *Scleropages inscriptus*, a new fish species from the Tananthayi or Tenasserim River Basin, Malay Peninsula of Myanmar (Osteoglossidae: Osteoglossiformes). Aqua, International Journal of Ichthyology 18: 113-118.

- Talwar PK, Jhingran AG (1991) Inland fishes of India and adjacent countries. Vol.1. A.A. Balkema, Rotterdam, 541 pp. URL: <http://zoobank.org/5ea4ea8d-c5a9-40e7-95ad-3daa476a6ab3>
- Vidthayanon C, Termvidchakorn A, Pe M (2005) Inland fishes of Myanmar. Secretariat, Southeast Asian Fisheries Development Center, Bangkok, 160 pp.
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PN (2005) DNA barcoding Australia's fish species. Philosophical Transactions of the Royal Society B: Biological Sciences 360 (1462): 1847-1857. <https://doi.org/10.1098/rstb.2005.1716>
- Watanabe K, Kano Y, Takahashi H, Mukai T, Kakioka R, Tominaga K (2009) GEDIMAP: a database of genetic diversity for Japanese freshwater fishes. Ichthyological Research 57 (1): 107-109. <https://doi.org/10.1007/s10228-009-0129-7>
- Win SS (2018) Assessment on fishery sustainability in Inle wetland, Nyaung Shwe township, southern Shan State. International International Journal of Avian & Wildlife Biology 3 (5): 345-350. <https://doi.org/10.15406/ijawb.2018.03.00118>




Figure 1.  
A map of Hopong and the surrounding region.





Figure 2.

Photos of the spring field in Hopong. **A** A small dam holding spring water, inhabited by *Sawbwa resplendens* and *Oreochromis niloticus*. **B** A shallow dam backwater, inhabited by *Danio margaritatus* and *D. erythromicron*. **C** A stream flowing from the dam, inhabited by *Devario sondhii* and *Inlecypis auropurpureus*. **D** A ditch and shallow wetland, inhabited by *Physoschistura rivulicola* and *Channa harcourtbutleri*.




Figure 3.

The haplotype networks of five species groups ("species endemic to Inle Lake") generated with the TCS algorithm, based on COI regions (640 bp). Each bar on a branch corresponds to a single nucleotide substitution. Small dots represent hypothetical haplotypes. **A** *Inlecyptris auropurpureus* obtained from Hopong and two other localities. **B** *Microrasbora rubescens* obtained from Hopong and two other localities. The Heho specimens were obtained from a local market and their original locality is unknown. **C** *Physoschistura brunneana* and *P. rivulicola* obtained from Hopong and two other localities. The Heho specimens were obtained from a local market and their original locality is unknown. **D** *Sawbwa resplendens* obtained from Hopong and Inle Lake. **E** *Petruichthys brevis* obtained from Hopong and two other localities and *Petruichthys* sp. obtained from Hopong.

Table 1.

Fish species collected in a spring field in Hopong, the Salween Basin, Myanmar

Order	Family	Species	N (DNA barcoding)	Accession number
Cypriniformes				
	Cyprinidae	<i>Cyprinus rubrofasciatus</i> *	1 (0)	—
		<i>Danio erythromicron</i> †	33 (12)	LC190280, LC190284– LC190286, LC190314, LC645170–LC645175, LC645189
		<i>Danio margaritatus</i>	21 (9)	LC190268, LC190311– LC190313, LC190315, LC190395–LC190398
		<i>Devario brawni</i>	6 (4)	LC190317, LC190318, LC190320, LC645169
		<i>Devario sondhii</i>	21 (20)	LC190281, LC190289– LC190304, LC645177– LC645179
		<i>Esomus danrica</i> *	2 (1)	LC645176
		<i>Inlecypis auropurpureus</i> †	19 (13)	LC190269–LC190279, LC645167, LC645168
		<i>Microrasbora rubescens</i> †	66 (4)	LC190305–LC190308
		<i>Pethia stoliczkana</i>	30 (2)	LC190282, LC190283
		<i>Sawbwa resplendens</i> †	2 (1)	LC645184
		<i>Systemus</i> sp. cf. <i>rubripinnis</i>	3 (2)	LC190316, LC645185
		<i>Lepidocephalichthys berdmorei</i>	9 (3)	LC190324–LC190326
	Nemacheilidae	<i>Petruichthys brevis</i> †	18 (7)	LC190287, LC190288, LC190309, LC190310, LC645165, LC645166, LC645186
		<i>Petruichthys</i> sp. (Hopong)	2 (1)	LC645164
		<i>Physoschistura brunneana</i> †	5 (4)	LC190399–LC190402
		<i>Physoschistura rivulicola</i>	16 (9)	LC190319, LC190321– LC190323, LC190403– LC190405, LC645181, LC645182
Cyprinodontiformes				
	Poeciliidae	<i>Gambusia affinis</i> *	1 (1)	LC645188

		<i>Poecilia reticulata</i> *	1 (0)	—
Cichliformes				
	Ambassidae	<i>Parambassis lala</i> *	10 (2)	LC645180, LC645187
	Cichlidae	<i>Oreochromis niloticus</i> *	1 (0)	—
Beloniformes				
	Adrianichthyidae	<i>Oryzias uwai</i> *	11 (0)	—
Synbranchiformes				
	Synbranchidae	<i>Monopterus javanensis</i>	1 (1)	LC190383
	Mastacembelidae	<i>Mastacembelus caudiocellatus</i> †	2 (1)	LC645163
Anabantiformes				
	Osphronemidae	<i>Trichogaster labiosa</i> *	7 (0)	—
	Channidae	<i>Channa harcourtbutleri</i> †	30 (5)	LC190327–LC190330, LC645183
* Putative introduced species; † Assumed as endemic to Inle Lake				