

Current cave monitoring practices, their variation and recommendations for future improvement in Europe: A synopsis from the 6th EuroSpeleo Protection Symposium

Alexander M. Weigand^{‡,§,|}, Szilárd-Lehel Bücs[¶], Stanimira Deleva^{#,□}, Lada Lukić Bilela^{«,»}, Pierrette Nyssen[^], Kaloust Paragiamian[˘], Axel Ssymank^{!|}, Hannah Weigand^{‡,?}, Valerija Zakšek[˘], Maja Zagmajster[˘], Gergely Balázs[©], Shalva Barjadze[‡], Katharina Bürger[‡], William Burn^P, Didier Cailhol[^], Amélie Decrolière[«], Ferdinando Didonna^{F,|}, Azdren Doli[‡], Tvrtko Drazina^N, Joerg Dreybrodt^K, Lana Đud[©], Csaba Egri[?], Markus Erhard^W, Sašo Finžgar^T, Dominik Fröhlich[§], Grant Gartrell^{‡‡}, Suren Gazaryan^{§§}, Michel Georges^{||}, Jean-Francois Godeau[^], Ralf Grunewald^{¶¶}, John Gunn^{###}, Jeff Hajenga[□], Peter Hofmann^{««}, Lee Ronald Frederick Dennis Knight^{»»}, Hannes Köble[§], Nikolina Kuharic^N, Christian Lüthi^{^^}, Cristian Mihai Munteanu^{˘˘}, Rudjer Novak^{||,|}, Dainis Ozols^{??}, Matija Petkovic^{˘˘}, Fabio Stoch^{©©,||}, Bärbel Vogel^{§,|}, Ines Vukovic[©], Meredith Hall Weberg^{‡,‡}, Christian Zaenker^{§,PP}, Stefan Zaenker^{PP,§}, Ute Feit^{¶¶}, Jean-Claude Thies^{!|}

‡ Musée National d'Histoire Naturelle, Luxembourg, Luxembourg

§ Verband der deutschen Höhlen- und Karstforscher e. V., München, Germany

| European Cave Protection Commission (ECPC), Luxembourg, Luxembourg

¶ Centre for Bat Research and Conservation, Cluj-Napoca, Romania

National Museum of Natural History, Bulgarian Academy of Sciences, Sofia, Bulgaria

□ Sede del Sur, Universidad de Costa Rica, Golfito, Costa Rica

« University of Sarajevo, Faculty of Science, Sarajevo, Bosnia and Herzegovina

» Biospeleological Society in Bosnia and Herzegovina, Sarajevo, Bosnia and Herzegovina

^ Ecofirst scl - Union Belge de Spéléologie, Namur, Belgium

˘ Hellenic Institute of Speleological Research, Irakleion, Crete, Greece

! Bundesamt für Naturschutz, Bonn, Germany

? Fondation faune-flore, Luxembourg, Luxembourg

˘ University of Ljubljana, Ljubljana, Slovenia

© Eötvös Loránd University, Budapest, Hungary

‡ Ilia State University, Tbilisi, Georgia

‡ Verband Österreichischer Höhlenforscher, Vienna, Austria

P British Caving Association, Eyam, United Kingdom

^ French association of karstology, Pierrefontaine-lès-Blamont, France

« MaSePas, Namur, Belgium

F Italian Speleological Society, Castellana Grotte, Italy

‡ Mendelova Univerzita, Brno, Czech Republic

N Croatian Biospeleological Society, Zagreb, Croatia

K European Speleological Federation; Swiss Society of Speleology, Zurich, Switzerland

© Ministry of Economy and Sustainable Development, Zagreb, Croatia

? Ministry of Agriculture, Nagykovácsi, Hungary

W European Environment Agency (EEA), Copenhagen, Denmark

T Speleological Association of Slovenia, Ljubljana, Slovenia

‡‡ The Blueberry Patch, Mount Compass, Australia

§§ Eurobats, Bonn, Germany

|| Commission Wallonne d'Etude et de Protection des Sites Souterrains (CWEPSS) asbl, Bruxelles, Belgium

¶¶ Bundesamt für Naturschutz, Putbus, Germany

University of Birmingham, Birmingham, United Kingdom

□□ West Virginia Division of Natural Resources, Berkeley, United States of America

«« Verein für Höhlenkunde in München e. V., Brannenburg, Germany

»» British and Irish Hypogean Crustacea Recording Scheme, Exeter, United Kingdom

^^ Swiss Speleological Society, Interlaken, Switzerland

vy Emil Racoviță Institute of Speleology (Karstology, Karst Inventory and Protection Department); Romanian Academy of Sciences, Bucharest, Romania

ii University of Zagreb, Zagreb, Croatia

77 Nature Protection Agency, Sigulda, Latvia

55 Faculty of Biology, University of Belgrade, Belgrade, Serbia

çç Italian Speleological Society, Bologna, Italy

ll Université Libre de Bruxelles, Evolutionary Biology and Ecology, Bruxelles, Belgium

44 Virginia Cave Board, Richmond, United States of America

PP Landesverband für Höhlen- und Karstforschung Hessen e.V., Fulda, Germany

Corresponding author: Alexander M. Weigand (alexander.weigand@mnhn.lu)

Abstract

This manuscript summarizes the outcomes of the 6th EuroSpeleo Protection Symposium. Special emphasis was laid on presenting and discussing monitoring activities under the umbrella of the Habitats Directive (EU Council Directive 92/43/EEC) for habitat type 8310 "Caves not open to the public" and the Emerald Network. The discussions revealed a high level of variation in the currently conducted underground monitoring activities: there is no uniform definition of what kind of underground environments the "cave" habitat should cover, how often a specific cave has to be monitored, and what parameters should be measured to evaluate the conservation status. The variation in spatial dimensions in national definitions of caves further affects the number of catalogued caves in a country and the number of caves to be monitored. Not always participants are aware of the complete national monitoring process and that data sets should be freely available or easily accessible. The discussions further showed an inherent dilemma between an anticipated uniform monitoring approach with a coherent assessment methodology and, on the contrary, the uniqueness of caves and subterranean biota to be assessed – combined with profound knowledge gaps and a lack of resources. Nevertheless, some good practices for future cave monitoring activities have been identified by the participants: (1) Cave monitoring should focus on bio- and geodiversity elements alike; (2) Local communities should be involved, and formal agreements envisaged; (3) Caves must be understood as windows into the subterranean realm; (4) Touristic caves should not be excluded ad-hoc from regular monitoring; (5) New digital tools and open FAIR data infrastructures should be implemented; (6) Cave biomonitoring should focus on a large(r) biological diversity; and (7) DNA-based tools should be integrated. Finally, the importance of the 'forgotten' Recommendation No. 36 from the Bern Convention as a guiding legal European document was highlighted.

Keywords

cave monitoring, Habitats Directive, habitat type 8310, Emerald Network, Bern Convention, Recommendation 36

Date and place

26/ 10 - 30/10/2021; International Academy for Nature Conservation (INA), Isle of Vilm, Germany.

List of participants

Please refer to the complete list of authors. No differentiation is made between physical and virtual participants.

Introduction

This publication summarizes the outcomes of the 6th EuroSpeleo Protection Symposium entitled "*Assessing, monitoring and protecting cave biotopes and geotopes through Natura 2000 or similar programs in Europe*", held from 26th to 30th October 2021 at the International Academy for Nature Conservation (INA) on the Isle of Vilm (Germany). This hybrid event, jointly organized by the German Federal Agency for Nature Conservation (BfN) and the European Cave Protection Commission (ECPC) of the European Speleological Federation (FSE), gathered 45 participants from 22 countries (16 EU, 4 non-EU Europe, Australia and United States of America).

Aims of the workshop

A questionnaire was sent to the participants before the meeting, with the aim to generate a descriptive overview of the status quo of cave monitoring activities in the participating countries (Suppl. material 1). The questionnaire, as well as the symposium, had a strong focus on monitoring activities under the umbrella of the Habitats Directive (EU Council Directive 92/43/EEC) for habitat type 8310 "Caves not open to the public" and the Emerald Network. To a lesser extent the topics of monitoring activities for individual cave-dwelling species of Annexes II and IV of the Habitats Directive (i.e., Microchiroptera, *Proteus anguinus*, *Speleomantes* spp., *Conger* spp. and *Leptodirus hochenwartii*) or the habitat type 8330 "Submerged or partially submerged sea caves" were discussed.

Key outcomes and discussions

A high degree of variation in current cave monitoring practices

The participant's feedback received via the questionnaire revealed a high level of variation in the currently conducted underground monitoring activities. On the one hand, these differences result from the different framework conditions the Habitats Directive and the

Emerald Network set for monitoring activities. Generally, Natura 2000 sites within the Habitats Directive are considered as the contribution of the EU member states to the global Emerald Network. However, as a more significant discrepancy, there is only a recommendation for reporting in the Emerald Network (e.g. H1 "Terrestrial underground caves, cave systems, passages and waterbodies"). In contrast, there is an obligation to report on and explicitly assess the conservation status of habitat types 8310/8330 as well as Annex species in the six-year reporting cycle of the Habitats Directive (Art. 17). On the other hand, the surveyed countries showed very strong discrepancies in terms of who carries out the monitoring activities (independent experts, speleological societies, research institutes, universities, museums, NGOs, bat specialists, national parks, ministries or regional administrations) and what resources can be relied on (financial, infrastructural, personnel, pool of knowledge). At site level the "Natura 2000" standard data form is harmonized with Emerald-site descriptions, however there is no explicit ruling how often this needs to be updated. Here the local degree of conservation should be reported. Moreover, there is no uniform definition of what kind of underground environments the "cave" habitat should cover (Table 1), how often a specific cave has to be monitored (monthly to once each three years), and what parameters should be measured (fauna, flora[sic], abiotic parameters) to evaluate the conservation status. The variation in spatial dimensions in national definitions of caves further affects the number of catalogued caves in a country and the number of caves to be monitored. The survey results also showed that not all participants were aware of the complete national monitoring process and that data sets were not freely available or easily accessible. While in the EU the Natura 2000 implementation in the past largely focused on site selection and protection there is generally limited EU-guidance in monitoring systems (Ellwanger et al. 2018) and in some Member States these systems are still not fully established.

Conservation status of caves

Anecdotally, caves are just holes in the ground and maybe important for bats. And, they should be in generally good conservation status since nobody has access to them or it is physically too demanding to enter them. These typical misunderstandings are challenged by the fact(s) that caves are under strong human impact. Environmental pollution and sports, tourism and leisure activities are frequently identified as the main pressures and threats, which are often further intensified by illegal collections of cave-dwelling species and speleothems in some countries or specific caves (Prelovšek and Zupan Hajna 2011, Simičević 2017, Mammola 2018).

Trends of illegal cave-dwellers trade are particularly prominent in developing and transition countries, mostly due to ineffective local and regional management systems and insufficient interest in governing structures on this issue. In Dinaric Karst speleological objects, collecting and illegal trade mainly dates back to the 19th and early 20th centuries (Sket 2012). But unfortunately, the collection and illegal trade of mostly endangered, rare, and relict fauna, as well as of palaeontological remains (mostly *Ursus spelaeus*) still exists, e.g. in Bosnia and Herzegovina (Lukić Bilela and Jelić 2018), particularly in faunistic rich

caves such as Vjetrenica or Megara (in Bjelašnica Mt.) known as one of the largest *U. spelaeus* finding sites in the Western Balkans (Lukić-Bilela et al. 2013).

Another severe problem, especially in Vjetrenica cave, was the sampling of numerous specimens within (already small) metapopulations of rare species. With or without a collecting permit, these negative examples have been observed even among well-known and established scientists, who are expected to have high ethical principles. Furthermore, obtaining permits is relatively simple and generally does not contain mechanisms to control the sampling methodology and/or number of samples, nor are fines imposed for non-compliance. Thus, transition countries often become a twilight zone for illegally collecting valuable samples, with holotypes often ending outside the home country. Furthermore, most of these countries are not signatories to the Nagoya Protocol, and have no impact to any benefit-sharing, while their subterranean biodiversity is directly threatened.

Based on the Habitats Directive 2013-2018 assessments (EEA Web Tool 2021), the evaluation of the conservation status of H8310 caves varies from country to country and between biogeographical regions, but usually falls in the categories FV (Favourable) and U1 (Unfavourable-Inadequate). However, many country reports highlighted that the monitoring data and evaluations relied on expert opinions with a very limited amount of systematically collected data available. This lack of a systematic data collection allows on the one hand a high degree of freedom in implementation procedures, and the authorities, institutions and individual persons to get involved in national cave monitoring activities, but on the other hand also creates a significant heterogeneity in the evaluation of the conservation status between countries. This shows that cave monitoring to a large extent relies on expert volunteers and their interests and, thus, is strongly dependent on motivated and skilled experts (e.g., speleologists, researchers, naturalists) as well as available resources. The lack of financial and infrastructural support by national public administrations also creates a risk of fluctuating expert knowledge for cave monitoring and finally renders it extremely difficult to systematically collect, store and analyse data on a full scheme and long-term basis.

As a result, lacking adequate underlying data for monitoring can lead to erroneous conclusions regarding the conservation status of habitat 8310 or resident species, creating further inadequate conservation measures, or in case of a false favorable status, to no actions at all.

Prospects for future cave monitoring

Modern trends in conservation (speleo)biology prioritise the protection of endemic, relict or permanently endangered animal groups and their habitats, which necessitates an accurate assessment of the actual faunal inventory. Apart from being the basis for the preparation of Red Lists or Red Books of subterranean fauna, an adequate inventory is crucial for professionally guided monitoring activities of the often extremely vulnerable cave-dwelling organisms and subterranean ecosystems as a whole.

In line with a proposed conservation roadmap for the subterranean biome (Wynne et al. 2021), the symposium participants were guided by three basic questions: How to optimally monitor cave environments? How to evaluate the collected data? And finally, how to adequately define the conservation status of a cave? For the Habitats Directive, two principles of assessment are usually executed in parallel: loss or deterioration of the habitat and the distance to predefined favourable conservation status. The discussions showed an inherent dilemma between an anticipated uniform monitoring approach with a coherent assessment methodology and, on the contrary, the uniqueness of caves and subterranean biota to be assessed combined with profound knowledge gaps and a lack of resources. Nevertheless, seven good practices for future cave monitoring activities have been identified by the participants:

(1) Cave monitoring should focus on bio- and geodiversity elements alike

Caves harbour valuable bio- and geodiversity, and as such, both measures should be monitored and their entities protected. There was also a strong agreement that geodiversity elements guide biodiversity elements in underground environments and it is fundamental to secure them for effectively protecting cave-dwelling communities, which is in line with the recently adopted Resolution WCC-2020-Res-074-EN. An aspect that is intrinsically understood but not always explicitly formulated. Hence, a strong link between Life and Earth Sciences is needed to conduct monitoring activities, which often stays in contrast with established departmental and community structures. Goal-oriented integrative expert groups and networks should form, involving biologists, geologists, anthropologists, archaeologists, speleologists, speleodivers, and public administrations. A multidisciplinary community is needed to merge our understandings of caves being important sites of cultural heritage and endangered habitats of specific cave-dwelling communities (Wynne et al. 2021).

(2) Local communities should be involved, and formal agreements envisaged

Cave protection is a multi-layered process. The local community is physically, and often also emotionally, most closely involved. As such, local communities should be centrally involved in protecting caves, as no one can be more interested in protecting and improving their natural heritage (assets). This can be done e.g., by highlighting emblematic cave organisms (e.g., cave animal of the year; Vogel et al. 2017b) or fascinating speleothems and by active integration in the monitoring process, e.g. as conducted during the monitoring of special interest geological features at sites of special scientific interest (SSSI) in the UK. Training of local cavers in monitoring methods and conservation approaches (e.g., Romanian bat monitoring, Bücs et al. (2019)) can also create favorable conditions for active involvement in the future. Moreover, formal agreements for cave protection or memorandums of understanding should be aimed for, and a shared point-of-view of management plans developed by all stakeholders involved. A good example here originates from the Vjetrenica cave, as one of the faunistically richest caves in the world (Culver et al. 2021; Ozimec and Lucic 2009) with 219 species recorded (incl. 41 new for science) and being the reference site for the definition of a novel habitat type - the cave

hygropetric (Sket 2004, Ozimec 2021). It has been declared a Protected Landscape and nominated for inscription on the UNESCO World Heritage List in 2021. This exceptional bio- and geodiversity also requires careful project management such as preparation of planning documents (strategy, program, and management plans), planning of future research but also financing and continuous work on promotion and education, with strong inclusion of the local community and the Municipality of Ravno, besides the governmental bodies. The cave is managed by the Public company "Vjetrenica", but the entire local community is directly or indirectly involved in cave protection and community prosperity. This includes working closely with speleologists, scientists from different fields of research, experts and business people. The local community is turning to sustainable tourism, sustainable agriculture and organic production to avoid toxic organophosphorus pesticides, supports scientific expeditions in the underground of Popovo polje (such as donations of local small business owners in food and other necessities). They also take an active part in different development projects in the Municipality of Ravno such as Vjetrenica infrastructure arrangement, revitalisation of the narrow railway and many other projects that directly contribute to sustainable tourism, agriculture and development. The local community supported the provision of space for the establishment of the Vjetrenica Biospeleological Museum, the first such museum in this part of Europe, and is actively working to create conditions for new development ideas that will enable further research in Vjetrenica cave but all other caves in Popovo polje.

(3) Caves must be understood as windows into the subterranean realm

Caves allow humans to more or less easily enter underground environments, and to act in them. However, caves as an anthropocentric term must be understood and seen as windows into the subterranean realm (Mammola 2018). As such, monitoring activities within caves and the protection of a cave should be only the beginning of environmental protection, and it is of paramount importance to save the whole ecosystem by preserving nearby aquifers and surface areas. The specific fauna of groundwater aquifers also has many local endemic species and their distribution is not linked to typical distribution patterns of above-ground biodiversity (Deharveng et al. 2009).

(4) Touristic caves should not be excluded ad-hoc from regular monitoring

The habitat type 8310 of the Habitats Directive excludes touristic caves from legal monitoring obligations. However, touristic caves are often the best-studied and understood underground environments in a country (Baradla Domica cave system in Hungary; Frasassi caves in Italy; Vjetrenica cave in Bosnia and Herzegovina; Grotte de Han in Belgium; Meziad cave in Romania) and harbour at least equally rich bio- and geodiversity elements compared to non-touristic caves (Culver and Sket 2002, Culver et al. 2021). Additionally, only small(er) parts of a touristic cave usually are open to the broader public with large(r) parts excluded from regular visits. Hence, the integration of touristic caves into regular monitoring activities seems logical, as there exists no valid scientific reason for their ad-hoc exclusion. At least, the non-touristic parts of caves open to the public could be considered as habitat type 8310. Many touristic caves play also an important role for bat

species of Annexes II and IV of the Habitats Directive, and in this case monitoring of the bat species including their habitat is necessary anyway.

(5) New digital tools and open FAIR data infrastructures should be implemented

The FAIR principles of data management and stewardship should be followed to more effectively use and interconnect existing information, i.e. fostering the **F**indability, **A**ccessibility, **I**nteroperability, and **R**euse of digital assets (Wilkinson et al. 2016). Good examples for open (bio)speleological data e.g. comprise the CroSpeleo database from Croatia (Institute for Environment and Nature of the Ministry of Economy and Sustainable Development 2022), the Cave Fauna of Greece Database (Paragamian et al. 2018), the Database of Romanian cave invertebrates (Moldovan et al. 2020), the SubBIOCODE database from Bosnia and Herzegovina (SubBioLab 2022) and the Cave Biodiversity of Georgia database (Barjadze et al. 2019). Furthermore, new smart tools such as the German CaveLife app (Vogel et al. 2017a) for ecological assessments of underground environments can be utilised to ensure direct in-field digital data entry.

(6) Cave biomonitoring should focus on a large(r) biological diversity

The protection of the subterranean realm and fauna of a region always begins with the protection of certain caves and groups or organisms (Culver and Sket 2002). However, subterranean habitats are often only monitored for their suitability as an important hibernation and nursery place for bats. Furthermore, the listed subterranean non-microchiropteran species in Annexes II and IV of the Habitats Directive are eutroglobiont low-range endemics and of high conservation concern for very few European countries. The direct link to the monitoring of the more complex cave-dwelling fauna is not explicitly made, although the description of habitat type 8310 states that the "specialised fauna" should be taken into account. It is a gap in the species Annexes of the Habitats Directive that at the time of their status nascendi several species groups were not considered, due to missing data and or understanding. However, many of these species are of European importance and fulfil all legal criteria of the Directive for their inclusion into the Annexes. Therefore, once an update of Annexes is foreseen in future, a systematic scan of overlooked taxa to include at least a few indicator species would help.

Underground habitats represent complex ecosystems, with specific trophic transfers and food-webs (Saccò et al. 2019, Zaenker et al. 2020, Premate et al. 2021). Yet, to understand and protect underground environments as habitats of taxonomically rich and seasonally variable communities (incl. members of the specialised cave fauna, seasonally cave-dwelling and epigeal animal taxa, as well as bacteria, protists and fungi), future biodiversity monitoring should go beyond the current limited faunistic elements and selected eutroglobiont species. The huge spectrum of cavernicolous fauna should be taken as a major part of the typical species inventory of the cave habitat, intrinsically be monitored and assessed as a "structures and functions" parameter (which includes typical species) and thus be understood as obligatory part of the Habitat assessment of Cave

habitat types 8310 and 8330. Typical species as such have to be reported as a list to the EU as part of the national reports every 6 years, however clear rules on how to integrate them into assessments are still missing.

(7) DNA-based tools should be integrated

Faunistic assessments of subterranean communities can widely benefit from more widespread implementation of DNA-based tools. The digital taxonomic knowledge deposited in DNA barcode reference libraries can be freely utilised, without the direct need or immediate availability of a morpho-taxonomic expert. Undeterminable developmental stages or complicated genders can be readily identified and members within cryptic species complexes, i.e. biological species currently taxonomically lumped under a single name, can be more precisely told apart, which is an often highlighted conservation concern for subterranean biota (Trontelj et al. 2009, Delić et al. 2017, Fišer et al. 2018, Malard et al. 2018). Furthermore, environmental DNA (eDNA) approaches can help to improve the knowledge on endangered subterranean target species, especially in difficult to reach underground environments (Saccò et al. 2022). Specific eDNA-based assays have been developed e.g. for the Cave Salamander *Proteus anguinus* (Gorički et al. 2017, Vörös et al. 2017), the Alabama Cave Crayfish *Cambarus speleocoopi* (Boyd et al. 2020) and Hay's Spring Amphipod *Stygobromus hayi* and its more widespread congener *S. tenuis potomacus* (Niemiller et al. 2017). Finally, DNA or eDNA metabarcoding-based surveys can help to reach a more comprehensive understanding of the subterranean community in a given ecosystem, e.g., by increasing taxonomic resolution and including so far underrepresented taxa such as fungi, protists, diatoms, bacteria, cyanobacteria, algae and mosses (Sohlberg et al. 2015, Miettinen et al. 2015, Nawaz et al. 2018, Korbelt et al. 2017, Pfendler et al. 2018, Alaoui-Sosse et al. 2021). Metabarcoding approaches have been also utilised to characterise the diets of bats (Galan et al. 2018) and to identify ancient rodent remains (Guimaraes et al. 2016).

Conclusion

Cave monitoring practices are far from being standardised in Europe. On the one hand, this is due to the fact that variable resources and levels of support are available to the nations, and on the other hand, imposed by the fact that central parameters of cave monitoring are poorly described in the Directive 92/43/EEC and very poorly interpreted in the EU Habitat Interpretation Manual (e.g. definition of "cave", definition "not open to the public", term "specialised species", term "endemic species"). Although, and due to natural constraints, cave monitoring needs to have a certain level of national flexibility to account for the variation of bio- and geodiversity elements, some basic standardisation guidelines and open data repositories are needed to compare data across space and time.

The most important European legal document on the conservation of underground habitats is the Recommendation No. 36 (Bern Convention 1992). This recommendation dates back to 1992, asking the Member States to list underground invertebrates and underground habitats that need to be protected and to proceed with their legal protection. The initial

documentation was provided by Christian Juberthie (1931-2019) and 24 distinguished speleobiologists from 20 European countries (Juberthie 1992). Unfortunately, the Recommendation remains unimplemented in the majority of Member States (Haslett 2007), and became widely forgotten in the European cave monitoring community. A first important step would be to revive the contents of the Recommendation No. 36, including criteria for selecting underground habitats of biological value and proposals for procedures of protection and management of underground habitats, in order to have a common denominator as a basis for future cave monitoring activities.

References

- Alaoui-Sosse B, Ozaki S, Barriquand L, De Luca D, Cennamo P, Vallot B, Alaoui-Sosse L, Bourgeade P, Boustia F, Aleya L, Pfendler S (2021) Does Microbial Diversity of Cave Ecosystems Differ from Outside? The Case of the Azé Show Cave (France). *Research Square* <https://doi.org/10.21203/rs.3.rs-471958/v1>
- Barjadze S, Arabuli T, Mumladze L, Maghradze E, Asanidze Z, Shavadze L, Gogshelidze M (2019) Cave Biodiversity of Georgia - Open Access Database. <https://cbg.iliauni.edu.ge/en/>. Accessed on: 2022-3-01.
- Bern Convention (1992) Recommendation No. 36, on the conservation of underground habitats. Council of Europe, Strasbourg. URL: https://search.coe.int/bern-convention/Pages/result_details.aspx?ObjectId=0900001680746962
- Boyd SH, Niemiller KDK, Dooley KE, Nix J, Niemiller ML (2020) Using environmental DNA methods to survey for rare groundwater fauna: Detection of an endangered endemic cave crayfish in northern Alabama. *PLOS ONE* 15 (12). <https://doi.org/10.1371/journal.pone.0242741>
- Bűcs SL, Gőnczi VI, Szigeti M, Jumanca M, Onodi H, Bodea F, Gherghel N, Ghercea A, Lucaci L, Ricman R, Dumbravă AR, Sándor DA, Barti L, Jére C (2019) Results of chiro-speleo collaboration in the 2016-2019 period in Romanian caves. National Caver Congress, Șuncuiuș, Romania.
- Culver DC, Sket B (2002) Biological Monitoring in Caves. *Acta Carsologica* 31 (1): 55-64. <https://doi.org/10.3986/ac.v31i1.403>
- Culver DC, Deharveng L, Pipan T, Bedos A (2021) An Overview of Subterranean Biodiversity Hotspots. *Diversity* 13 (10). <https://doi.org/10.3390/d13100487>
- Deharveng L, Stoch F, Gibert J, Bedos A, Galassi D, Zagmajster M, Brancelj A, Camacho A, Fiers F, Martin P, Giani N, Magniez G, Marmonier P (2009) Groundwater biodiversity in Europe. *Freshwater Biology* 54 (4): 709-726. <https://doi.org/10.1111/j.1365-2427.2008.01972.x>
- Delić T, Trontelj P, Rendoš M, Fišer C (2017) The importance of naming cryptic species and the conservation of endemic subterranean amphipods. *Scientific Reports* 7 (1). <https://doi.org/10.1038/s41598-017-02938-z>
- EEA Web Tool (2021) Biogeographical assessments of conservation status of species and habitats under Article 17 of the Habitats Directive . <https://nature-art17.eionet.europa.eu/article17/>. Accessed on: 2022-3-01.
- Ellwanger G, Runge S, Wagner M, Ackermann W, Neukirchen M, Frederking W, Müller C, Ssymank A, Sukopp U (2018) Current status of habitat monitoring in the European Union according to Article 17 of the Habitats Directive, with an emphasis on habitat

- structure and functions and on Germany. *Nature Conservation* 29: 57-78. <https://doi.org/10.3897/natureconservation.29.27273>
- Eriņš G (2004) *Alas Latvijā*. [Caves in Latvia]. Zvaigzne ABC, Riga, 76 pp.
 - Fišer C, Robinson CT, Malard F (2018) Cryptic species as a window into the paradigm shift of the species concept. *Molecular Ecology* 27 (3): 613-635. <https://doi.org/10.1111/mec.14486>
 - Galan M, Pons JB, Tournayre O, Pierre É, Leuchtman M, Pontier D, Charbonnel N (2018) Metabarcoding for the parallel identification of several hundred predators and their prey: Application to bat species diet analysis. *Molecular Ecology Resources* 18 (3): 474-489. <https://doi.org/10.1111/1755-0998.12749>
 - Gorički Š, Stanković D, Snoj A, Kuntner M, Jeffery WR, Trontelj P, Pavičević M, Grizelj Z, Năpăruș-Aljančić M, Aljančić G (2017) Environmental DNA in subterranean biology: range extension and taxonomic implications for *Proteus*. *Scientific Reports* 7 (1). <https://doi.org/10.1038/srep45054>
 - Guimaraes S, Pruvost M, Daligault J, Stoetzel E, Bennett EA, Côté NM, Nicolas V, Lalis A, Denys C, Geigl E-, Grange T (2016) A cost-effective high-throughput metabarcoding approach powerful enough to genotype ~44 000 year-old rodent remains from Northern Africa. *Molecular Ecology Resources* 17 (3): 405-417. <https://doi.org/10.1111/1755-0998.12565>
 - Haslett JR (2007) European Strategy for the conservation of invertebrates. Council of Europe, *Nature and Environment* 145: 1-91.
 - Institute for Environment and Nature of the Ministry of Economy and Sustainable Development (2022) Nature protection Information System – CroSpeleo (Cave registry of the Republic of Croatia). <https://crospeleo.mingor.hr/>. Accessed on: 2022-3-01.
 - Juberthie C (1992) *Underground Habitats and Their Protection*. Council of Europe, Strasbourg, 166 pp.
 - Korbel K, Chariton A, Stephenson S, Greenfield P, Hose GC (2017) Wells provide a distorted view of life in the aquifer: implications for sampling, monitoring and assessment of groundwater ecosystems. *Scientific Reports* 7 (1). <https://doi.org/10.1038/srep40702>
 - Lukić Bilela L, Jelić D (2018) *Caves and other Subterranean Habitats*. In: Brūmelis G, Troumbis A (Eds) Chapter 3: Status, trends and future dynamics of biodiversity and ecosystems underpinning nature's contributions to people; Chapters of the regional assessment report on biodiversity and ecosystem services for Europe and Central Asia of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. 320-326 pp. URL: <https://www.ipbes.net/assessment-reports/eca>
 - Lukić-Bilela L, Ozimec R, Miculinić K, Basara D (2013) A comprehensive valorisation of Megara cave with a view to preservation and protection. *Natura Montenegrina* 12 (3): 1-1.
 - Malard F, Grison P, Eme D, Fiser C, Flot J-, Konecny-Dupré L, Lefébure T, Stoch F, Zagmajster M, Douady C (2018) Integrating multiple species criteria and species hypotheses in subterranean biology. ARPHA Conference Abstracts 1 <https://doi.org/10.3897/aca.1.e30399>
 - Mammola S (2018) Finding answers in the dark: caves as models in ecology fifty years after Poulson and White. *Ecography* 42 (7): 1331-1351. <https://doi.org/10.1111/ecog.03905>

- Miettinen H, Kietäväinen R, Sohlberg E, Numminen M, Ahonen L, Itävaara M (2015) Microbiome composition and geochemical characteristics of deep subsurface high-pressure environment, Pyhäsalmi mine Finland. *Frontiers in Microbiology* 6 <https://doi.org/10.3389/fmicb.2015.01203>
- Moldovan OT, Iepure S, Brad T, Kenesz M, Mirea IC, Năstase-Bucur R (2020) Database of Romanian cave invertebrates with a Red List of cave species and a list of hotspot/coldspot caves. *Biodiversity Data Journal* 8 <https://doi.org/10.3897/bdj.8.e53571>
- Nawaz A, Purahong W, Lehmann R, Herrmann M, Totsche KU, Küsel K, Wubet T, Buscot F (2018) First insights into the living groundwater mycobiome of the terrestrial biogeosphere. *Water Research* 145: 50-61. <https://doi.org/10.1016/j.watres.2018.07.067>
- Niemiller ML, Porter ML, Keany J, Gilbert H, Fong DW, Culver DC, Hobson CS, Kendall KD, Davis MA, Taylor SJ (2017) Evaluation of eDNA for groundwater invertebrate detection and monitoring: a case study with endangered *Stygobromus* (Amphipoda: Crangonyctidae). *Conservation Genetics Resources* 10 (2): 247-257. <https://doi.org/10.1007/s12686-017-0785-2>
- Ozimec R, Lucic I (2009) The Vjetrenica cave (Bosnia & Herzegovina) – one of the world's most prominent biodiversity hotspots for cave-dwelling fauna. *Subterranean Biology* 7: 17-24.
- Ozimec R, et al. (2021) Vjetrenica – Centar špiljske bioraznolikosti Dinarida . [Vjetrenica – Cave Biodiversity Hotspot of the Dinarides]. JP Vjetrenica – ADIPA, 356 pp. [ISBN 9789926856403]
- Paragamian K, Poulinakis M, Paragkamian S, Nikoloudakis I (2018) Cave fauna of Greece database - Hellenic Institute of Speleological Research. <https://database.inspee.gr/>. Accessed on: 2022-3-01.
- Pfendler S, Karimi B, Maron P-, Ciadamidaro L, Valot B, Bousta F, Alaoui-Sosse L, Alaoui-Sosse B, Aleya L (2018) Biofilm biodiversity in French and Swiss show caves using the metabarcoding approach: First data. *Science of The Total Environment* 615: 1207-1217. <https://doi.org/10.1016/j.scitotenv.2017.10.054>
- Prelovšek M, Zupan Hajna N (2011) Pressures and protection of the underground karst –cases from Slovenia and Croatia. Karst Research Institute ZRC SAZU, Postojna. [ISBN 9789612542856]
- Premate E, Zagamjster M, Fiser C (2021) Inferring predator-prey interaction in the subterranean environment: a case study from Dinaric caves. *Scientific reports* 11 (1): 1-9. <https://doi.org/10.1038/s41598-021-01249-8>
- Saccò M, Blyth A, Bateman PW, Hua Q, Mazumder D, White N, Humphreys WF, Laini A, Griebler C, Grice K (2019) New light in the dark - a proposed multidisciplinary framework for studying functional ecology of groundwater fauna. *Science of The Total Environment* 662: 963-977. <https://doi.org/10.1016/j.scitotenv.2019.01.296>
- Saccò M, Guzik MT, van der Heyde M, Nevill P, Cooper SJ, Austin AD, Coates PJ, Allentoft ME, White NE (2022) eDNA in subterranean ecosystems: Applications, technical aspects, and future prospects. *Science of The Total Environment* 820 <https://doi.org/10.1016/j.scitotenv.2022.153223>
- Simičević V (2017) Poachers threaten Balkans' underground biodiversity. *Science* 358 (6367): 1116-1117. <https://doi.org/10.1126/science.358.6367.1116>
- Sket B (2004) The cave hygropetric - a little known habitat and its inhabitants. *Archiv für Hydrobiologie* 160 (3): 413-425. <https://doi.org/10.1127/0003-9136/2004/0160-0413>

- Sket B (2012) *Diversity patterns in Dinaric karst*. In: White WB, Culver D (Eds) *Encyclopedia of Caves*. 2. Academic Press, 228-238 pp. <https://doi.org/10.1016/B978-0-12-383832-2.00031-1>
- Sohlberg E, Bomberg M, Miettinen H, Nyssönen M, Salavirta H, Vikman M, Itävaara M (2015) Revealing the unexplored fungal communities in deep groundwater of crystalline bedrock fracture zones in Olkiluoto, Finland. *Frontiers in Microbiology* 6 <https://doi.org/10.3389/fmicb.2015.00573>
- Ssymank A, et al. (in press) Das europäische Schutzgebietssystem Natura 2000. BfN-Handbuch zur Umsetzung der Fauna-Flora-Habitat-Richtlinie (92/43/EWG) und der Vogelschutzrichtlinie (2009/147/EG). Band 2.2: Lebensraumtypen der Meere und Küsten, der Binnengewässer sowie der Heiden und Gebüsche. - 2. A. Naturschutz und Biologische Vielfalt 172.
- SubBioLab (2022) SubBIOCODE database. <https://db.subbiocode.net/>. Accessed on: 2022-3-01.
- Trontelj P, Douady CJ, Fiser C, Gibert J, Goricki Š, Lefébure T, Sket B, Zaksek V (2009) A molecular test for cryptic diversity in ground water: how large are the ranges of macro-stygobionts? *Freshwater Biology* 54 (4): 727-744. <https://doi.org/10.1111/j.1365-2427.2007.01877.x>
- Vogel B, Zaenker S, Zaenker C, Ipsen A, Wolf A, Knolle F (2017a) Smart Monitoring of Cave Habitats - the VdHK Smartphone App. In: Moore K, White S (Eds) *Proceedings of the 17th International Congress of Speleology*, 1. 17th International Congress of Speleology, Sydney, 22.-28.07.2017. Australian Speleological Federation Inc, Sydney [ISBN 978-0-9808060-5-2-07-6].
- Vogel B, Zaenker S, Zaenker C, Ipsen A, Wolf A, Knolle F (2017b) Cave Animal Of The Year – From National To International? In: Moore K, White S (Eds) *Proceedings of the 17th International Congress of Speleology*, 1. 17th International Congress of Speleology, Sydney, 22.-28.07.2017. Australian Speleological Federation Inc, Sydney [ISBN 978-0-9808060-5-2-07-6].
- Vörös J, Márton O, Schmidt BR, Gál JT, Jelić D (2017) Surveying Europe's Only Cave-Dwelling Chordate Species (*Proteus anguinus*) Using Environmental DNA. *PLOS ONE* 12 (1). <https://doi.org/10.1371/journal.pone.0170945>
- Wilkinson M, Dumontier M, Aalbersberg IJ, et al. (2016) The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data* 3: 160018. <https://doi.org/10.1038/sdata.2016.18>
- Wynne JJ, Howarth FG, Mammola S, Ferreira RL, Cardoso P, Di Lorenzo T, Galassi DMP, Medellín RA, Miller BW, Sánchez-Fernández D, Bichuette ME, Biswas J, BlackEagle CW, Boonyanusith C, Amorim IR, Borges PAV, Boston PJ, Cal RN, Cheeptham N, Deharveng L, Eme D, Faille A, Fenolio D, Fišer C, Fišer Ž, 'Ohukani'ōhi'a Gon SM, Goudarzi F, Griebler C, Halse S, Hoch H, Kale E, Katz AD, Kováč L, Lilley TM, Manchi S, Manenti R, Martínez A, Meierhofer MB, Miller AZ, Moldovan OT, Niemi ML, Peck SB, Pellegrini TG, Pipan T, Phillips-Lander CM, Poot C, Racey PA, Sendra A, Shear WA, Silva MS, Taiti S, Tian M, Venarsky MP, Pakarati SY, Zagmajster M, Zhao Y (2021) A conservation roadmap for the subterranean biome. *Conservation Letters* 14 (5). <https://doi.org/10.1111/conl.12834>
- Zaenker S, Bogon K, Weigand A (2020) *Die Höhlentiere Deutschlands: Finden – Erkennen – Bestimmen*. Quelle & Meyer, 448 pp. [ISBN 3494018316]

Table 1.

List of selected legal definitions of "a cave" for monitoring purposes.

| Geographical coverage | Definition of a cave | Defined spatial criteria of a cave | Legal source |
|---|--|---|---|
| EU member states | Caves not open to the public, including their water bodies and streams, hosting specialised or high endemic species, or that are of paramount importance for the conservation of Annex II species (e.g. bats, amphibians) | Not defined | Interpretation Manual of European Union Habitats, version EUR 28 |
| Bosnia and Herzegovina (Entity: Federation of BiH and Entity: Republic of Srpska) | Speleological objects are naturally formed underground cavities longer than 5 meters, that can be entered by man, and the dimensions of the entrance are less than the depth or length of object (caves, pits, abysses, estavels, etc.) | More than 5 m long; object can be entered by man, with dimensions of the entrance less than the depth or length of object | Zakon o zaštiti prirode („Službene novine Federacije BiH“, broj:66/13); and Zakon o zaštiti prirode (“Službeni glasnik Republike Srpske, broj:113/08) |
| Bulgaria | The cave is an underground cavity with a volume, big enough to fit a man | Big enough to fit a man | Source used in the speleology manual: Malcolm S. Field (2002) A Lexicon of Cave and Karst Terminology with Special Reference to Environmental Karst Hydrology. Office of Research and Development, U.S. Environmental Protection Agency, Washington, DC |
| Croatia | Speleological objects/spaces are naturally formed underground spaces (caves, pits, abysses, estavelles, etc.), as well as their parts | Not defined | Point 46, paragraph 1, article 9 of the Nature protection act (Official Gazette, No. 80/2013, 15/2018, 14/2019 and 127/19) |
| Germany | Caves, including their waterbodies, are usually inhabited by specialized typical species (cavernicolous fauna), which may include rare local or regional endemics, restricted to one or a few cave systems. Semi-caves ("Balmen") are included if they have specific vegetation or fauna | Not defined | Legal source refers only to EU Habitats Directive itself, BNatSchG § 31-36 (Bundesnaturschutzgesetz, framework law) at National level, different implementations in the 16 Federal states with additional laws on nature conservation (Landesnaturschutzgesetze). See also Ssymank (in press) |
| Greece | Caves are not defined/ interpreted in Greece's legislation | Not defined | Not available |

| | | | |
|------------|--|--|---|
| Hungary | Cave is a natural cavity formed in the earth's crust, which exceeds the length of 2 meters along its longitudinal axis and - after the removal of its current or natural deposit - it is possible for a human to enter | More than 2 m long (longitudinal axis); broad enough to enter | 1996. évi LIII. törvény a természet védelméről (Law on nature protection), Tvt. 23§ |
| Italy | Both natural and artificial cavities are not defined/ interpreted in Italy's legislation. They are not protected at the origin but only by special laws on extractive activities, mineral waters, archaeology and fine arts, or by parks, reserves and regional laws | Not defined | Underground mines stay under the dictates of the D.P.R. no. 128 of 09/04/1959 (Police regulations of mines and quarries, updated with Legislative Decree no. 624/96) while cavities of historical-archaeological interest or of particular geological singularity are protected by the Code of Cultural Heritage and Landscape (Legislative Decree 22.01.2004 no. 42) |
| Latvia | Natural or naturalised subterranean entity within bedrock or cemented Quaternary sediments more than 3 m long and broad enough to enter | More than 3 m long; broad enough to enter; length > width of the entrance | Page 12 in Eniņš (2004) |
| Luxembourg | Caves and semi-caves not used for tourism, including their water bodies. Requirements: 1) naturally formed underground cavity with a dark zone. 2) minimum size: cavities into which a person can enter, larger than 5m ³ , with a depth of at least 5m. 3) animal burrows are explicitly excluded | At least 5 m long; broad enough to enter; volume larger than 5m ³ | Monitoring sheet and national interpretation manual |
| Romania | The cave is a natural cavity formed in the earth's crust, wide and deep enough to allow a human to enter. By extension, the cave can be a system that can have more than one entrance and consists of several galleries, halls, wells and chimneys. The term also applies to the cavities defined above, which are partially or completely submerged or which have been opened by demolition or clearing | Wide and deep enough to allow a human to enter | Art. 43(3) from Governmental Emergency Ordinance 57/2007, approved by Law 49/2011 |

Supplementary material

Suppl. material 1: Questionnaire for the 6th EuroSpeleo Protection Symposium 2022_

Authors: Jean-Claude Thies, European Cave Protection Commission (ECPC)

Data type: questionnaire

Brief description: *This questionnaire was mandatory for each applicant. It should yield important information about the Natura 2000 monitoring and conservation of cave habitats in the country of origin. The information derived was presented at the Symposium and was an important foundation for discussion.*

[Download file](#) (1.54 MB)