

Ticks on the Edible Dormouse (*Glis glis*)

Karolína Srbová[‡], Pavlína Paclíková[‡], Luděk Žůrek[§], Peter Adamik[‡]

[‡] Department of Zoology, Palacký University Olomouc, Olomouc, Czech Republic

[§] University of Veterinary and Pharmaceutical Sciences Brno, Brno, Czech Republic

Corresponding author: Karolína Srbová (karolina.srbova1@gmail.com)

Abstract

Rodents in central Europe, such as the Edible Dormouse, are common hosts for many ectoparasites, including ticks. One of the most extensively studied tick species is *Ixodes ricinus*, whose geographic distribution extends from Scandinavia to Morocco. In 2014, a new North African species *I. inopinatus* was described and found in Morocco, Tunisia, the Iberian Peninsula, and peripherally also in Romania, Austria, and southern Germany. Both *Ixodes* species were reported in sympatry in southern Europe (Spain and Portugal) where they feed mainly on lizards. During the period from early June to October in 2016-2019, we captured 4597 dormice and calculated the rate of tick infestation. Our results show that the mean prevalence of ticks in the dormouse population was 31.9 %. In the next step, 400 ticks from 2019 (255 larvae, 144 nymphs, 1 adult) from dormice were identified morphologically and by multiplex-PCR to distinguish *I. ricinus* and *I. inopinatus*. The results of our analysis show the first report of *I. inopinatus* feeding on rodents.

Keywords

dormouse, ticks, rate of infestations, crossbreeding

Presenting author

Karolína Srbová

Presented at

Oral presentation at the 11th International Dormice Conference 2022 (May 9 – 13, 2022)

Conflicts of interest