

Movement Paradigm for Hazara Torrent Frog *Allopaa hazarensis* and Murree Hills Frog *Nanorana vicina* (Anura: Dicroglossidae)

Ayesha Akram[‡], Muhammad Rais[§], Muhammad Saeed[‡], Waseem Ahmed[‡], Sumbul Gill[‡], Jibran Haider[‡]

[‡] Herpetology Lab, Department of Zoology, Wildlife and Fisheries, Pir Mehr Ali Shah-Arid Agriculture University Rawalpindi, Rawalpindi 46000, Pakistan, Rawalpindi, Pakistan

[§] PMAS Arid Agriculture University Rawalpindi, Rawalpindi, Pakistan

| Gilgit-Baltistan Forest, Wildlife and Environment Department, Gilgit 15100, Pakistan, Gilgit, Pakistan

Corresponding author: Muhammad Rais (sahil@uaar.edu.pk)

Academic editor: Truong Nguyen

Abstract

Endemic anurans are particularly vulnerable to environmental changes, and are susceptible to population declines because of their restricted distribution ranges. The Murree Hills Frog *Nanorana vicina* and Hazara Torrent Frog *Allopaa hazarensis* are associated with the torrential streams and nearby clear water pools in subtropical chir pine forest and other forest types, at elevations higher than 1000 m in Pakistan. In this study, we have provided data on the extent of movement of these frog species for the first time. We installed radio transmitters on a total of 13 Murree Hills Frogs and 13 Hazara Torrent Frogs during eight consecutive days in September 2017 and 2018. Our results showed that these frogs did not move long distances along the stream or away from the stream into the forest. All the radio-tracked frogs showed movement of < 3 m. We found a significant differences only in the distance moved by Murree Hills Frogs between the two years studied. Based on our findings, we propose a movement paradigm that focuses on conservation implications for these endemic frogs.

Keywords

conservation, dispersal, endemic, frogs, habitat, radio telemetry

Introduction

Conservation of amphibians is becoming more vital due to the increasing global extinction rate in this group (Rais et al. 2021). Amphibians respond to geophysical characteristics at broad spatial scales, but few studies have examined their response to changes in

landscape structure and climate change (Dupuis et al. 2000, Adams and Bury 2002). The ultimate structure of amphibian populations depends on the success of dispersal, spatial distribution of water bodies and connectivity of breeding sites (Semlitsch and Bodie 1998, Marsh and Trenham 2001, Moilanen and Nieminen 2002). Structural and functional landscape connectivity is essential for dispersal of species across the landscape (Taylor et al. 2006).

Studies on habitat use provide useful information for species conservation (Manly et al. 2002, Brunjes et al. 2006, Aarts et al. 2008). Endemic amphibians are particularly vulnerable to environmental changes and are susceptible to population declines (Lecis and Norris 2004, Moore et al. 2004) because of their restricted distribution range. The Murree Hills Frog and Hazara Torrent Frog are endemic to South Asia and Pakistan, respectively (Khan 2006, Rais et al. 2021). These frogs are associated with the torrential streams and nearby clear water pools in subtropical chir pine forest and Himalayan moist temperate forest at elevations higher than 1000 m (Ahmed et al. 2020), and their breeding season is from July-August (Saeed et al. 2021). The two species are categorized as Least Concern by the Red List of Threatened Species by the International Union for Conservation of Nature (Ahmed et al. 2020). Habitat degradation, urbanization and climate change are the known threats to these species (Ohler and Dutta 2004, Khan et al. 2008). Currently, there are no published data on the movement of these species. Given this, the present study used radio telemetry to assess for the first time the movements of *Allopaa hazarensis* and *Nanorana vicina* and proposed a movement paradigm that focuses on conservation implications for these endemic frogs.

Materials and methods

Study area and species.— We conducted the present study on Murree Hills Frog (*Nanorana vicina*) and Hazara Torrent Frog (*Allopaa hazarensis*) at a natural freshwater stream (33.8432°N, 73.4694°E; 1693 m elevation), located in Village Parhanna, Tehsil Murree, District Rawalpindi, Punjab Province, Pakistan. This stream cascades over rapids and has a few associated ponds (Fig. 1). The stream is part of the Murree-Kotli Sattiyar and Kahuta National Park, Punjab, Pakistan. The topography of the National Park at higher altitude is mainly composed of rugged terrain with narrow valleys. The hilly area contains valleys created by the fast flowing running water of streams and rivers (Atta-ur-Rahman et al. 2010). Most of the vegetation in the area consists of sub-tropical chir pine forest and Himalayan moist temperate (Khan 2006).

Installing radio transmitters.—We captured the frogs using dip nets. We used Holohil BD-2A transmitters (0.49 g) and followed the attachment method by Muths (2003). We arranged the transmitter so that the battery was orientated towards the rear of the individual to allow it to move easily in the water. We made adjustments to the assembly system to make it easier for frogs to carry the transmitter. We constructed the radio transmitter belts by using a very thin, soft elastic thread and light-weight, brightly-coloured plastic beads. For each assembly, we placed a transmitter in the center of the elastic thread, with an equal number of beads on both sides of the transmitter and tied a knot in

the elastic thread. We ensured that the belt was not so tight that it constricted the frog's body and not so loose that it could easily slip off. To attach the transmitter, we stretched the legs of the frog and adjusted the belt as needed to fit around the frog's waist (Muths 2003) (Fig. 2). We used a vernier caliper (Insize Precision Measurement Vernier Caliper SL-1112) to measure snout-vent length (mm) and a digital weighing balance (BL 60001-5) to weigh individuals (grams). We ensured that the attachment assembly would not exceed 10% of the frog's total body mass (Richards et al. 1994).

Radio tracking.—We installed radio transmitters on three Murree Hills Frogs (1 ♂, 2 ♀) and five Hazara Frogs (2 ♂, 3 ♀) in September 2017 and 10 Murree Hills Frogs (5 ♂, 5 ♀) and eight Hazara Frogs (6 ♂, 2 ♀) in September 2018 (non-breeding season) for eight consecutive days during each session. Details on specimens (sex, snout-vent length and weight) and transmitters are given in Table 1. Since the two studied frog species are nocturnal, we located and observed the tracked frogs three times, every three hours from sunset to sunrise. We recorded the distance moved by each frog and calculated mean distances (m) moved for males and females of each species for the entire session.

After testing normality of our data ($P > 0.05$ for Shapiro-Wilk test) in SPSS 25, we used the Mann-Whitney test to compare distances (median) moved by males and females of each species in a given year and distances moved by radio-tracked frogs (pooled data for males and females) of each species between 2017 and 2018 ($\alpha = 0.05$).

Results

We did not observe much movement (limited to < 3 m) along the stream or away from the stream into the forest by either species. The mean distance (m) moved by radio-tracked males and females of Hazara Frogs and Murree Hills Frogs in 2017 and 2018 is given (Table 1). We lost a few transmitters (*Allopaa hazarensis*, ♂ = 4, ♀ = 1; *Nanorana vicina*, ♂ = 2, ♀ = 3) during the study period and, consequently, distance data were not recorded in these cases (Table 1). We found a significant difference between the distance (median) travelled by Murree Hills Frogs (pooled data for the two sexes) ($U = 27$; $P = 0.046$) in 2017 and 2018 with more distance travelled in 2017 ($n = 6$; 0.60 m) as compared to 2018 ($n = 19$; 0.30 m). We did not find significant differences between any of the radio-tracked males and females of Hazara Torrent Frogs in 2017 ($U = 0.00$; $P = 0.58$), 2018 ($U = 7.50$; $P = 0.32$) and of Murree Hills Frogs in 2017 ($U = 1.00$; $P = 0.10$) and 2018 ($U = 34.50$; $P = 0.43$) and Hazara Torrent Frogs (pooled data for the two sexes) ($U = 48$; $P = 0.31$) between 2017 and 2018.

Discussion

We provided data on the movement pattern of two frogs endemic to Himalayan region, Hazara Torrent Frog (*Allopaa hazarensis*) and Murree Hills Frog (*Nanorana vicina*), for the first time. Our data showed that these frog species exhibit limited movement during the observed time period. A synthesis of review on movement and dispersal in amphibians by

Rais and Ahmed (2021) suggests that movement and dispersal of short distance (< 1 km) are common, of medium distance (2-4 km) are uncommon and of long distances (> 5 km) are very rare. Funk et al. (2005) reported that only 4% of marked Columbia Spotted Frog (*Rana luteiventris*) adults moved distances greater than 200 m. Berven and Grudzien (1990) reported only two Wood Frogs (*Rana sylvatica*) moved distances of 2,530 m. De Villiers and Measey (2017) reported that only 5% of marked African Clawed Frogs (*Xenopus laevis*) made over-land movements with distances of ~ 150 m and only 91 individuals moved distances of 2.4 km. The limited extent of movement exhibited by the Murree Hills Frog and the Hazara Torrent Frog during the present study indicates that these endemic frogs depend on a specific stream to live and reproduce, which have critical implications for their conservation.

The Murree Hills Frogs and Hazara Torrent Frogs are facing anthropogenic threats, such as habitat degradation, urbanisation and natural threats, as well as climate change (Ohler and Dutta 2004, Khan et al. 2008). Accordingly, we propose a movement paradigm for these two frog species. In response to such anthropogenic threats and/or climate change, these frog species might face local extinction if they cannot move greater distances or move over-land through open forest to colonise nearby streams (which are, in most cases, > 300 m away from the studied area; Muhammad Rais, pers. obs). These species may have two options for the dispersal and establishment of a metapopulation:

1. move upstream, which would require use of energy reserves and may subject newly dispersed/immigrants to competition with individuals already inhabiting upstream areas,
2. move downstream into unsuitable habitat in lower elevations, with more urban settlements, pollution and deforestation.

The species are not expected to take the risk of dispersal into subtropical scrub streams located further south due to unfavourable habitat and unsuitable environmental conditions. Increase in the air and water temperature or water withdrawal from the streams by the local community could seriously impact populations of these species. The species might be forced to perform over-land migration through the forest to occupy nearby streams, which are situated at a distance difficult to travel by amphibians or perform upstream migration that would require considerable energy reserves and may cause stress in the individuals (Fig. 3). Various mitigation approaches, such as construction of artificial wetlands, enhanced habitat connectivity and wetland restoration have been proposed to mitigate such effects elsewhere in the world (Brand and Snodgrass 2010, Lehtinen and Galatowitsch 2001, Rais and Ahmed 2021). These could be tested in the study area in future for the conservation of the two studied species.

We could have provided a more detailed data of movement in these frogs if we had not lost 38% of our fitted transmitters. The terrain is hilly and the stream had large boulders which were difficult to move. It was unclear whether the transmitters were lost while the frogs hid beneath the boulders, due to predation or to heavy rains which created flooding in the stream. For future research, we suggest using conventional mark-recapture study techniques or using PIT tags. It will reduce cost and more frogs could be included in the

study. Likewise, we also suggest carrying out additional studies by incorporating multiple adjacent stream systems to better understand dispersal and colonisation by these frogs.

Acknowledgements

We thank Paul Freed for proof reading our manuscript. We are also thankful to Punjab Forest Department for facilitating field visits.

Funding program

The study was funded by the International Foundation of Science, Sweden (Grant No. D5459/1) and the Higher Education Commission, Pakistan (NRPU Project 3170).

Ethics and security

We did not carry out any invasive methods for species identification. All specimens were handled with the utmost care and were released in their habitat after installing radio transmitters.

Author contributions

Ayesha Akram: carried out the research as part of her Ph.D study, conducted field surveys, collected data and wrote the original manuscript; Muhammad Rais: designed the study, assisted in site selection and standardisation of the methods and submitted the manuscript; Muhammad Saeed: conducted field surveys and collected data; Waseem Ahmed: conducted field surveys, collected data, conducted literature search and formatted the manuscript; Sumbul Gill: conducted field surveys, collected data, and archived field data; Jibran Haider: conducted field surveys and collected data.

Conflicts of interest

The authors have declared that no competing interests exist.

References

- Aarts G, MacKenzie M, McConnell B, Fedak M, Matthiopoulos J (2008) Estimating space-use and habitat preference from wildlife telemetry data. *Ecography* 31 (1): 140-160. <https://doi.org/10.1111/j.2007.0906-7590.05236.x>
- Adams M, Bury RB (2002) The endemic headwater stream amphibians of the American Northwest: associations with environmental gradients in a large forested preserve.

- Global Ecology and Biogeography 11 (2): 169-178. <https://doi.org/10.1046/j.1466-822x.2002.00272.x>
- Ahmed W, Rais M, Saeed M, Akram A, Khan IA, Gill S (2020) Site occupancy of two endemic stream frogs in different forest types in Pakistan. 15(3):506–511. Herpetological Conservation and Biology 15 (3): 506-511.
 - Atta-ur-Rahman, Khan AN, Collins A, Qazi F (2010) Causes and extent of environmental impacts of landslide hazard in the Himalayan region: a case study of Murree, Pakistan. Natural Hazards 57 (2): 413-434. <https://doi.org/10.1007/s11069-010-9621-7>
 - Berven K, Grudzien T (1990) Dispersal in the Wood Frog (*Rana sylvatica*): Implications for Genetic Population Structure. Evolution 44 (8). <https://doi.org/10.2307/2409614>
 - Brand A, Snodgrass J (2010) Value of artificial habitats for amphibian reproduction in altered landscapes. Conservation Biology 24 (1): 295-301. <https://doi.org/10.1111/j.1523-1739.2009.01301.x>
 - Brunjes K, Ballard WB, Humphrey M, Harwell F, McIntyre N, Krausman P, Wallace M (2006) Habitat Use by Sympatric Mule and White-Tailed Deer in Texas. Journal of Wildlife Management 70 (5): 1351-1359. [https://doi.org/10.2193/0022-541x\(2006\)70\[1351:hubsma\]2.0.co;2](https://doi.org/10.2193/0022-541x(2006)70[1351:hubsma]2.0.co;2)
 - De Villiers FA, Measey J (2017) Overland movement in African clawed frogs (*Xenopus laevis*): empirical dispersal data from within their native range. PeerJ 5 <https://doi.org/10.7717/peerj.4039>
 - Dupuis LA, Bunnell FL, Friele PA (2000) Determinants of tailed frog's range in British Columbia, Canada. Northwest Science 74:109–115. 74: 109-115.
 - Funk WC, Greene AE, Corn PS, Allendorf FW (2005) High dispersal in a frog species suggests that it is vulnerable to habitat fragmentation. Biology Letters 1 (1): 13-16. <https://doi.org/10.1098/rsbl.2004.0270>
 - Khan, et al. (2006) Amphibians and Reptiles of Pakistan. Krieger Publishing Company, Malabar, Florida, USA.
 - Khan MS, Dutta S, Ohler A (2008) *Allopaia hazarensis*: The IUCN Red List of Threatened Species 2008. International Union for Conservation of Nature. IUCN Red List of Threatened Species <https://doi.org/10.2305/iucn.uk.2008.rlts.t58426a11779666.en>
 - Lecis R, Norris K (2004) Habitat correlates of distribution and local population decline of the endemic Sardinian newt *Euproctus platycephalus*. Biological Conservation 115 (2): 303-317. [https://doi.org/10.1016/s0006-3207\(03\)00149-6](https://doi.org/10.1016/s0006-3207(03)00149-6)
 - Lehtinen R, Galatowitsch S (2001) Colonization of restored wetlands by amphibians in Minnesota. The American Midland Naturalist 145 (2): 388-396. [https://doi.org/10.1674/0003-0031\(2001\)145\[0388:CORWBA\]2.0.CO;2](https://doi.org/10.1674/0003-0031(2001)145[0388:CORWBA]2.0.CO;2)
 - Manly BF, McDonald LL, Thomas DL, McDonald TL, Erickson WP (2002) Resource Selection by Animals: Statistical Design and Analysis for Field Studies. 2nd Edition. Kluwer, Dordrecht, The Netherlands..
 - Marsh D, Trenham P (2001) Metapopulation Dynamics and Amphibian Conservation. Conservation Biology 15 (1): 40-49. <https://doi.org/10.1111/j.1523-1739.2001.00129.x>
 - Moilanen A, Nieminen M (2002) Simple connectivity measures in spatial ecology. Ecology 83 (4): 1131-1145. [https://doi.org/10.1890/0012-9658\(2002\)083\[1131:scmise\]2.0.co;2](https://doi.org/10.1890/0012-9658(2002)083[1131:scmise]2.0.co;2)

- Moore RD, Griffiths RA, Román A (2004) Distribution of the Mallorcan midwife toad (*Alytes muletensis*) in relation to landscape topography and introduced predators. *Biological Conservation* 116 (3): 327-332. [https://doi.org/10.1016/S0006-3207\(03\)00202-7](https://doi.org/10.1016/S0006-3207(03)00202-7)
- Muths E (2003) Home Range and Movements of Boreal Toads in Undisturbed Habitat. *Copeia* 2003 (1): 160-165. [https://doi.org/10.1643/0045-8511\(2003\)003\[0160:hramob\]2.0.co;2](https://doi.org/10.1643/0045-8511(2003)003[0160:hramob]2.0.co;2)
- Ohler A, Dutta S (2004) *Nanorana vicina*: IUCN Red List of Threatened Species. International Union for Conservation of Nature. IUCN Red List of Threatened Species <https://doi.org/10.2305/iucn.uk.2004.rlts.t58443a11781868.en>
- Rais M, Ahmed W (2021) Amphibian Dispersal Among Terrestrial Habitats and Wetlands in a Landscape. In: Leal Filho W., Azul A.M., Brandli L., Lange Salvia A., Wall T. (Eds) *Life on Land. Encyclopedia of the UN Sustainable Development Goals*. Springer, Cham. https://doi.org/10.1007/978-3-319-71065-5_151-1.
- Rais M, Ahmed W, Sajjad A, Akram A, Saeed M, Hamid HN, Abid A (2021) Amphibian fauna of Pakistan with notes on future prospects of research and conservation. *ZooKeys* 1062: 157-175. <https://doi.org/10.3897/zookeys.1062.66913>
- Richards SJU., Sinsch U, Alford RA (1994) Radio tracking. Pp 155-158 *In* *Measuring and Monitoring Biological Diversity. Standard Methods for Amphibians*. Heyer W.R., M.A. Donnelly, R.W. Mc- Diarmid, L.C. Hayek, and M.S. Foster (Eds.). Washington and London, Smithsonian Institution Press..
- Saeed M, Rais M, Gray R, Ahmed W, Akram A, Gill S, Fareed G (2021) Rise in temperature causes decreased fitness and higher extinction risks in endemic frogs at high altitude forested wetlands in northern Pakistan. *Journal of Thermal Biology* 95 <https://doi.org/10.1016/j.jtherbio.2020.102809>
- Semlitsch R, Bodie JR (1998) Are Small, Isolated Wetlands Expendable? *Conservation Biology* 12 (5): 1129-1133. <https://doi.org/10.1046/j.1523-1739.1998.98166.x>
- Taylor P, Fahrig L, With K (2006) Landscape connectivity: A return to basics. Pp. 29–43 *In* *Connectivity Conservation*. Crooks, K.R., and M. Sanjayan (Eds.). Cambridge University Press, Cambridge, UK.29-43.



Figure 1.

Freshwater stream at the Village Parhanna, Tehsil Murree, District Rawalpindi, Punjab Province, Pakistan. Photographed by Muhammad Saeed.



Figure 2.

Making the adjustable belt (A-B), measuring the snout-vent length (C), attaching the BD-02 radio transmitter assembly on the frog's waist (D), Hazara Torrent Frog (*Allopaa hazarensis*), (E) Murree Hills Frog (*Nanorana vicina*) (F).




Figure 3.

Movement paradigm for endemic frogs: Hazarda Torrent Frog (*Allopaia hazarensis*) and Murree Hills Frog (*Nanorana vicina*) in subtropical pine forest, based on our radio tracking data.

Table 1.

Mean distance (meter) moved by Hazara Torrent Frog (*Allopaa hazarensis*) and Murree Hills Frog (*Nanorana vicina*) along the studied stream in Village Parhanna, Tehsil Murree, District Rawalpindi, Punjab Province, Pakistan, during eight days in September 2017 and 2018.

*missing data due to the loss of the transmitter

2017					
Hazara Torrent Frog (<i>Allopaa hazarensis</i>)					
Sex	ID	Transmitter Frequency (Hz)	Snout-vent Length (mm)	Weight (gm)	Distance (m)
♂	99	150.712	58	152.82	1.5
♂	97	150.550	57	151.9	*
				Mean (♂)	1.5
♀	100	150.755	59	159.04	1.51
♀	98	150.670	64	158.25	1.2
♀	102	150.867	75	176.4	2.41
				Mean (♀)	1.7± 0.31
Murree Hills Frog (<i>Nanorana vicina</i>)					
♂	101	150.831	81	172.5	1.5
				Mean (♂)	1.5± 0.17
♀	103	150.904	87	207.96	*
♀	104	150.948	98	256.3	2.72
				Mean (♀)	2.72± 0.30
2018					
Hazara Torrent Frog (<i>Allopaa hazarensis</i>)					
♂	368	150.593	56	158.5	0.6
♂	99	150.712	53	149.56	0.6
♂	101	150.832	50	153.9	1.5
♂	373	150.895	49	150.34	*
♂	365	150.396	36	140.2	*
♂	367	150.575	37	148	*
				Mean (♂)	0.9± 0.14
♀	370	150.695	64	161.33	*
♀	366	150.533	60	160.4	0.9
				Mean (♀)	0.9± 0
Murree Hills Frog (<i>Nanorana vicina</i>)					
♂	369	150.614	58	164.18	2.11
♂	374	150.975	79	189.08	1.5
♂	371	150.795	75	187.91	*

♂	100	150.755	89	222.35	*
♂	102	150.867	81	213.41	1.2
				Mean (♂)	1.6± 0.27
♀	104	150.949	73	173.05	*
♀	363	150.352	61	179.31	0.6
♀	372	150.813	72	187.85	0.6
♀	98	150.670	88	212.41	2.41
♀	362	150.313	74	173.05	*
				Mean (♀)	1.2± 0.33