

Diversity of culturable thermophilic bacteria from Tata Pani hot spring of Kotli Azad Jammu and Kashmir

Kazima Ishaq[‡], Asad Hussain Shah[§], Anila Fariq[‡], Sajida Rasheed[‡], Sammyia Jannat[‡]

[‡] Department of Biotechnology, Faculty of Basic and Applied Sciences, University of Kotli Azad Jammu and Kashmir, Kotli, Pakistan

[§] Senior Research Fellow, Faculty of Biology Medicine and Health The University of Manchester The Michael Smith Building Oxford Road Manchester M13 9PT, Manchester, United Kingdom

Corresponding author: Asad Hussain Shah (asad.shah@manchester.ac.uk)

Academic editor: Anna Sandionigi

Abstract

Hot water springs are unique areas populated by mesophiles, thermotolerant and hyperthermophiles. They are the source of diversity of thermophiles, mainly belonging to archaea and bacteria domains. The diversity of thermophiles gives an outline of the huge biological potential that can be exploited for industrial applications. To this end, this study was aimed to isolate and characterise the unexplored thermophilic microorganisms from hot water spring in Tatapani, Tehsil & District Kotli AJK, Pakistan. Around 10 bacterial isolates were identified using morphological, biochemical, physiological and molecular attributes. Sequencing of the 16S rDNA gene of the isolates followed by BLAST search revealed that the strain MBT008 has 100% similarity with *Anoxybacillus kamchatkensis*. MBT012 showed 99.57% similarity with *A. mongoliensis*, MBT014 was affiliated with *A. tengchongensis* with 99.43% similarity, MBT009 showed 99.83% homology with *A. gonensis* and MBT018, 98.70% similarity with *A. karvacharensis*. The presence of all this microbial diversity in one common source is of immense importance related to environmental and industrial aspects in general and extraction of thermostable enzymes from these thermophiles specifically opens new horizons in the field of industrial biotechnology. These thermophiles are revealing new capabilities and are being manipulated by biotechnologists in utilizing them in different unique ways.

Keywords

Hot water springs, thermophiles, diversity, bacteria, archaea, 16S rDNA

Materials and methods

Sample collection

A total of ten water/sediment samples from 10 different locations separated by a distance of approximately 1.0 ft (0.3 m) were collected from different locations of the hot water spring of Tatapani (District Kotli) in sterilised screw-capped bottles and were carried to the laboratory of Biotechnology and processed immediately. The physiochemical characteristics of the hot water spring was examined. The electrical conductivity was monitored by using an EC meter and pH of water samples was measured using a pH meter.

Isolation of thermophilic strains

The sediment and water samples were incubated at 70°C overnight in nutrient broth medium for the isolation of thermophiles. The growth was observed by analysing the turbidity of the growth medium. Next day, the suspended cultures were poured by spread plate technique on the agar plates for the isolation of the mixed population of microorganisms. For calculating the colonyforming unit of viable bacterial cells, several serial dilutions were made in 1:10 ratios. From these dilutions, 0.1 ml was plated on to nutrient agar and, after 24 hours incubation, colonies were observed and counted on the plate. The number of bacteria (CFU) per millilitre of sample was calculated by dividing the number of colonies by the dilution factor. To purify bacterial colonies, single colony streaking is done by picking individual colonies with a sterilised inoculation loop and touching it to the nutrient agar plate.

Morphological characterisation

Colony morphology like colour, shape, elevation and texture of pure bacterial cultures obtained through single colony streaking was observed visually as well as examined under the microscope. Gram's staining was performed to differentiate bacteria according to cell wall composition through a series of staining and decolourising steps.

Growth conditions optimisation

Growth conditions were optimised for pH, temperature, inoculum size and incubation time. Temperature range for incubation varied from 40, 50, 60, 70 and 80°C and the pH dependence of growth was tested in the pH range of 6.0-9.0 in nutrient broth medium, while the effect of varied incubation time intervals for these thermophiles' growth was noted for 24 hours, 48 hours and 72 hours. The growth was also monitored at different inoculum size i.e. 25 µl, 50 µl, 75 µl and 100 µl. The optical density in all conditions was measured to measure the growth of strains at 600 nm on a double beam UV/VIS scanning spectrophotometer (Model: AE-S90-MD).

Biochemical characterisation

For biochemical characterisation, API 20E strips were used to examine the isolates for numerous substrates utilisation like adonitol, glycerol, D-arabinose, erythritol, L-arabinose, D-xylose, L-xylose, ribose, B-methyl xyloside, mannose, glucose, L-sorbose, dulcitol, rhamnose, inositol, mannitol, sorbitol, D-methyl-D-glucoside, amygdalin, N-acetylglucosamine, aesculin, arbutin, salicin, maltose, cellobiose, lactose, trehalose, sucrose, melibiose, raffinose, starch, glycogen, D-turanose, D-fucose, inulin, L-fucose, D-arabitol, D-lyxose, L-arabitol, gluconate, 2,5-ketogluconate and xylitol. Suspended bacterial cultures (100 µl) grown for 24 hours in 0.8% sterilised saline solution were injected into the wells of strips containing the substrate and incubated at 70°C for 24 hours. The colour change was studied according to kit instructions. After the incubation period, all the results were recorded and then revealed the tests which require the addition of reagents. The presence of oxidase and catalase enzymes was examined according to the procedures designed by Prescott et al. (1997).

Antibiotic Sensitivity

The sensitivity profiles of the bacteria isolated from water/sediment samples against various antibiotics like ciprofloxacin, levofloxacin, azithromycin, cefixime, linezolid etc. was also done by the disc diffusion method.

DNA extraction and amplification of the 16S rDNA gene

Genomic DNA was extracted from bacteria consistent with the protocol described by Cheng and Jiang (2006). The 16S rDNA gene amplification from the purified genomic DNA was performed in 0.2 ml PCR tubes with 20 µl reaction volume using the forward primer 27F AGAGTTTGATCCTGGCTCAG and the reverse primers 1492R TACGGCTACCTTGTTACGACTT and all the PCR amplifications were carried out by using a thermal cycler (Model: TC-TE).

Sequencing analysis

After amplification, the PCR products obtained were sequenced by using same upstream and downstream primers, by a commercial sequencing facility i.e. MacroGen. The entire 16S rDNA sequences of these bacterial strains obtained were blasted using the online NCBI BLAST programme (Waterhouse et al. 2009). Phylogenetic analyses were done to express evolutionary associations of these isolates. The analysis started with aligning of sequences by the Clustal W tool and after alignment, the phylogenetic tree was built using MEGAX software.

Introduction

Hot springs comprise of the environment with high temperature ranges and formed due to geothermal movement and are host to variety of thermophiles (Zhang et al. 2004, Yohandini et al. 2015). A hydrothermal spring is characterised by warm groundwater which is heated by intense geothermal heat, the indispensable heat beneath the Earth's surface. The life in hot springs started long before reaching the surface and the hot water

serves as a habitat for large number of bacteria and algae which were then termed as thermophiles (Adiguzel et al. 2009). These thermophilic microorganisms prefer to grow at high temperature which normally does not exist in nature. Thermophilic microorganisms were the first living organisms which develop and evolve in the ancient days when the Earth's surface temperature was relatively high, so they were called the "Universal Ancestor" (Kumar 2010). Moreover, a study of thermophiles resulted in the discovery of novel species (Kozina et al. 2010).

Thermophiles are characterised into moderate (optimum temperature, 55–60°C), extremophiles (optimum temperature, 60–80°C) and hyperthermophiles (optimum temperature, 80–110°C) (Saha et al. 2005). Nowadays, extremophilic microorganisms with their different classes, thermophiles, alkaliphiles, acidophiles, halophiles and psychrophiles, attracted the attention of many researchers in various areas, because of their capability to resist and perform life functions under extreme environmental conditions (Tango and Islam 2010). Due to their potential to produce thermostable enzymes (like DNA polymerases, chitinases, cellulases, amylases, pectinases, proteases, lipases and xylanases), thermophilic bacteria have attained great consideration amongst extremophiles. These thermostable enzymes are more stable and possess unique characteristics that make them appropriate for performing biotechnological procedures at higher temperatures (Singh et al. 2011). The earliest thermophilic spores forming bacteria capable of surviving at 70°C were discovered by Miquel in 1888 for the first time (Miquel 1888). Since that, numerous spore forming strains of thermophilic bacteria, mainly related to *Bacillus* and *Clostridium* genera, have been characterised (Maugeri et al. 2001, Belduz et al. 2003).

Thermophiles have attracted the attention of many scientists for their potential in biotechnological processes (Adiguzel et al. 2009). These thermophilic bacteria have been characterised phenotypically and genotypically in particular from many hot springs in different regions of the world, including Turkey (Gul-Guven et al. 2008), Bulgaria (Derekova et al. 2008), Italy (Maugeri et al. 2001), China (Poli et al. 2009), Greece (Sievvert et al. 2000, Lau et al. 2008), Iceland (Takacs et al. 2001) and India (Sharma et al. 2008). Due to development in techniques of molecular biology, such as gene sequencing of 16S rRNA, tremendous opportunities have become available for identifying and characterising microbes at species level.

In recent times, the bio-chemical and molecular characterisation of thermophilic microorganisms from geothermal springs has been reported to explain the mechanism and molecular cause of the adaptation of these organisms to extreme temperature (Panda et al. 2016).

Tatapani is situated in Kotli, Azad Kashmir (Pakistan) at a latitude and longitude of 33.60 N, 73.94 E, respectively and TattaPani hot springs are located on the right bank of Poonch River as shown in Fig. 1 and can be reached from Kotli by covering a distance of 26 km. The hot spring of Tatapani has been known for its flowing hot water on the western bank of Poonch River at an elevation of 682 m (2,237 ft). Due to its exclusive hot water, it attracts the attention of tourists who benefit from its steam in winter. Locals and tourists

enjoy this hot water bath which is rich in sulphur. The water is believed to have high medical value and good for the skin. However, the microbial diversity of this very unique repository of microorganisms has not been explored. The tremendous capability of such important bacterial strains can be a substantial source of thermostable enzymes.

Tatapani hot spring has not yet been explored from the microbiological characteristics. The specific purpose of this study was to identify and characterise the bacterial strains from the hot water spring of Tatapani by using various morphological markers and to carry out the biochemical and molecular characterisation of the isolated bacteria. Due to limited scientific knowledge existing on the research being done in this field and to explore this unseen and under-utilised source of thermostable bacteria, this study was performed to isolate and characterise the thermophilic microorganisms from various locations of the hotwater spring situated in Tatapani, District Kotli AJK, Pakistan. The advancement of the scientific information regarding the biodiversity of these delicate ecosystems underlines the requirement for their preservation; in addition, this microbial diversity possibly will be the basis for further biotechnological utilizations representing an important keystone for a developing region.

Results

Isolation and Characterisation of the Samples

In this study, the hot spring represented a moderate to high temperature (39.9–75°C) and neutrophilic to alkalophilic (pH 7.03–8.6) environment with varying electrical conductivity (0.51– 3.27 μscm^{-1}). After enrichment, visible turbidity was observed in almost all samples within 24 to 48 hours, frequently appearing clumpy or as a surface pellicle. After the turbidity had become equally thicker, the colour of some of the tubes containing the microbial sample was yellow or orange. The total cell count varied from one sampling site to another (Table 1). The enrichment samples were given codes viz. MBT006, MBT007, MBT008, MBT009, MBT010, MBT011, MBT012, MBT013, MBT014 and MBT018.

The variation of colonies count was observed at different sampling sites. These ten samples were cultured at 70°C. It was noted that the number of CFUs in sample 1 i.e. MBT006 is greater (6.0×10^3) and decreased as we move away from the main hot water source towards sites 2, 3, 4 and so on.

Identification and differentiation of isolates

The bacterial strains exhibited different colony and cell morphology. Under light microscope, all were detected as rods either single or were arranged in chains. Within 24 hours of incubation at 70°C, compact spreading colonies were seen which were yellow, white, creamy, brown and greyish white in colour.

Morphologically, the strains exhibited some difference in the colour, shape, texture and margin of the colonies (Table 2). They appeared creamy, white and grey; translucent or

opaque; smooth or rough; with regular or irregular edges. The colonies appeared either raised or flat on the agar surface.

Biochemical and Metabolic Characterisation of the bacterial Isolates

The biochemical characterisation was done by API 20E strips for identifying phenotypic diversity and none of the ten isolates shared the common phenotypic characters (Table 3). All the isolates were positive for β -galactosidase and citrate utilization. Most of the isolates were unable to metabolise many sugars i.e. nositol, mannitol, sorbitol, rhamnose, melibiose, sucrose and amygdalin. However, a few isolates showed glucose and arabinose utilization (Table 4). Only two isolates i.e. MBT011 and MBT012 exhibited positive results for indole production. All except two (MBT006 and MBT012) were able to produce gelatinase enzyme.

Antibiotic sensitivity

The antibiotic sensitivity profile of the bacterial isolates showed that all isolates were sensitive to amoxicillin (Ax) and levofloxacin (LEV) and resistant towards metronidazole (MET). The strains MBT009 and MBT012 were found resistant to clarithromycin (CLR) and linezolid (LNZ), while most of the strains showed resistance against AZM (Table 5).

Physiological characteristics

Physiological studies were performed with all the ten strains. Temperature (a physiological limiting factor) controls the microbial cells multiplication. The temperature of water governs distribution of microbes within hot springs. The association between growth rate and temperature is shown in Fig. 2. The optimum temperature for their growth was 70°C, the maximum growth temperature was 80°C and the minimum temperature was about 40°C. The maximum growth of bacterial isolates was observed when incubated overnight at 70°C. At 40°C and 80°C, no growth was observed; minimum growth was appeared after incubation at 50°C and 60°C. These results depict that the optimal temperature for growth of these isolates was 70°C.

In addition to temperature, pH of water is an important factor defining the microbial diversity in hot water springs. All isolated strains were inoculated in LB broth with different pH ranges from 6.0-9.0. Good growth was noted at pH 7 and it is considered as the optimum pH for the growth of thermophilic bacteria as shown in Fig. 2.

The growth rate was also measured at varying time intervals in order to obtain the optimum growth period. Best growth was observed after 48 hours of incubation. From these results, the optimum time for growth of thermophiles was considered at 48 hours. Similarly, the growth rate of thermophilic bacteria was observed best when incubated with 50 μ l of inoculum.

Molecular characterisation of the Isolates

For the ultimate identification and phylogenetic study of the strains, 16S rDNA gene sequencing was done following the Basic Local Alignment Search Tool (BLAST)

programme. The 16S rDNA gene sequences of the five isolates were aligned with their associated bacterial sequences from the GenBank databases. Sequence analysis revealed high affiliation with those of the linked strains available in GenBank. The sequence alignment showed that all the strains belonged to genus *Anoxybacillus* with significant sequence similarity as listed in Table 6.

The phylogenetic tree was constructed for all the isolated strains, based upon 16S rDNA gene sequence alignment. BLAST analysis showed the strongest similarity (100%) of the strain MBT008 with *A. kamchatkensis* and MBT009 (99.83%) with *A. gonensis*. MBT012 showed 99.57% similarity with *A. mongoliensis*, MBT014 with *A. tengchongensis* (99.43%) and MBT018 was closely related to *A. karvacharensis* (98.70%) (Fig. 3).

The genus *Anoxybacillus* was abundant in the majority of the explored locations; the presence of *Anoxybacillus* in the majority of sampled sites is ascribed to the capability of the genus to pass at higher rates, as well as their potential to resist extreme environmental stresses.

Discussion

Geothermal environments are enriched by the diversity of thermophilic archaea and bacteria. Many thermophilic and hyperthermophilic archaea have been isolated from geothermal and hydrothermal systems. In order to obtain a better understanding of microbial ecosystems and roles in the geothermic community, many studies have been performed to reveal the link between microbial niches, diversity and physicochemical factors, such as temperature, pH and water chemistry (Skirnisdottir et al. 2000, Purcell et al. 2007, Pearson et al. 2008, Everroad et al. 2012, Wang et al. 2013). The hot spring of Tattapani is a habitat for the diversity of thermophilic microorganisms and it represents the functional adaptations of these microbes to withstand extreme temperatures.

One of the studies (Nakagawa and Fukui 2002) reported that the bacterial communities in streamers with a temperature above 65°C was abundant in *Thermus*, *Thermodesulfobacteria*, *Crenarchaeota* and *Aquificae*. In 1995, Miquel firstly characterised the aerobic thermophilic, spore forming bacteria, which were able to grow at 70°C (Miquel 1995).

Bacteria belonging to the genus *Thermus* had become important in thermophile research after the isolation of *Thermus aquaticus* carried out by Brock and Freeze (1969). Later on, numerous additional *Thermus* species have been discovered. However, in this study, the existence of thermophilic bacteria related to genus *Anoxybacillus* were investigated in thermal spring of Tattapani, Pakistan. All of the isolated thermophiles belong to the domain bacteria, phylum firmicutes, class bacilli, order bacillales, within family bacillaceae (<http://rdp.cme.msu.edu/seqmatch>). Therefore, it is clear that the water sample enriched with NB media was appropriate for growth of the genus *Bacillus*.

The decrease in CFU with change in altitude is attributed to the gradual decrease in temperature of water when the sample was taken some distance away from the hot

spring. The temperature significantly influenced the bacterial richness. Occurrence of bacteria, at elevated temperatures, is due to numerous adaptations in physiological conditions and genetics as the stress response to maintain homeostasis (Wang et al. 2015).

Microorganisms grow across a wide temperature range, pH, salinity and oxygen levels. Extremophilic microorganisms have the potential to grow in diverse extreme environments, such as low or high temperature, alkaline and acidic pH, high radiations and high salinity (Mirete et al. 2016).

The optimal temperature observed for maximum growth of isolates in our study was 70°C, optimal pH observed was 7.0 and optimal time interval was 48 hours. Large-scale comparisons of genomes of mesophiles and thermophiles had confirmed that genomes of the thermophiles contain a higher guanine and cytosine (GC) content than that of mesophiles (Takami et al. 2004, Wang et al. 2015). It has been established that a high GC content is related to the thermal stability of the genome and optimal growth temperature of microbes (Musto et al. 2005, Musto et al. 2006). In addition, rRNAs and tRNAs, that serve as the translational machinery for most of the thermophiles, were reported to contain high GC contents as well (Higgs and Ran 2008, Satapathy et al. 2010).

The results of phenotypic and physiological characters of thermophilic bacterial isolates, MBT010 and MBT008 are in line with those reported by Bisht and Panda (2011), whereas the bacterial strain MBT011 is in agreement with outcomes of the studies by Shida et al. (1997) and for the thermophilic bacterial strains MBT006 and MBT018, our findings also correlate with the study of Mohammed in 2012, who isolated the three new species of moderately thermophilic bacteria from three hot springs, located in Jazan District, southwestern of Saudi Arabia. They did the phylogenetic analysis of these strains using their 16S rDNA sequence and also studied the morphological, biochemical and physiological characteristics of the isolates (Mohammad A. Khiyami 2012). On the basis of phenotypical and physiological features, these isolated thermophiles also depicted consistent results with those of Rastogi et al. (2009).

In another study from Pakistan, Zahra et al in 2020 isolated strain AK9 from the hot water spring of Tattapani Azad Kashmir, Pakistan; extracted and purified cellulase enzyme which reserved its activity from 50-70°C and 3–7 pH. They reported that *B. amyloliquefaciens* AK9 can be used in bioconversion of lignocellulosic biomass to fermentable sugar (Amin et al. 2016a, Amin et al. 2016b, Zahra et al. 2020).

The phylogenetic analysis, based on 16S rRNA gene similarity, depicted that the strain MBT009 was affiliated with the species *Anoxybacillus gonensis* with 99.83% sequence similarity as shown in Fig. 3(A). Our results are consistent with the results of Dai et al. (2011) who had isolated the E13T strain from water-sediment samples from springs in Yunnan Province of China, closely related to the species of *A. flavithermus* (99.2% sequence similarity). These findings also resemble the study in which Chan et al. (2015) identified bacterial strains affiliated with genera *Anoxybacillus* and *Geobacillus* from a Malaysian hot spring. Their results revealed a clear domination of the genus

Anoxybacillus represented by *A. thermarum*, *A. flavithermus* and *A. pushchinoensis* (Chan et al. 2015).

Our study correlates with the study of Balsam et al. (2017) in which they reported the morphological, microscopic, biochemical, molecular and physiological characterisation of ten isolates isolated from hot springs in Jordan; out of which nine isolates belonged to the genus *Bacillus*.

Anoxybacillus a relatively new genus, is alkali tolerant, Gram-positive rod, tested positive for catalase and oxidase activity. In Pakistan, this genus has been identified and characterised by Jabeen et al. in 2019 from the hot water spring in Chakwal (Jabeen et al. 2019).

In our study, one of the isolates namely MBT009 had nearest similarity (99.83%) with *Anoxybacillus gonensis* which was related to the study of Belduz et al. (2003), who had isolated the thermophilic strain G2T (which showed strong homology > 97% with *A. gonensis*) from mud and water samples from the Gonen and Diyadin hot springs, respectively, located in the Turkish provinces of Balikesir and Agri. They had screened the strains for thermostable enzyme production and found that both strains had the ability to produce industrially-important enzymes (Belduz et al. 2003).

Conclusion

Thermophiles are unique organisms with tremendous diversity and biological significance. Azad Jammu Kashmir is an unexplored area with blessed sources of hot springs. The microbial diversity of hot springs from Kotli AJK has been explored for the first time on the basis of morphological, biochemical and molecular markers. In this study, ten (10) strains of thermophilic bacteria were isolated from different locations of geothermal hot springs of Tatapani. The 16s gene sequence revealed striking genetic variability in these isolates. These strains were related to genus *Anoxybacillus* with significant levels of similarity. It was clearly demonstrated that the thermal water of Tatapani (Kotli) hot spring can be an important source of diversity of thermophilic bacteria which is of great significance for biotechnological processes at elevated temperatures. It is important to report that this is the first study to reveal this hot spring from a single common source for a diversity of numerous thermophilic bacteria, so their applications in the field of biotechnology can be exploited for the production of thermostable enzymes, as well as their metabolites can be used in various biotechnological processes. The study provided an authentic base for studies on complete genomic characterisation and are highly recommended for generating more detailed information of novel isolates.

Acknowledgements

The authors acknowledge the financial and logistic support provided by Finance and Transport sections of University of Kotli AJ&K.

Hosting institution

University of Kotli Azad Jammu and Kashmir

Author contributions

Asad Hussain Shah conceived the idea and led the project.

Kazima Ishaq performed the experiments and wrote the manuscript.

Anila Fariq performed the experiments and wrote the manuscript.

Sajida Rasheed analysed the data.

Sammyia Jannat interpreted the data and reviewed the manuscript.

Conflicts of interest

None.

References

- Adiguzel A, Ozkan H, Baris O, Inan K, Gulluce M, Sahin F (2009) Identification and characterization of thermophilic bacteria isolated from hot springs in Turkey. *Journal of Microbiological Methods* 79 (3): 321-328. <https://doi.org/10.1016/j.mimet.2009.09.026>
- Amin A, Ahmed I, Habib N, Abbas S, Xiao M, Hozzein W, Li W (2016a) *Nocardioides pakistanensis* sp. nov., isolated from a hot water spring of Tatta Pani in Pakistan. *Antonie van Leeuwenhoek* 109 (8): 1101-1109. <https://doi.org/10.1007/s10482-016-0711-8>
- Amin A, Ahmed I, Khalid N, Osman G, Khan IU, Xiao M, Li W (2016b) *Streptomyces caldifontis* sp. nov., isolated from a hot water spring of Tatta Pani, Kotli, Pakistan. *Antonie van Leeuwenhoek* 110 (1): 77-86. <https://doi.org/10.1007/s10482-016-0778-2>
- Balsam M, Al Daghistani H, Jaouani A, Abdel-Latif S, Kennes C (2017) Isolation and characterization of thermophilic bacteria from Jordanian hot springs: *Bacillus licheniformis* and *Thermomonas hydrothermalis* isolates as potential producers of thermostable enzymes. *International Journal of Microbiology* 2017: 1-12. <https://doi.org/10.1155/2017/6943952>
- Belduz AO, Dulger S, Demirbag Z (2003) *Anoxybacillus gonensis* sp. nov., a moderately thermophilic, xylose-utilizing, endospore-forming bacterium. *International Journal of Systematic and Evolutionary Microbiology* 53 (Pt 5): 1315-1320. <https://doi.org/10.1099/ijs.0.02473-0>
- Bisht SS, Panda AK (2011) Biochemical characterization and 16S rRNA sequencing of few lipase-producing thermophilic bacteria from Taptapani hot water spring, Orissa, India. *Biotechnology Research International* 2011: 452710. <https://doi.org/10.4061/2011/452710>

- Brock TD, Freeze H (1969) *Thermus aquaticus* gen. n. and sp. n., a nonsporulating extreme thermophile. *Journal of Bacteriology* 98 (1): 289-97. <https://doi.org/10.1128/jb.98.1.289-297.1969>
- Chan CS, Chan K, Tay Y, Chua Y, Goh KM (2015) Diversity of thermophiles in a Malaysian hot spring determined using 16S rRNA and shotgun metagenome sequencing. *Frontiers in Microbiology* 6 <https://doi.org/10.3389/fmicb.2015.00177>
- Cheng H, Jiang N (2006) Extremely rapid extraction of DNA from bacteria and yeasts. *Biotechnology Letters* 28 (1): 55-9. <https://doi.org/10.1007/s10529-005-4688-z>
- Dai J, Liu Y, Lei Y, Gao Y, Han F, Xiao Y, Peng H (2011) A new subspecies of *Anoxybacillus flavithermus* ssp. yunnanensis ssp. nov. with very high ethanol tolerance. *FEMS Microbiology Letters* 320 (1): 72-78. <https://doi.org/10.1111/j.1574-6968.2011.02294.x>
- Derkova A, Mandeva R, Kambourova M (2008) Phylogenetic diversity of thermophilic carbohydrate degrading bacilli from Bulgarian hot springs. *World Journal of Microbiology and Biotechnology* 24 (9): 1697-1702. <https://doi.org/10.1007/s11274-008-9663-0>
- Everroad RC, Otaki H, Matsuura K, Haruta S (2012) Diversification of bacterial community composition along a temperature gradient at a thermal spring. *Microbes and Environments* 27 (4): 374-381. <https://doi.org/10.1264/jisme2.me11350>
- Gul-Guven R, Guven K, Poli A, Nicolaus B (2008) *Anoxybacillus kamchatkensis* subsp. *asaccharedens* subsp. nov., a thermophilic bacterium isolated from a hot spring in Batman. *The Journal of General and Applied Microbiology* 54 (6): 327-34. <https://doi.org/10.2323/jgam.54.327>
- Higgs PG, Ran W (2008) Coevolution of codon usage and tRNA genes leads to alternative stable states of biased codon usage. *Molecular Biology and Evolution* 25 (11): 2279-91. <https://doi.org/10.1093/molbev/msn173>
- Jabeen F, Muneer B, Qazi JI (2019) Characterization of thermophilic bacteria *Anoxybacillus rupiensis* and cultivation in agroindustrial wastes isolated from hot spring in Chakwal, Pakistan. *Pakistan Journal of Zoology* 51 (4). <https://doi.org/10.17582/journal.pjz/2019.51.4.1243.1250>
- Kozina K, Kolganova TV, Chernyh NA, Bonch-Osmolovskaya EA (2010) *Caldanaero bacteruzonensis* sp. nov., an anaerobic, thermophilic, heterotrophic bacterium isolated from a hot spring. *International Journal of Systematic and Evolutionary Microbiology* 60: 1372-5. <https://doi.org/10.1099/ijs.0.012328-0>
- Kumar, et al. (2010) Characterization of hot water spring source isolated clones of bacteria and their industrial applicability. *International Journal of Chemical Research* 2: 01-07. <https://doi.org/10.9735/0975-3699.2.1.1-7>
- Lau MY, Aitchison J, Pointing S (2008) Bacterial community composition in thermophilic microbial mats from five hot springs in Central Tibet. *Extremophiles* 13 (1): 139-149. <https://doi.org/10.1007/s00792-008-0205-3>
- Maugeri T, Gugliandolo C, Caccamo D, Stackebrandt E (2001) A polyphasic taxonomic study of thermophilic bacilli from shallow, Marine Vents. *Systematic and Applied Microbiology* 24 (4): 572-587. <https://doi.org/10.1078/0723-2020-00054>
- Miquel A (1995) De l'« École » au « Collège ». L'apprentissage du savoir vivant 209-214. <https://doi.org/10.3917/puf.vi.1995.01.0209>
- Miquel P (1888) Monographie d'un bacille vivant au-delà de 70 °C. *Ann Micrographic* 1 (3).

- Mirete S, Morgante V, González-Pastor JE (2016) Functional metagenomics of extreme environments. *Current Opinion in Biotechnology* 38: 143-9. <https://doi.org/10.1016/j.copbio.2016.01.017>
- Mohammad A. Khiyami (2012) Thermo-aerobic bacteria from geothermal springs in Saudi Arabia. *African Journal of Biotechnology* 11 (17). <https://doi.org/10.5897/ajb11.3339>
- Musto H, Naya H, Zavala A, Romero H, Alvarez-Valin F, Bernardi G (2005) The correlation between genomic G + C and optimal growth temperature of prokaryotes is robust: A reply to Marashi and Ghalanbor. *Biochemical and Biophysical Research Communications* 330 (2): 357-360. <https://doi.org/10.1016/j.bbrc.2005.02.133>
- Musto H, Naya H, Zavala A, Romero H, Alvarez-Valin F, Bernardi G (2006) Genomic GC level, optimal growth temperature, and genome size in prokaryotes. *Biochemical and Biophysical Research Communications* 347 (1): 1-3. <https://doi.org/10.1016/j.bbrc.2006.06.054>
- Nakagawa T, Fukui M (2002) Phylogenetic characterization of microbial mats and streamers from a Japanese alkaline hot spring with a thermal gradient. *The Journal of General and Applied Microbiology* 48 (4): 211-22. <https://doi.org/10.2323/jgam.48.211>
- Panda AK, Bisht SS, De Mandal S, Kumar NS (2016) Bacterial and archeal community composition in hot springs from Indo-Burma region, North-east India. *AMB Express* 6 (1): 111. <https://doi.org/10.1186/s13568-016-0284-y>
- Pearson A, Pi Y, Zhao W, Li W, Li Y, Inskeep W, Perevalova A, Romanek C, Li S, Zhang C (2008) Factors controlling the distribution of archaeal tetraethers in terrestrial hot springs. *Applied and Environmental Microbiology* 74 (11): 3523-3532. <https://doi.org/10.1128/aem.02450-07>
- Poli A, Romano I, Cordella P, Orlando P, Nicolaus B, Ceschi Berrini C (2009) *Anoxybacillus thermarum* sp. nov., a novel thermophilic bacterium isolated from thermal mud in Euganean hot springs, Abano Terme, Italy. *Extremophiles* 13 (6): 867-874. <https://doi.org/10.1007/s00792-009-0274-y>
- Prescott, John P, Harley, Donald A. Klein, Delihias (1997) *Microbiology. The Quarterly Review of Biology* 72 (4): 472-473. <https://doi.org/10.1086/419992>
- Purcell D, Sompong U, Yim LC, Barraclough T, Peerapornpisal Y, Pointing S (2007) The effects of temperature, pH and sulphide on the community structure of hyperthermophilic streamers in hot springs of Northern Thailand. *FEMS Microbiology Ecology* 60 (3): 456-466. <https://doi.org/10.1111/j.1574-6941.2007.00302.x>
- Rastogi G, Muppidi G, Gurram R, Adhikari A, Bischoff K, Hughes S, Apel W, Bang S, Dixon D, Sani R (2009) Isolation and characterization of cellulose-degrading bacteria from the deep subsurface of the homestake gold mine, Lead, South Dakota, USA. *Journal of Industrial Microbiology & Biotechnology* 36 (4): 585-598. <https://doi.org/10.1007/s10295-009-0528-9>
- Saha P, Mondal AK, Mayilraj S, Krishnamurthi S, Bhattacharya A, Chakrabarti T (2005) *Paenibacillus assamensis* sp. nov., a novel bacterium isolated from a warm spring in Assam, India. *International Journal of Systematic and Evolutionary Microbiology* 55 (Pt 6): 2577-2581. <https://doi.org/10.1099/ijs.0.63846-0>
- Satapathy SS, Dutta M, Ray SK (2010) Higher tRNA diversity in thermophilic bacteria: a possible adaptation to growth at high temperature. *Microbiological Research* 165 (8): 609-16. <https://doi.org/10.1016/j.micres.2009.12.003>
- Sharma A, Pandey A, Shouche YS, Kumar B, Kulkarni G (2008) Characterization and identification of *Geobacillus* spp. isolated from Soldhar hot spring site of Garhwal

- Himalaya, India. *Journal of Basic Microbiology* 49 (2): 187-194. <https://doi.org/10.1002/jobm.200800194>
- Shida O, Takagi H, Kadowaki K, Nakamura LK, Komagata K (1997) Transfer of *Bacillus alginolyticus*, *Bacillus chondroitinus*, *Bacillus curdlanolyticus*, *Bacillus glucanolyticus*, *Bacillus kobensis*, and *Bacillus thiaminolyticus* to the genus *Paenibacillus* and emended description of the Genus *Paenibacillus*. *International Journal of Systematic Bacteriology* 47 (2): 289-98. <https://doi.org/10.1099/00207713-47-2-289>
 - Sievert S, Ziebis W, Kuever J, Sahm K (2000) Relative abundance of archaea and bacteria along a thermal gradient of a shallow-water hydrothermal vent quantified by rRNA slot-blot hybridization. *Microbiology* 146 (6): 1287-1293. <https://doi.org/10.1099/00221287-146-6-1287>
 - Singh G, Bhalla A, Kaur P, Capalash N, Sharma P (2011) Laccase from prokaryotes: A new source for an old enzyme. *Reviews in Environmental Science and Biotechnology* 10 (4): 309-326. <https://doi.org/10.1007/s11157-011-9257-4>
 - Skirnisdottir S, Hreggvidsson GO, Hjörleifsdottir S, Marteinsonn VT, Petursdottir SK, Holst O, Kristjánsson JK (2000) Influence of sulfide and temperature on species composition and community structure of hot spring microbial mats. *Applied and Environmental Microbiology* 66 (7): 2835-41. <https://doi.org/10.1128/AEM.66.7.2835-2841.2000>
 - Takacs CD, Ehringer M, Favre R, Cermola M, Eggertsson G, Palsdottir A, Reysenbach A (2001) Phylogenetic characterization of the blue filamentous bacterial community from an Icelandic geothermal spring. *FEMS Microbiology Ecology* 35 (2): 123-128. <https://doi.org/10.1111/j.1574-6941.2001.tb00795.x>
 - Takami H, Takaki Y, Chee G, Nishi S, Shimamura S, Suzuki H, Matsui S, Uchiyama I (2004) Thermodaptation trait revealed by the genome sequence of thermophilic *Geobacillus kaustophilus*. *Nucleic Acids Research* 32 (21): 6292-303. <https://doi.org/10.1093/nar/gkh970>
 - Tango MSA, Islam MR (2010) Potential of extremophiles for biotechnological and petroleum applications. *Energy Sources* 24 (6): 543-559. <https://doi.org/10.1080/00908310290086554>
 - Wang Q, Cen Z, Zhao J (2015) The survival mechanisms of thermophiles at high temperatures: an angle of omics. *Physiology (Bethesda, Md.)* 30 (2): 97-106. <https://doi.org/10.1152/physiol.00066.2013>
 - Wang S, Hou W, Dong H, Jiang H, Huang L, Wu G, Zhang C, Song Z, Zhang Y, Ren H, Zhang J, Zhang L (2013) Control of temperature on microbial community structure in hot springs of the Tibetan Plateau. *PLOS One* 8 (5): e62901. <https://doi.org/10.1371/journal.pone.0062901>
 - Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ (2009) Jalview Version 2 A multiple sequence alignment editor and analysis workbench. *Bioinformatics (Oxford, England)* 25 (9): 1189-91. <https://doi.org/10.1093/bioinformatics/btp033>
 - Yohandini H, Julinar, Muharni (2015) Isolation and phylogenetic analysis of thermophile community within Tanjung Sakti hot spring, South Sumatera, Indonesia. *Hayati Journal of Biosciences* 22 (3): 143-148. <https://doi.org/10.1016/j.hjb.2015.10.006>
 - Zahra T, Irfan M, Nadeem M, Ghazanfar M, Ahmad Q, Ali S, Siddique F, Yasmeen Z, Franco M (2020) Cellulase production by *Trichoderma viride* in submerged fermentation using response surface methodology. *Punjab University Journal of Zoology* 35 (2). <https://doi.org/10.17582/journal.pujz/2020.35.2.223.228>

- Zhang H, Kallimanis A, Koukkou AI, Drinas C (2004) Isolation and characterization of novel bacteria degrading polycyclic aromatic hydrocarbons from polluted Greek soils. *Applied Microbiology and Biotechnology* 65 (1): 124-131.




Figure 1.
Map of the hot spring Tatapani, AJK, Pakistan.




Figure 2. Measuring growth rate of bacteria. **(A)** Growth rate of isolates at different temperatures. **(B)** Growth rate of isolates at different pH; **(C)** Growth rate of isolates at different incubation periods; **(D)** Growth rate of isolates at different inoculum size.




Figure 3.

Phylogenetic trees of Tatapani hot spring isolates which have the highest homology with the genus *Anoxybacillus*. The phylogenetic trees were constructed using the neighbour-joining method, based on total 16S rDNA sequencing with 100 bootstrap replicates (software MEGAX). **(A)** MBT009 showing similarity with *A. gonensis*; **(B)** MBT012 showing similarity with *A. mongoliensis*; **(C)** MBT014 with *A. tengchongensis*; **(D)** MBT008 with *A. karvacharensis*; **(E)** MBT018 was closely related to *A. karvacharensis*.

Table 1.

Total cell count of live and cultivable cells in samples.

Sample	Temperature	pH	CFU/ml
MBT006	67	7	6.0×10^3
MBT007	52	6.9	5.2×10^3
MBT008	49	7	4.7×10^3
MBT009	52	7.1	5.1×10^3
MBT010	51	6.9	Not Determined
MBT011	53	7	2.9×10^3
MBT012	54	6	2.1×10^3
MBT013	52	6	1.4×10^4
MBT014	51	7	5.7×10^3
MBT018	49	7	5.2×10^3

Table 2.

Morphological characteristics.

Sample	Shape	Margin	Visual colour	Colour under microscope	Elevation	Texture
MBT006	Regular	Smooth	White-creamy	Dark brown	Raised	Creamy
MBT007	Irregular	Smooth	Grey-yellow	Dark brown	Raised	Creamy
MBT008	Regular	Smooth	white	Golden	Flat	Creamy
MBT009	Irregular	Rough	White-creamy	Dark brown	Raised	Sticky
MBT010	Irregular	Smooth	Grey-brown	Dark brown	Raised	Creamy
MBT011	Regular	Rough	White	Brown	Raised	Creamy
MBT012	Irregular	Smooth	Off-white	Dark brown	Flat	Creamy
MBT013	Regular	Smooth	Yellow	Pale	Raised	Sticky
MBT014	Regular	Rough	Pale	Brown	Raised	Creamy
MBT018	Irregular	Smooth	Off-white	Golden	Raised	Creamy

Table 3.

Cell morphology of bacterial isolates.

Strains	Cell shape	Gram staining	Arrangement
MBT006	Rod	+ve	Pairs/clusters
MBT007	Rod	+ve	Pairs
MBT008	Rod	+ve	Pairs
MBT009	Rod	+ve	Clusters
MBT010	Rod	+ve	Pairs/clusters
MBT011	Rod	+ve	Pairs/clusters
MBT012	Rod	+ve	Pairs/clusters
MBT013	Rod	+ve	Pairs
MBT014	Rod	+ve	Pairs/clusters
MBT018	Rod	+ve	Clusters

Table 4.

Biochemical characteristics of bacterial isolates.

Tests	MBT006	MBT007	MBT008	MBT009	MBT010	MBT011	MBT012	MBT013	MBT014	MBT018
β-galactosidase	+	+	+	+	+	+	+	+	+	+
L-arginine	-	+	-	-	+	+	-	+	+	+
L-lysine	-	-	-	+	+	-	-	-	-	-
Citrate utilisation	+	+	+	+	+	+	+	+	+	+
H₂S production	-	+	+	+	-	-	+	+	-	-
L-tryptophane	+	+	-	+	+	+	+	-	+	-
Indole production	-	-	-	-	-	+	+	-	-	-
Acetoin production	+	+	+	-	+	+	+	-	+	+
Gelatinase	-	+	+	+	+	+	-	+	+	+
D-glucose	-	-	+	+	-	+	+	+	-	-
D-manitol	-	-	-	-	-	-	-	-	-	-
Inositol	-	-	-	-	-	-	-	-	-	-
D-sorbitol	-	-	-	-	-	-	-	-	-	-
L-rhamnose	+	-	-	-	+	-	-	-	-	-
D-sucrose	-	-	-	-	-	-	-	-	-	-
D-melibiose	+	-	-	+	+	-	-	-	+	-
Amygdalin	+	-	+	+	-	-	-	-	-	-
L-arabinose	+	+	+	-	-	+	-	-	-	-
Cytochrome-oxidase	+	+	+	-	+	-	+	+	+	+

Table 5.

Antibiotic resistance spectra of thermophilic bacterial strains by the disc diffusion method.

Strains	Ax	MET	LEV	SXT	Cip	CFM	PRL	AZM	CLR	LNZ
MBT006	24 (S)	0 (R)	32 (S)	25 (S)	30 (S)	16 (S)	18 (S)	0 (R)	41 (S)	17 (S)
MBT007	29 (S)	0 (R)	25 (S)	16 (S)	42 (S)	0 (R)	34 (S)	21 (S)	0 (R)	20 (S)
MBT008	17 (S)	0 (R)	24 (S)	0 (R)	16 (S)	24 (S)	19 (S)	0 (R)	30 (S)	0 (R)
MBT009	26 (S)	0 (R)	24 (S)	19 (S)	0 (R)	17 (S)	0 (R)	23 (S)	0 (R)	0 (R)
MBT010	16 (S)	0 (R)	22 (S)	24 (S)	41 (S)	0 (R)	18 (S)	0 (R)	26 (S)	16 (S)
MBT011	30 (S)	0 (R)	20 (S)	26 (S)	30 (S)	19 (S)	16 (S)	0 (R)	32 (S)	26 (S)
MBT012	21 (S)	16 (S)	24 (S)	0 (R)	17 (S)	30 (S)	36 (S)	18 (S)	0 (R)	0 (R)
MBT013	24 (S)	0 (R)	32 (S)	20 (S)	0 (S)	30 (S)	18 (S)	0 (R)	26 (S)	32 (S)
MBT014	17 (S)	0 (R)	30 (S)	16 (S)	28 (S)	18 (S)	26 (S)	0 (R)	32 (S)	30 (S)
MBT018	18 (S)	0 (R)	18 (S)	34 (S)	26 (S)	18 (S)	30 (S)	17 (S)	16 (S)	20 (S)

Table 6.

List of thermophilic bacteria identified on the basis of 16S rRNA genes.

Isolate code	Closest affiliation	Accession number	Percentage similarity
MBT008	<i>Anoxybacillus kamchatkensis</i>	OM918284	100
MBT009	<i>Anoxybacillus gonensis</i>	OM918285	99.83
MBT012	<i>Anoxybacillus mongoliensis</i>	OM918286	99.57
MBT014	<i>Anoxybacillus tengchongensis</i>	OM918287	99.43
MBT018	<i>Anoxybacillus karvacharensis</i>	OM918288	98.70