

New species of *Lathrolestes* Förster (Hymenoptera: Ichneumonidae) from Côte d'Ivoire

Alexey Reshchikov ‡

‡ Department of Zoology, Swedish Museum of Natural History, Stockholm, Sweden

Corresponding author:

Academic editor: Jose Fernandez-Triana

ZooBank: [urn:lsid:zoobank.org:pub:318A4D11-CEF2-4EC2-A8D2-319B0B858171](https://doi.org/10.21203/rs.3.rs-1111111/v1)

Abstract

Lathrolestesivoriensis **sp. n.** is described from Côte d'Ivoire. This is the second record of the genus from the Afrotropical region and the first record of the genus and the subfamily for the country. Illustrated re-description of *L.ruwenzoricus* (Benoit 1955) is also provided.

Keywords

Ctenopelmatinae, Perilissini, *Lathrolestes*, *Lathrolestes ivoriensis*, *Lathrolestes ruwenzoricus*, Afrotropical region, Côte d'Ivoire.

Introduction

Very few is known about the diversity of Afrotropical Ichneumonidae, excluding the excellent revision of Ophioninae by Gauld & Mitchell (Gauld and Mitchell 1978), seminal work by Seyrig (Seyrig 1935) and a series of recent papers (Bennett and Barnes 2011, Rousse et al. 2013, Yu et al. 2012). However most of those paper were focused in Madagascar or South Africa. Only 1979 species of ichneumonids had been recorded from the whole Afrotropical region (Yu et al. 2012). This is less than half as many as are known from Germany! The African fauna of the large subfamily Ctenopelmatinae (Hymenoptera, Ichneumonidae) is very poorly known – there are currently only ten species known from tropical Africa (Aubert 1987, Benoit 1955, Cameron 1911, Morley 1926, Seyrig 1935, Szépligeti 1908, Townes and Townes 1973).

The genus *Lathrolestes* Förster, 1869 (Hymenoptera, Ichneumonidae) is a large genus with 99 described species (Reshchikov et al. 2012, Reshchikov 2013). Only one species was previously described from the Afrotropical region (Benoit 1955). No species of *Lathrolestes* had been recorded in Côte d'Ivoire or anywhere else in Africa except

Democratic republic of the Congo (Benoit 1955) before this work. One species, *Lathrolestes ivoriensis* sp. n. is described in this paper as new. Taking into consideration how rare species of the genus *Lathrolestes* are in Africa, and the distinguishing combination of character states, this species is described from only one specimen.

Materials and methods

This work is based on the material of the Royal Museum for Central Africa (RMCA), Tervuren, Belgium. Morphological terminology used in the study follows that of Gauld (Gauld 1997). The female holotype specimen of *Lathrolestes ruwenzoricus* (Benoit, 1955) was examined, re-described and illustrated. Photographs were taken with a Canon Digital Camera 7D, combined with Zerene ®. The following collections were checked for Afrotropical material: American Entomological Institute, Gainesville; Academy of Natural Sciences of Philadelphia; Hungarian Museum of Natural History, Budapest; Natural History Museum, London; National Museum of Natural History, Paris; "Naturalis" Biodiversity Centre, Leiden; Swedish Museum of Natural History, Stockholm; University of Tartu. Not any specimens of the genus *Lathrolest* were found in these collections.

Taxon treatments

Lathrolestes ivoriensis Reshchikov sp. nov.

- ZooBank [urn:lsid:zoobank.org:act:B7B1AEA3-2DDD-48DE-B9AF-71384AFCFE5A](https://zoobank.org/act:B7B1AEA3-2DDD-48DE-B9AF-71384AFCFE5A)

Material

Holotype:

- a. country: Côte d'Ivoire; stateProvince: Abengourou; verbatimLocality: Amangouakro; verbatimLatitude: 6°52'3.79"N; verbatimLongitude: 3°45'48.97"W; eventDate: xii.1962; individualCount: 1; sex: female; recordedBy: J. Decelle; institutionCode: RMCA; occurrenceID: C5BE5EEB-1B29-5AA7-81CE-EAAD58589960

Description

Body length 10 mm (Fig. 1a). Antenna with 35 flagellomeres. Scape 1.54 times as long as wide (Fig. 1b). Head narrowed behind eyes (Fig. 2b), matt, sparsely and shallowly punctate on granulated surface. Maximum length of temple 0.63X transverse eye diameter; minimum length of temple 0.53X transverse eye diameter. Width of face 1.18X height of eye (Fig. 1b), in dorsolateral profile very slightly convex, with bulge, sparsely (more densely in the middle) and shallowly punctate on granulated surface; frons the same; interspace between hind half of lateral ocellus and eye and vertex matt, with shallow sparse punctures, 3.36X of transverse ocellus diameter (Fig. 2b). Clypeus not separated from face, sparsely punctate, very slightly projecting anteriorly (Fig. 1b); apical margin of clypeus obtuse. Tentorial pit large and elongate. Malar space 0.76X basal mandible width, its margin banded by rugosity starting from tentorial pit.

Lower mandible tooth longer than upper. Occipital carina medially complete, reaching hypostomal carina at base of mandible.

Mesosoma matt. Notauli vestigial (Fig. 2b). Mesopleuron matt, granulated, with weak sparse punctures. Tarsus with apical article not enlarged (Fig. 2a). Claws pectinate till its half with not high teeth (Fig. 2a). Wings infusate (Fig. 1a). Fore wing with areolet petiolate. Rs intercepting pterostigma before its middle. 2m-cu straight, with single bulla. Hind wing with cu-a intercepted below middle. Metapleurum granulated. Propodeum matt, impunctate, basal transverse carina and costula absent, area superomedia elongate (Fig. 2d). Body with fine moderately dense setae.

Metasoma matt, evenly covered with moderately short dense setae, shallowly punctate. First metasomal tergite 2.6X as long as apically wide; without shallow median longitudinal impression; bordered by lateral longitudinal carinae and dorsal longitudinal carinae (rather good defined in middle (Fig. 2e). Second metasomal tergite slightly elongate (Fig. 2f). Ovipositor straight, stout dorsally, without notch (Fig. 2c), as long as metasomal height.

Coloration (Figs 1, 2). Face, basal part of clypeus, malar space, basal part of mandible, temples around eye yellow. Apical part of clypeus, mandible, frons, vertex and outer part of temple black. Antenna, palps, mostly pronotum, notum (excluding horseshoe spot in its hind part), tegula, veins and pterostigma, coxae, trochanters, tarsi, hind legs entirely, hind part of 1st and 2nd metasomal tergites and entirely further tergites, ovipositor sheath dark brown. Lower part of pronotum, horseshoe spot in hind part of notum, scutellum, postscutellum, propodeum, mesopleurum, femur and tibia of fore and middle legs, 1st and 2nd (excluding their hind parts) reddish.

Diagnosis

This species differs from other species of the genus by the combination of the following character states: head narrowed behind eyes, clypeus not separated from face, margin of malar space banded by rugosity starting from tentorial pit, wings infusate, claws pectinate till its half with not high teeth (Fig. 2a), basal transverse carina and costula absent, area superomedia elongate (Fig. 2d), ovipositor straight, stout dorsally, without notch (Fig. 2c), as long as metasomal height. This species morphologically closely related to *Lathrolestesjennyae* Gauld, 1997, *L.xochiquetzalis* Reshchikov, 2011 and *L. kukulkanis* Reshchikov, 2011 (all from Central America) with which it shares similar coloration of clypeus (yellow and black in apical part) and well defined dorsal longitudinal carinae of 1st metasomal tergite. The new species clearly differs from Mexican species by the longer first metasomal tergite (2.6X as long as apically wide versus shorter than 1.6–1.9 X as long in the Mexican species), absence of basal transverse carina of the propodeum and costula (Fig. 2d), ovipositor as long as metasomal height, straight, stout dorsally, without notch (Fig. 2c) and unique coloration (Figs 1, 2) (see description). From the single known Afrotropical species, *L. ruwenzoricus* (Benoit, 1955) the new species is distinguished on the basis of colour (*L. ruwenzoricus* is entirely black except for reddish fore legs).

Etymology

The species epithet refers to the name of the country where it was collected.

Distribution

Côte d'Ivoire.

Lathrolestes ruwenzoricus (Benoit, 1955)

Description

Body length 12 mm (Fig. 3). Scape 1.08 times as long as wide (Fig. 4a). Head narrowed behind eyes (Fig. 4b), matt, sparsely and shallowly punctate on smooth surface. Maximum length of temple 1.3X transverse eye diameter; minimum length of temple 1.07X transverse eye diameter. Width of face 1.18X height of eye; in dorsolateral profile very slightly convex, with bulge, sparsely and shallowly punctate on smooth surface; frons the same; interspace between hind half of lateral ocellus and eye and vertex matt, with shallow sparse punctures, 2.58X of transverse ocellus diameter. Clypeus not separated from face, sparsely punctate, very slightly projecting anteriorly; apical margin of clypeus obtuse. Tentorial pit large and elongate. Malar space equal basal mandible width, its margin banded by rugosity starting from tentorial pit. Lower mandible tooth longer than upper. Occipital carina medially complete, reaching hypostomal carina at base of mandible.

Mesosoma matt. Notauli shallowly imprinted at base. Mesopleuron matt, smooth, with weak sparse punctures. Tarsus with apical article not enlarged (Fig. 4c). Claws pectinate (Fig. 4c). Wings hyaline (Fig. 3). Fore wing with areolet petiolate. Rs intercepting pterostigma far before its middle. 2m-cu straight, with single bulla. Hind wing with cu-a intercepted below middle. Metapleurum slightly wrinkled. Propodeum matt, impunctate, only apical areas defined (Fig. 4d). Body with fine moderately dense setae.

Metasoma matt, evenly covered with moderately short dense setae, shallowly punctate. First metasomal tergite 1.52X as long as apically wide (Fig. 4e); with very weak shallow median longitudinal impression; bordered by lateral longitudinal carinae, dorsal longitudinal carinae absent. Second metasomal tergite square (Fig. 4f). Ovipositor straight, without notch, as long as metasomal height.

Coloration (Figs 3, 4). Body mostly black. Fore femur, tibia and basal tarsomer reddish.

Diagnosis

This species differs from other members of the genus by the combination of the following character states: body mostly black excluding fore femur, tibia and basal

tarsomerus which are reddish; matt, sparsely and shallowly punctate on smooth surface; Rs intercepting pterostigma far before its middle; propodeum matt and impunctate with only apical areas defined. This species very easily can be recognized from *L. ivoriensis* Reshchikov, sp. n. by unusually black and dim coloration of body, propodeal carinae with only apical areas present and absence of dorsal longitudinal carinae of 1st metasomal tergite.

Distribution

Democratic Republic of the Congo

Acknowledgements

I would like to thank Dr. Eliane De Coninck and *Stéphane Hanot* (RMCA) for provided material, Dr. Andrew Bennett and Dr. Jose Fernandez-Triana (CNC) for valuable suggestions and corrections of the manuscript.

References

- Aubert JF (1987) Deuxième prélude à une révision des Ichneumonides Scolobatinae. Bulletin de la Société Entomologique de Mulhouse 1987: 33-40.
- Bennett A, Barnes D (2011) Revision of the world species of *Zambion* (Hymenoptera, Ichneumonidae, Tryphoninae). ZooKeys 159: 19. <https://doi.org/10.3897/zookeys.159.2219>
- Benoit PL (1955) Contributions a l'étude des Ichneumonidae africains (Hymenoptera). Annales du Musée Royal du Congo Belge 38: 1-55.
- Cameron P (1911) On the parasitic Hymenoptera collected by Mr. A.J.T. Janse, Transvaal. Annals of the Transvaal Museum 2: 173-217.
- Gauld ID (1997) The Ichneumonidae of Costa Rica, 2. Introduction and keys to species of the smaller subfamilies, Anomaloninae, Ctenopelmatinae, Diplazontinae, Lycorininae, Phrudinae, Tryphoninae (excluding Netelia) and Xoridinae, with an appendices on the Rhyssinae. 57. Memoirs of the American Entomological Institute, 585 pp.
- Gauld ID, Mitchell PA (1978) Taxonomy, Distribution and Host Preferences of African Parasitic Wasps of the Sub-family Ophioninae. Imprint unknown (December 1978), 186 pp. [ISBN 0851984096]
- Morley C (1926) On some South African Ichneumonidae in the collection of the South African Museum. Annals of the South African Museum 23: 435-481.
- Reshchikov A (2013) Two new species of *Lathrolestes* (Hymenoptera, Ichneumonidae) from Norway, northern Russia and Finland with a key to western Palaearctic species. Zootaxa 3681 (1): 59. <https://doi.org/10.11646/zootaxa.3681.1.3>
- Reshchikov A, Veijalainen A, Saaksjarvi I (2012) A new species of *Lathrolestes* (Hymenoptera, Ichneumonidae) from Ecuadorian Amazonia, with a key to the Neotropical species of the genus. ZooKeys 251: 21-27. <https://doi.org/10.3897/zookeys.251.3709>

- Rousse P, Villemant C, Seyrig A (2013) Ichneumonid wasps from Madagascar. VI. The genus *Pristomerus* (Hymenoptera: Ichneumonidae: Cremastinae). European Journal of Taxonomy 49: 1-38. <https://doi.org/10.5852/ejt.2013.49>
- Seyrig A (1935) Mission scientifique de l'Omo. Tome III. Fascicule 18. Hymenoptera, II. Ichneumonidae: Cryptinae, Pimplinae, Tryphoninae et Ophioninae. Mémoires du Muséum National d'Histoire Naturelle 4: 1-100.
- Szépligeti G, Y. S (1908) Hymenoptera: Braconidae & Ichneumonidae. Wissenschaftliche Ergebnisse der Schwedischen Zoologischen Expedition nach dem Kilimandjaro, dem Meru und den umgebenden Massaisteppe. Tryckt hos P. Palmquists aktiebolag, 1910, Stockholm, 25-96 pp.
- Townes HK, Townes M (1973) A catalogue and reclassification of the Ethiopian Ichneumonidae. 19. Memoirs of the American Entomological Institute, 416 pp.
- Yu DS, Horstmann K, van Achterberg K (2012) Taxapad. URL: <http://www.taxapad.com>




Figure 1.
Holotype female of *L. ivoriensis* Reshchikov, sp. n.
a: habitus;
b: face.




Figure 2.

Holotype female of *Livoriensis* Reshchikov, sp. n.

a: claw;

b: head;

c: ovipositor;

d: propodeum;

e: 1st metasomal tergite;
f: 2nd metasomal tergite.



Figure 3.
Holotype female of *L. ruwenzoricus* (Benoit 1955), habitus.




Figure 4.
Holotype female of *L. ruwenzoricus* (Benoit 1955)

a: face;
b: head;
c: claw;
d: propodeum;
e: 1st metasomal tergite;
f: 2nd metasomal tergite.