

A new record of *Potamanthellus caenoides* Ulmer 1939 (Ephemeroptera: Neoephemereidae) from the southern Western Ghats of India

C. Selvakumar[‡], K.G. Sivaramakrishnan[§], S. Janarthanan[‡]

[‡] Department of Zoology, University of Madras, Guindy Campus, Chennai-600025, India

[§] Flat 3, Door No. 7, Gokulam Apartments, Gokulam Colony, West Mambalam, Chennai-600033, India

Corresponding author:

Academic editor: Benjamin Price

Abstract

Background

As part of ongoing exploration of the mayflies of hill streams of the southern Western Ghats of India, we establish a new record of mayfly.

New information

Potamanthellus caenoides Ulmer 1939 is newly recorded based on larval collection from the upstream of Silent Valley National Park of the southern Western Ghats. Brief ecological notes are appended.

Keywords

Ephemeroptera, new record, *Potamanthellus caenoides*, southern Western Ghats, India

Introduction

Ephemeroptera is a biogeographically significant archaic order of aquatic insects abounding in several enigmatic families in the pantropical region, especially in the Oriental Realm. Neoephemereidae is a small group of mayflies presently confined to Holarctic and Oriental regions. Bae 1998 recognized only three genera viz., *Potamanthellus* Lestage 1930 (seven species), *Neoephemera* McDunnough 1925 (five species) and *Ochernova* Bae 1998 (one species), synonymising *Leucorhoenanthus* Lestage 1930 (one species) with *Neoephemera*. This is not accepted by Kluge 2004 and

Bauernfeind and Soldán 2012. Larvae of Neophemeridae have unique operculate gills on the second abdominal segment that are fused medially. The larvae of *Potamanthellus* are distinguished from those of *Neophemera* and *Ochernova* by their densely setate mouthparts, by their lack of well developed lateral expansions of the pronotum and mesonotum, and by their possession of rows of long setae on the caudal filaments (Bae 1998). However, the larvae of *Potamanthellus* cannot be differentiated from *Leucorhoenanthus* by any plesiomorphic or apomorphic larval traits. Based on our larval collections of Ephemeroptera of the Western Ghats, we establish a new record of *Potamanthellus caenoides* (Ulmer 1939), which is a significant range extension to south Asia from its known range viz., Sumatra Island located in Southeast Asia. Differential diagnosis of *P. caenoides* is verified based on the larval descriptions given (Ulmer 1939) and subsequent revisionary studies (Bae 1998). Brief ecological notes are appended.

Materials and methods

The material used for this study was collected from the up-streams of Silent Valley National Park in south western region of the Western Ghats of peninsular India. The specimens were preserved in 85% ethanol. Some specimens were mounted on slides to enable detailed microscopic observations. Photographs were taken on the stereozoom and brightfield microscopes (Magnus and Nikon Eclipse 80i).

Taxon treatment

Potamanthellus caenoides Ulmer 1939

Material

- a. taxonomicStatus: accepted; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Ephemeroptera; family: Neophemeridae; taxonRank: species; genus: *Potamanthellus*; specificEpithet: *caenoides*; country: INDIA; stateProvince: Kerala; municipality: Silent Valley National Park; locality: Poochipara; verbatimElevation: 935 m; verbatimLatitude: 11°06'49.5" N; verbatimLongitude: 76°25'52.4" E; samplingProtocol: Hand picking; year: 2013; month: April; day: 18; habitat: Cascade; individualCount: 7; sex: male & female; lifeStage: Larva; identifiedBy: C. Selvakumar & K. G. Sivaramakrishnan; occurrenceID: 29C0045E-2D6B-54F7-AAB1-F38DA6F7FE0E

Diagnosis

Potamanthellus caenoides is distinguished from other species of *Potamanthellus* by the following combination of characters in larvae: (i) a distinct diagonal ridge on operculate gills (Fig. 1); (ii) distinct tubercles on abdominal terga 6–8 (Fig. 1); (iii) dorsal forefemora with transverse row of setae (Fig. 2); (iv) relatively small body size (<8 mm) (Fig. 1) and (v) relatively short caudal filaments that possess strongly developed lateral setae (Fig. 1). *P. caenoides* is distinguished from closely related species *P. ganges* by the following characters: (i) posteromedian tubercle on

abdominal terga 1–2 and 6–8 distinct (Fig. 1); (ii) rows of hairlike setae strongly developed and mature body ca. 6-8 mm (Fig. 1) and (iii) dorsal forefemora with transverse row of setae (Fig. 2).

Distribution

Indonesia (Sumatra (Ulmer 1939), Java, Bali, Lombok and Flores), Malaysia (Malay peninsula, Sabah and Sarawak), Philippines (Mindanao), Thailand (Bae 1998), Vietnam (Nguyen and Bae 2003) and India (southern Western Ghats).

Biology

The larvae of *P. caenoides* occur in moderately fast flowing mountain streams and rivers ranging 850-935 m in altitude. The streams and rivers are canopied by predominant riparian trees. The substrates consist of relatively coarse particles (boulder 30%, cobble 20%, pebble 20% and gravel and sand 30%), fallen leaves and detritus. The water temperature in April ranges 18-23°C. Larvae were collected by Kick samples and hand picking.

Taxon discussion

Presently this genus consists of seven species viz. *P. amabilis* (Eaton 1892), *P. caenoides* (Ulmer 1939), *P. chinensis* (Hsu 1936), *P. edmundsi* (Bae 1998), *P. ganges* (Bae 1998), *P. shaowuensis* (Gui et al. 1999) and *P. unicutibius* (Nguyen and Bae 2003). However, only one species viz., *P. ganges* is known from India from the tributary of Ganges (Bae 1998). *P. caenoides* (Ulmer 1939) is new record from the southern Western Ghats and second species from India.

Acknowledgements

The authors are grateful to Jobin C. Tharian for donation of larval material from his collections. They thank Dr. K. A. Subramanian, Zoological Survey of India, Kolkata for the photo of entire larva. C. Selvakumar thanks University Grants Commission (UGC), New Delhi, India for the award of Dr. D. S. Kothari Post Doctoral Fellowship [No.F.4-2/2006 (BSR)/13-670/2012 (BSR)].

Author contributions

All authors are equally contributed.

References

- Bae YJ (1998) Phylogenetic Systematics and Biogeography of the Neoephemeridae (Ephemeroptera: Pannota). *Aquatic Insects* 20: 35-68. <https://doi.org/10.1076/aqin.20.1.35.4489>
- Bauernfeind E, Soldán T (2012) *The mayflies of Europe (Ephemeroptera)*. Apollo Books, Ollerup, 781 pp. <https://doi.org/10.1163/9789004260887>
- Eaton AE (1892) New species of Ephemeridae from the Tenasserim Valley. *Transactions of the Entomological Society of London* 1892: 185-190. <https://doi.org/10.1111/j.1365-2311.1892.tb02045.x>
- Gui H, Zhou CF, Sc C (1999) Ephemeroptera. In: Huang BK (Ed.) *Fujian Insect Fauna*. 1. Fujian Science and Technology Press, Fuzhou.
- Hsu Y (1936) New Chinese mayflies from Kiangsi Province (Ephemeroptera). *Peking Natural History Bulletin* 10: 119-126.
- Kluge N (2004) *The Phylogenetic system of Ephemeroptera*. Kluwer Academic Publishers, Dordrecht, 442 pp. <https://doi.org/10.1007/978-94-007-0872-3>
- Lestage J (1930) Contribution à l'étude des larves des Éphéméroptères. VI. - Les larves dites fouisseuses. - Le regime des larves. - Les larves et les poisons. *Bulletin et Annales de la Société Entomologique de Belgique* 70: 79-89.
- McDunnough J (1925) New Canadian Ephemeridae with Notes III. *Canadian Entomology* 57: 168-176. <https://doi.org/10.4039/Ent57168-7>
- Nguyen VV, Bae YJ (2003) Taxonomic review of the Vietnamese Neoephemeridae (Ephemeroptera) with description of *Potamanthellus unicutibius*, new species. *Pan-Pacific Entomologist* 79: 230-236.
- Ulmer G (1939) Eintagsfliegen (Ephemeropteren) von den Sunda-Inseln. *Archiv für Hydrobiologie* 16: 443-692.



Figure 1.
Dorsal view of *Potamanthellus caenoides* Ulmer 1939.




Figure 2.

Potamanthellus caenoides Ulmer 1939.

a: Foreleg

b: Dorsal forefemora with transverse row of setae