

# Checklist of bees (Hymenoptera: Apoidea) from managed emergent wetlands in the lower Mississippi Alluvial Valley of Arkansas

Phillip L Stephenson<sup>‡</sup>, Terry L Griswold<sup>§</sup>, Michael S Arduser<sup>l</sup>, Ashley P G Dowling<sup>¶</sup>, David G Krementz<sup>#</sup>

<sup>‡</sup> Arkansas Cooperative Fish and Wildlife Research Unit, Department of Biological Sciences, University of Arkansas, Fayetteville, AR, United States of America

<sup>§</sup> USDA ARS Pollinating Insects Research Unit, Utah State University, Logan, UT, United States of America

<sup>l</sup> Missouri Department of Conservation - Retired, Webster Groves, MO, United States of America

<sup>¶</sup> Department of Entomology, University of Arkansas, Fayetteville, AR, United States of America

<sup>#</sup> U.S. Geological Survey, Arkansas Cooperative Fish and Wildlife Research Unit, Department of Biological Sciences, University of Arkansas, Fayetteville, AR, United States of America

Corresponding author: Phillip L Stephenson ([phillipleestephenson@gmail.com](mailto:phillipleestephenson@gmail.com)), Terry L Griswold ([terry.griswold@ars.usda.gov](mailto:terry.griswold@ars.usda.gov)), Michael S Arduser ([michael.arduser@mdc.mo.gov](mailto:michael.arduser@mdc.mo.gov)), Ashley P G Dowling ([adowling@uark.edu](mailto:adowling@uark.edu)), David G Krementz ([krementz@uark.edu](mailto:krementz@uark.edu))

Academic editor: Gilberto M. M. Santos

## Abstract

## Background

Here we present the results from a two-year bee survey conducted on 18 managed emergent wetlands in the lower Mississippi Alluvial Valley of Arkansas, USA. Sample methods included pan traps, sweep netting and blue-vane traps. We document 83 bee species and morphospecies in 5 families and 31 genera, of which 37 species represent first published state records for Arkansas. The majority of species were opportunistic wetland species; only a small number were wetland-dependent species or species largely restricted to alluvial plains.

## New information

We present new distributional records for bee species not previously recorded in managed emergent wetlands and report specimens of thirty-seven species for which no published Arkansas records exist, expanding the known ranges of *Ceratina cockerelli*, *Diadasia enavata*, *Lasioglossum creberrimum*, *Svastra cressonii* and *Dieunomia triangulifera*. We also distinguish opportunistic wetland bee species from wetland-dependent and alluvial plain-restricted species.

## Keywords

Apoidea, Arkansas, bee, biodiversity, emergent wetland, Mississippi Alluvial Valley, native species, state record, range expansion

## Introduction

Wetlands of one type or another occur throughout North America and, in some parts of the country, dominate the landscape (Mitsch and Gosselink 2015). Wetlands typically have a unique biota, with numerous obligate and opportunistic species (Niering 1988, Kingsbury and Gibson 2012), including various plants (Lichvar et al. 2016) that provide cover and food for many vertebrates such as migratory birds (Kross et al. 2008, Bellrose 1976). A number of these plants are insect-pollinated or experience enhanced reproduction as a consequence of insect visitation (Harder and Barrett 1992, Lippok et al. 2000, Sohmer and Sefton 1978, Loose et al. 2005, Reader 1975, Klips 1999, Estes and Thorp 1974), indicating that pollination services in wetlands are an important part of wetland systems and their function.

While bees are considered the most important pollinators in most North American communities (Michener 2007), relatively little is known about bee faunas occupying or servicing wetland communities in North America. Some wetland communities have been surveyed, amongst these the Florida Everglades (Pascarella et al. 2000), some north-central Florida wetlands (Hall and Ascher 2010), fens in southern Michigan (Fiedler et al. 2011), playa lakes in Nebraska (Park et al. 2017), cranberry bogs in the northeast US (Loose et al. 2005) and wet flatwoods in Louisiana (Bartholomew and Prowell 2006). These studies and others have demonstrated that the vast majority of bee species found in wetlands also occur in terrestrial habitats and are therefore opportunistic wetland species. In fact, many bees found foraging in wetlands may nest in adjacent terrestrial habitats or in parts of the wetland complex that would not be delineated as, or considered, wetlands based on current definitions (Rust 1980, Cowardin et al. 1979). Nevertheless, a small number of bee species in North America are largely or entirely dependent on wetland communities, either because they depend on the pollen of certain wetland-obligate plants (e.g. *Ptilothrix bombiformis* depends on *Hibiscus* spp. pollen) or have certain nesting or development requirements other than pollen that may be provided only by wetlands, such as certain algae, which may play a role in providing oxygen to soil-nesting immature bees in seasonally flooded sites (Norden et al. 2003).

Reported here is the bee species list from the two-year study monitoring bee communities initiated in 2015 throughout the lower Mississippi Alluvial Valley (LMAV), an area dominated by agriculture and isolated wetlands. Our project represents the only work reported in the emergent wetlands of the LMAV, a region thought to have impaired bee species richness (Koh et al. 2015). Some properties surveyed are managed by the U.S. Fish and Wildlife Service, the Arkansas Game and Fish Commission, the Arkansas Natural

Heritage Commission, while some are privately owned and managed. The purpose of our study was to compile an inventory of the bee fauna of emergent wetlands in the LMAV of Arkansas.

## Materials and methods

### Study Site

The LMAV in Arkansas is bounded on the southwest by the West Gulf Coastal Plain and Ouachita Mountains, on the northwest by the Ozark Plateau and, on the east, by the Mississippi River. The elevation of the LMAV varies by only 46 m throughout the entire 402 km length of the LMAV in Arkansas (Crow 1974). The region is now dominated by agriculture (61% coverage; soybean, rice, corn, sorghum and cotton) with fragments of remnant emergent wetland (1%) and bottomland hardwood forest (17%) (King et al. 2006, USDA National Agricultural Statistics Service (NASS) 2016). The LMAV averages 118-134 cm of rainfall annually, with an average of 35 cm of rainfall between June and September (Scott et al. 1998). All of the 18 sites surveyed (Fig. 1) were used for agricultural or aquacultural production in the past 20 years before being restored to emergent wetlands. All sites had been impounded and were either being managed as moist-soil units, re-established to functioning emergent wetlands through the Wetland Reserve Program (WRP) or were naturally returning to emergent wetlands (Table 1). Wetlands ranged in size from 1 hectare to 50.5 hectares and periodically had standing water based on natural hydrology or water control structures.

Palustrine emergent wetlands are classified as areas <8 ha in size, lacking active wave-formed or bedrock shoreline features, water depth in the deepest part of the basin <2.5 m at low water and salinity due to ocean-derived salts less than 0.5 ppt (Cowardin et al. 1979). This wetland type is sometimes managed using soil disturbance (disking) or water level manipulation (control structures) to produce persistent or non-persistent vegetation for migratory birds (Fredrickson 1991). Persistent vegetation will remain erect when inundated with water and usually include rushes (*Juncus* spp.), cattails (*Typha* spp.), marsh-mallow (*Hibiscus* spp.) and perennial smartweeds (*Persicaria* spp.), while non-persistent vegetation will break over at the water line when inundated with water and usually include grasses (Poaceae), forbs (Asteraceae) and annual smartweeds (*Persicaria* spp.) (Cowardin et al. 1979: 41).

### Collection Methods

We captured bees by placing 10 pan trap stations approximately 20 m apart throughout managed emergent wetlands along a permanent transect following an opportunistic path avoiding open water. Pan trap station platforms held 3,266 ml Solo brand cups that were painted fluorescent blue, fluorescent yellow, or white (Droege et al. 2010, Kirk 1984, Leong and Thorp 1999). These cups were filled  $\frac{3}{4}$  full with soapy water. Pan trap station platforms were adjusted to the average vegetation height at every collection point. We placed traps out at all sites between 0700-0900 hrs and collected them the same day between

1800-2000 hrs. We strained pan trapped bees using a 180  $\mu$ m sieve in the field and transferred them to Whirl-Pak bags with 70% ethanol. We used one blue-vane trap (1.89 l jar) per field site suspended from a shepherd's hook pole, with the bottom of the trap 1 m above the ground (Kimoto et al. 2012, Stephen and Rao 2005). The blue-vane trap was filled with 475 ml of soapy water. These blue-vane traps were placed and collected on the same schedule as the pan traps and samples similarly extracted. We used indirect sweep netting to sample for bees that were not attracted to either pan or blue-vane traps. We conducted 5 random transects of 50 sweeps apiece within each wetland per collection period to capture bees. Sweeps were conducted between 1100-1345 hrs (Stephen and Rao 2007) in 2015 and between 0900-1000 hrs (Roulston et al. 2007) in 2016. Sweep net collection periods were altered between years because we observed bees were more active between 0730-1000 hrs during the previous year. All sweep net samples were placed in 3.8 l Ziploc bags and were placed in the freezer until processed. We sampled each site 4-7 times in 2015 (19 May-18 September) and 8 times each in 2016 (22 May-9 September).

## Species Identification

Bee specimens were washed, dried, pinned and labelled with location information (Stephenson 2017). We identified bees to species when possible or to genus using identification guides and [DiscoverLife.org](https://www.discoverlife.org) (Ascher and Pickering 2017). We confirmed identifications with Harold Ikerd, Katherine Parys, Sam Droege and John Ascher. Voucher specimens are deposited at the University of Arkansas Arthropod Museum, Fayetteville, AR and at the U.S. National Pollinating Insect Collection, Logan, UT USA.

## Range

Species ranges and state records were determined using primary literature and other published accounts (see Literature Cited, below), the North American bee database available at [DiscoverLife.org](https://www.discoverlife.org) (Ascher and Pickering 2017) and, in a few cases, the bugguide website (bugguide.net).

## Wetland Affiliation

We classified bee species as "opportunistic," "wetland-dependent," or "alluvial plain-restricted" based on published accounts and the ongoing surveys of one of the authors (MSA) in selected National Wildlife Refuges on the alluvial plains of the upper and middle Mississippi, lower Missouri and lower Ohio Rivers. The wetland-dependent and alluvial plain-restricted species are indicated by asterisks in the checklist below.

## Checklist

### Family Colletidae

***Colletes nudus* Robertson, 1898**

**Notes:** Widespread east of the Rocky Mountains but not previously recorded from Arkansas (Stephen 1954). Opportunistic (Table 1: Site 15).

***Hylaeus (Hylaeus) mesillae* (Cockerell, 1896)**

**Notes:** Transcontinental but not previously recorded from Arkansas (Snelling 1970). Opportunistic (Table 1: Site 6).

***Hylaeus (Prosopis) affinis* (Smith, 1853)**

**Notes:** Widespread but not previously recorded from Arkansas (Hurd 1979). Opportunistic (Table 1: Site 1-3, 6, 14, 17, 18).

***\*Hylaeus (Prosopis) nelumbonis* (Robertson, 1890)**

**Notes:** New record for Arkansas; previously recorded from Illinois and Maryland south to Florida and Louisiana (Mitchell 1960, Hurd 1979). Wetland specialist (Table 1: Sites 1-7, 10, 12, 14, 15).

***\*Hylaeus (Prosopis) ornatus* Mitchell, 1951**

**Notes:** New record for Arkansas; previously recorded from North Carolina and Florida (Mitchell 1960, Hurd 1979). Wetland specialist (Table 1: Site 4, 5, 14, 15, 17).

***Hylaeus sp. 1***

**Notes:** (Table 1: Site 1).

***Hylaeus sp. 2***

**Notes:** (Table 1: Site 18).

**Family Andrenidae**

***Andrena (Callandrena s.l.) rudbeckiae* Robertson, 1891**

**Notes:** Known from the Great Plains east to North Carolina but not previously recorded from Arkansas (LaBerge 1967). Opportunistic (Table 1: Sites 5, 8).

***Andrena (Leucandrena) macra* Mitchell, 1951**

**Notes:** Known from the southeast to Texas but not previously recorded from Arkansas (LaBerge 1987). Opportunistic (Table 1: Sites 5, 7, 11, 14, 15, 17, 18).

***Andrena (Scapteropsis) imitatrix* Cresson, 1872**

**Notes:** Opportunistic (Table 1: Site 5).

***Andrena (Simandrena) nasonii* Robertson, 1895**

**Notes:** Widespread in eastern North America west to Colorado and Texas but not previously recorded from Arkansas (LaBerge 1989). Opportunistic (Table 1: Site 17).

***Calliopsis (Calliopsima) coloradensis* Cresson, 1878**

**Notes:** Opportunistic (Table 1: Sites 3, 17, 18).

***Panurginus polytrichus* Cockerell, 1909**

**Notes:** Known from the adjacent states of Mississippi, Louisiana, and Texas but not previously recorded from Arkansas (Hurd 1979). Opportunistic (Table 1: Site 5).

***Perdita (Hexaperdita) foveata* Timberlake, 1956**

**Notes:** Opportunistic (Table 1: Site 4).

***Perdita* sp. 1**

**Notes:** (Table 1: Sites 2, 7, 12, 15, 16).

**Family Halictidae**

***Agapostemon angelicus/texanus***

**Notes:** These specimens are most likely *A. texanus*, as the closest records to Arkansas of the predominantly western *A. angelicus* are from SE Oklahoma, while there are a number of *A. texanus* records from Arkansas (Roberts 1972). Opportunistic (Table 1: Sites 1-9, 11-18).

***Agapostemon (Agapostemon) sericeus* (Forster, 1771)**

**Notes:** Opportunistic (Table 1: Sites 1-15, 17-18).

***Agapostemon (Agapostemon) splendens* (Lepeletier, 1841)**

Notes: Opportunistic (Table 1: Sites 4-6, 13).

***Agapostemon (Agapostemon) virescens* (Fabricius, 1775)**

Notes: Opportunistic (Table 1: Sites 1,2, 4, 5, 7-12, 17).

***Augochlora (Augochlora) pura* (Say, 1837)**

Notes: Opportunistic (Table 1: Sites 5, 18).

***Augochlorella aurata* (Smith, 1853)**

Notes: Opportunistic (Table 1: All Sites).

***Augochloropsis (Paraugochloropsis) fulgida* (Smith, 1853)**

Notes: *Augochloropsis fulgida* and *A. metallica* (below) are here recognised as separate species, rather than subspecies as this has been the traditional interpretation (Moure and Hurd 1987). Studies by one of us (MSA) indicate that the two are largely sympatric and their distinguishing morphological features stable; molecular data appear to support this (S. Droege, in litt.). A short key separating the two is available at [DiscoverLife.org](http://DiscoverLife.org), under *Augochloropsis metallica*. Opportunistic (Table 1: Sites 1, 11, 15, 18).

***Augochloropsis (Paraugochloropsis) metallica* (Fabricius, 1793)**

Notes: Opportunistic (Table 1: Sites 1-3, 5, 7, 8, 11, 12, 14).

***\*Dieunomia (Epinomia) triangulifera* (Vachal, 1897)**

Notes: New species record for Arkansas. Common in the central US usually on the alluvial plains of major rivers (Missouri, Arkansas) and their tributaries, east to the Mississippi River corridor and its tributaries in Missouri, Illinois and Indiana, but not recorded any further south along the Mississippi corridor until now (Cross 1958). A primary oligolege of *Helianthus* spp. and an important pollinator of *Helianthus annuus* and commercial sunflowers (Minckley *et al.* 1994). Primarily associated with alluvial plains of large rivers, not wetlands per se (Table 1: Site 8).

***Halictus (Nealictus) parallelus* Say, 1837**

Notes: Opportunistic (Table 1: All Sites).

***Halictus (Odontalictus) ligatus* Say, 1837**

Notes: Opportunistic (Table 1: All Sites).

***Halictus (Protohalictus) rubicundus* (Christ, 1791)**

**Notes:** This appears to be the first published Arkansas record of this common, widespread Holarctic species (Moure and Hurd 1987). Opportunistic (Table 1: Sites 1, 4, 9, 15, 18).

***Lasioglossum (Dialictus) bruneri* (Crawford, 1902)**

**Notes:** New species record for Arkansas; widespread in the eastern US (Gibbs 2011). Opportunistic (Table 1: Sites 1-3, 8, 9, 15).

***Lasioglossum (Dialictus) callidum* (Sandhouse, 1924)**

**Notes:** New species record for Arkansas; widespread in the eastern US (Gibbs 2011). Opportunistic (Table 1: Site 4).

***Lasioglossum (Dialictus) creberrimum* (Smith, 1853)**

**Notes:** New species record for Arkansas. *L. creberrimum* is a southeastern species, occurring largely along the coast from southeast Texas up to Maryland, with scattered inland records (Gibbs 2011). The Arkansas specimens represent the furthest inland occurrence of this species to date. Opportunistic (Table 1: All Sites).

***Lasioglossum (Dialictus) cressonii* (Robertson 1890)**

**Notes:** New species record for Arkansas; these specimens may represent the southernmost records for this common and widespread species (Gibbs 2011). Opportunistic (Table 1: Sites 1, 2, 4, 5, 7, 9, 12, 15).

***Lasioglossum (Dialictus) hartii* (Robertson, 1892)**

**Notes:** Restricted to alluvial plains and riparian corridors (Table 1: All Sites).

***Lasioglossum (Dialictus) hitchensi* Gibbs, 2012**

**Notes:** New species record for Arkansas. Widespread in the eastern US (Gibbs 2011, as *L. mitchelli* Gibbs 2010). Opportunistic (Table 1: Site 13).

***Lasioglossum (Dialictus) pilosum* (Smith, 1853)**

**Notes:** New species record for Arkansas. Occurs over much of the eastern US (Gibbs 2011). Opportunistic (Table 1: All Sites).



***Lasioglossum (Dialictus) sp. 1***

Notes: (Table 1: Sites 9, 13, 15, 16).

***Lasioglossum (Dialictus) sp. 2***

Notes: (Table 1: Sites 2, 10).

***Lasioglossum (Hemihalictus) lustrans* (Cockerell, 1897)**

Notes: Opportunistic (Table 1, Sites 2, 3, 5, 7, 9, 15, 18).

***Lasioglossum (Hemihalictus) nelumbonis* (Robertson, 1890)**

Notes: New species record for Arkansas. Occurs throughout much of the eastern US (Gibbs et al. 2013); primarily associated with alluvial plains and wetlands, but does occur in upland wetlands, upland pond margins, riparian areas etc (Table 1: All Sites).

***Lasioglossum sp. 3***

Notes: (Table 1: Sites 5, 9).

***Nomia (Acunomia) nortoni* Cresson, 1868**

Notes: Opportunistic (Table 1: Sites 5, 12-14).

***Sphecodes mandibularis* Cresson, 1872**

Notes: Opportunistic (Table 1: Site 5).

**Family Megachilidae**

***Dianthidium (Dianthidium) subrufulum* Timberlake, 1943**

Notes: Opportunistic (Table 1: Site 18).

***Megachile (Acentron) albitarsis* Cresson, 1872**

Notes: Widespread in eastern United States to Arizona but previously unrecorded from Arkansas (Mitchell 1937b). Opportunistic (Table 1: Sites 2, 3, 5, 7, 8, 10, 14, 15, 17).

***Megachile (Chelostomoides) campanulae* (Robertson, 1903)**

**Notes:** Widespread in eastern United States west into Great Plains but previously unrecorded from Arkansas (Mitchell 1937a, Snelling 1990). Opportunistic (Table 1: Site 13).

***Megachile (Leptorachis) petulans* Cresson, 1878**

**Notes:** Widespread in eastern United States to Arizona but previously unrecorded from Arkansas (Mitchell 1937b). Opportunistic (Table 1: Sites 5, 7, 14, 17, 18).

***Megachile (Litomegachile) brevis* Say, 1837**

**Notes:** Widespread but previously unrecorded for Arkansas (Mitchell 1935, Bzdyk 2012). Opportunistic (Table 1: Sites 3-13, 15-18).

***Megachile (Litomegachile) mendica* Cresson, 1878**

**Notes:** Opportunistic (Table 1: Sites 1, 2, 10, 13-15).

***Megachile (Litomegachile) texana* Cresson, 1878**

**Notes:** Widespread but previously unrecorded for Arkansas (Mitchell 1935, Bzdyk 2012). Opportunistic (Table 1: Sites 2, 3, 15).

**Family Apidae**

***Anthophorula (Anthophorisca) asteris* (Mitchell, 1962)**

**Notes:** New species record for Arkansas; a fairly widespread (Texas to Georgia to Indiana) but infrequently-collected species (Ascher and Pickering 2017, Timberlake 1980). Opportunistic (Table 1: Site 15).

***Apis (Apis) mellifera* Linnaeus, 1758**

**Notes:** Opportunistic (Table 1: All Sites).

***Bombus (Cullumanobombus) fraternus* (Smith, 1854)**

**Notes:** Opportunistic (Table 1: Sites 5, 6, 14).

***Bombus (Cullumanobombus) griseocollis* (De Geer, 1773)**

**Notes:** Opportunistic (Table 1: Sites 1, 4, 8, 15).

***Bombus (Pyrobombus) bimaculatus* Cresson, 1863**

Notes: Opportunistic (Table 1: Sites 3, 8, 14, 15).

***Bombus (Pyrobombus) impatiens* Cresson, 1863**

Notes: Opportunistic (Table 1: Sites 3, 4, 13-15, 18).

***Bombus (Thoracobombus) pensylvanicus* (De Geer, 1773)**

Notes: Opportunistic Table 1: Sites 1, 2, 5-12, 16, 17).

***Ceratina (Ceratinula) cockerelli* H.S. Smith, 1907**

Notes: New species record for Arkansas; a southern and southeastern species, occurring from Texas to South Carolina (Daly 1973). Opportunistic (Table 1: Sites 7, 16).

***Ceratina (Zadontomerus) dupla* Say, 1837**

Notes: New species record for Arkansas, based on a male specimen. Common throughout much of the eastern half of the US (Daly 1973, Rehan and Sheffield 2011). Opportunistic (Table 1: Site 2).

***Ceratina (Zadontomerus) sp. 1***

Notes: (Table 1: Sites 1-3, 9, 12, 14, 15, 18).

***Ceratina sp. 2***

Notes: (Table 1: Site 2).

***Diadasia (Diadasia) enavata* (Cresson, 1872)**

Notes: New species record for Arkansas. This Asteraceae specialist occurs throughout most of the western half of the US; our specimens represent the easternmost location of the species published to date (Hurd et al. 1980), but they have been collected in some parts of Mississippi (Dr. Katherine Parys 2018, pers. comm., 8 February). Opportunistic (Table 1: Sites 5, 7-11, 15-18).

***Eucera (Synhalonia) hamata* (Bradley, 1942)**

Notes: New species record for Arkansas. Occurs throughout much of the eastern two-thirds of the US, but is absent from the states south and west of Arkansas (Hurd 1979, Timberlake 1969). Opportunistic (Table 1: Sites 2, 5, 7-9).

### ***Eucera (Synhalonia) rosae* (Robertson, 1900)**

Notes: Opportunistic (Table 1: Sites 2, 3, 10, 12, 13, 17, 18).

### **\**Florilegus (Florilegus) condignus* (Cresson, 1878)**

Notes: New species record for Arkansas. A very widespread species, occurring throughout much of the eastern two-thirds of the US into Mexico, Central America and well into South America. No other native North American bee species has a similar or as extensive range. Populations in eastern North America are strongly associated with pickerelweed (*Pontedaria cordata* L.), alluvial plains and natural and constructed wetlands, including upland wetlands. The mouthparts of this species are festooned with hooked hairs (as are the mouthparts of the pickerelweed oligolege *Melissodes apicata* Robertson). *Florilegus condignus* females collect pollen from pickerelweed with their mouthparts as they hover, quickly “stabbing” the mouthparts in and out of the corolla. However, this species is not a strict oligolege of pickerelweed, as it also occurs in wetlands etc. where pickerelweed is absent. LaBerge and Ribble (1966) report western populations of this species to be potentially important pollinators of alfalfa. Primarily a wetland-dependent species in the eastern US portion of its range (Table 1: All Sites).

### ***Melissodes (Eumelissodes) agilis* Cresson, 1878**

Notes: New species record for Arkansas (LaBerge 1961). Opportunistic (Table 1: Sites 6, 7, 9).

### ***Melissodes (Eumelissodes) boltoniae* Robertson, 1905**

Notes: Opportunistic (Table 1: Sites 1, 4, 5, 7, 12-16, 18).

### ***Melissodes (Eumelissodes) denticulatus* Smith, 1854**

Notes: Opportunistic (: Site 17).

### ***Melissodes (Eumelissodes) druriellus* (Kirby, 1802)**

Notes: New species record for Arkansas (LaBerge 1956). Opportunistic (Table 1: Sites 1, 3, 4, 7, 11-15, 18).

### ***Melissodes (Eumelissodes) niveus* Robertson, 1895**

Notes: Opportunistic (Table 1: Site 5).

### ***Melissodes (Eumelissodes) trinodis* Robertson, 1901**

Notes: Opportunistic (Table 1: Site 7, 14, 15).

***Melissodes (Melissodes) bimaculatus* (Lepeletier, 1825)**

Notes: Opportunistic (Table 1: All Sites).

***Melissodes (Melissodes) communis* Cresson, 1878**

Notes: Opportunistic (Table 1: All Sites).

***Melissodes (Melissodes) comptoides* Robertson, 1898**

Notes: Opportunistic (Table 1: All Sites).

***Melissodes (Melissodes) tepaneca* Cresson, 1878**

Notes: Opportunistic (Table 1: Site 5, 8, 10, 11, 13, 15, 16).

***Melitoma taurea* (Say, 1837)**

Notes: Specimens from our study are the only Arkansas specimens we are aware of, but an Arkansas (Newton Co.) image of this species has recently been identified by JSA on bugguide.net, see <https://bugguide.net/node/view/1259116>. Opportunistic (Table 1: Sites 1, 3-6, 10-13, 15, 17).

***\*Ptilothrix bombiformis* (Cresson, 1878)**

Notes: Specimens from our study are the only Arkansas specimens we are aware of, but an Arkansas (Poinsett Co.) image of this species has recently been identified by JSA on bugguide.net, see <https://bugguide.net/node/view/1422145>. A wetlands specialist and Hibiscus oligolege (Rust 1980), but is occasionally found in developed areas some distance from wetlands visiting flowers of ornamental Hibiscus, or cultivated okra (Rau 1930). (Table 1: All Sites).

***Svastra (Brachymelissodes) cressonii* (Dalla Torre, 1896)**

Notes: New species record for Arkansas (LaBerge 1956). Opportunistic (Table 1: Sites 1, 5-7, 10, 13).

***Svastra (Epimelissodes) atripes* (Cresson, 1872)**

Notes: Opportunistic (Table 1: Sites 1-7, 9, 12-15, 17, 18).

***Svastra (Epimelissodes) obliqua* (Say, 1837)**

**Notes:** Opportunistic (Table 1: Sites 1-10, 12-18).

### ***Svastra (Epimelissodes) petulca* (Cresson, 1878)**

**Notes:** Opportunistic (Table 1: Sites 9, 16).

### **\**Triepeolus quadrifasciatus* (Say, 1823)**

**Notes:** The type specimen, which is presumed lost or destroyed, was described from “Arcansa” in 1823 by Thomas Say (Rightmyer 2008). However, the “Arcansas” (Arkansas) of that era was a much larger piece of real estate than the Arkansas of today, then including most of what is now Oklahoma. No additional specimens from Arkansas are mentioned in Rightmyer (2008), thus we consider this specimen the first documentation of the species in Arkansas. Primary host bee is *Svastra atripes*. Opportunistic (Table 1: Site 1).

### ***Xenoglossa (Eoxenoglossa) strenua* (Cresson, 1878)**

**Notes:** New species record for Arkansas (Hurd and Linsley 1967). Opportunistic (Table 1: Sites 2, 6).

### ***Xylocopa (Xylocopoides) virginica* (Linnaeus, 1771)**

**Notes:** Opportunistic (Table 1: Sites 2-4, 8, 9, 11, 12, 14-16, 18).

## **Analysis**

During 201 collection events, between 2015 and 2016, we collected 17,860 bees representing 83 species and morphospecies across 31 genera and five families. Thirty-seven species captured represent new Arkansas state records.

## **Discussion**

Our study expands the known distribution of several of the bee species collected because of the limited documentation in emergent wetlands and especially for the LMAV. Our species list is relevant to other emergent wetlands in the LMAV, but may not reflect bee species in other ecoregions in Arkansas, especially in urban and upland areas along the Arkansas River Valley (see Little 2013). Most of the species collected are widespread in North America and many have been recorded from states that border Arkansas. While their presence in the state may not be surprising, the fact they have not been recorded highlights the lack of published data and surveys performed in this physiographic region and for this state.

### ***Ceratina cockerelli***

*Ceratina cockerelli* is commonly associated with the Gulf Coastal Plains and the lower Piedmont ecoregions, but has been recorded outside of these ecoregions in West Texas. This species is the smallest *Ceratina* in eastern North America and is a generalist often associated with coastal habitats. The specimens collected represent a new state record for Arkansas and have expanded the known range of this species >200 km north into the Mississippi Alluvial Plain of Arkansas from its closest record in southwest Mississippi. These specimens were collected in Monroe and Woodruff Counties, Arkansas (Table 1: Sites 7, 16).

### ***Diadasia enavata***

*Diadasia enavata* is commonly found in the western portion of the United States of America. This species is known to be restricted to plants in the Asteraceae family (Hurd et al. 1980, Linsley and MacSwain 1958). The wetlands surveyed occasionally have *Coreopsis tinctoria* on their edges being visited frequently by *Diadasia enavata*. *Coreopsis tinctoria* is found throughout the continental United States of America often in bottomland areas (USDA, NRCS 2017). The specimens collected represent a new state record for Arkansas and have considerably expanded the known range of this species east into the Mississippi Alluvial Plain of Arkansas from its closest records in Missouri, Oklahoma and Texas (Hurd et al. 1980). These specimens were collected in Arkansas, Lawrence, Monroe and Woodruff Counties, Arkansas (Table 1: Sites 5, 7-11, 15-18).

### ***Dieunomia triangulifera***

*Dieunomia triangulifera* is a specialist of the sunflower genus *Helianthus* and is mainly found west of the Mississippi River and in the Great Plains of the United States of America (Cross 1958, Minckley et al. 1994). The wetlands surveyed had *Coreopsis tinctoria* present in the unit and on the levee. This specimen, collected in Arkansas County, represents a new state record for Arkansas (Table 1: Site 8).

### ***Lasioglossum creberrimum***

*Lasioglossum creberrimum* is commonly associated with the Gulf Coastal Plains and Piedmont ecoregions of the United States of America, but has been recorded outside of these ecoregions in rare cases. This species is considered a generalist and prefers open lands. *Lasioglossum creberrimum* was also collected in remnant prairies and an urban park in the Arkansas River Valley in 2011-2012 (Little 2013). The specimens collected in the present study represent a new state record for Arkansas that expands the known range of this species >200 km north into the Mississippi Alluvial Plain of Arkansas from its closest record in southwest Mississippi. These specimens were collected in Arkansas, Cross, Jackson, Lawrence, Monroe, Prairie, White and Woodruff Counties, Arkansas (Table 1: All Sites).

## ***Svastra cressonii***

*Svastra cressonii* is a species in the subgenus *Brachymelissodes* that is commonly found in the plains states of the south-central portion of the United States of America. This species has been collected as far north as Iowa (LaBerge 1956) and as far south as Louisiana (Ascher and Pickering 2017). The floral preferences of this species are unclear, but *Asteraceae* sp. and *Ludwigia peploides* have been mentioned (Estes and Thorp 1974, Ascher and Pickering 2017). Estes and Thorp (1974) documented *Svastra cressonii* foraging on *Ludwigia peploides* on the edges of farm ponds in Texas. This species of *Ludwigia* was present at all sites surveyed in our study. The collected *Svastra cressonii* specimens have expanded the known range >250 km east of previous documented occurrences. The specimens collected represent a new state record for Arkansas and expand the known range of this species into the Mississippi Alluvial Plain. These specimens were collected in Cross, Monroe, Prairie, White and Woodruff Counties, Arkansas (Table 1: Sites 1, 5-7, 10, 11, 13).

## **Acknowledgements**

Our research was funded by the U.S. Fish and Wildlife Service, the U.S. Department of Agriculture, the U.S. Geological Survey Arkansas Cooperative Fish and Wildlife Research Unit, Arkansas Game and Fish Commission, Arkansas Audubon Society and the University of Arkansas. We would like to thank Harold Ikerd, Katherine Parys, Sam Droege and John Ascher for their insight and identification assistance. Special thanks to Brandon Burdette, Jenny Courtway, Azlee Goode, Philip Mariage and Erik Ostrum for their diligent work in the field collecting data and, in the laboratory, processing samples. Special thanks to Diane Moler for logistical and budgetary assistance. Any use of trade, firm or product names is for descriptive purposes only and does not imply endorsements by the U.S. Government. All work was completed under Special Use Permits from Arkansas Natural Heritage Commission (S-NHCC-16-005), Arkansas Game and Fish Commission (030820161) and the U.S. Fish and Wildlife Service (43513-5-45, 43670-2016-024).

*This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2014-38640-22155 through the Southern Sustainable Agriculture Research and Education program, under sub-award number RD309-129/S000844. USDA is an equal opportunity employer and service provider.*

*Any opinions, findings, conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture.*



## Author contributions

Stephenson helped design the project, conducted the surveys, secured funding and prepared the specimens for identification and curation. Griswold and Arduser identified specimens, archived specimen data and consulted literature review. Dowling and Krementz helped design the project, interpreted data and secured funding. All authors participated in writing this manuscript.

## References


- Ascher JS, Pickering J (2017) Discover Life bee species guide and world checklist (Hymenoptera: Apoidea: Anthophila). Draft-47. [http://www.discoverlife.org/mp/20q?guide=Apoidea\\_species](http://www.discoverlife.org/mp/20q?guide=Apoidea_species). Accessed on: 2017-12-29.
- Bartholomew CS, Prowell D (2006) Comparison of bee diversity in upland and wet flatwood longleaf pine savannas in Louisiana (Hymenoptera: Apoidea). *Journal of the Kansas Entomological Society* 79 (2): 199-206. <https://doi.org/10.2317/0411.01.1>
- Bellrose RC (1976) Ducks, Geese and Swans of North America. 2nd Edition. Stackpole Books, Harrisburg, PA, 540 pp pp.
- Bzdyk EL (2012) A revision of the *Megachile* subgenus *Litomegachile* Mitchell with an illustrated key and description of a new species (Hymenoptera, Megachilidae, Megachilini). *ZooKeys* 221: 31-61. <https://doi.org/10.3897/zookeys.221.3234>
- Cowardin LM, Carter V, Golet FC, LaRoe ET (1979) Classification of wetlands and deepwater habitats of the United States. U. S. Fish and Wildlife Service <https://doi.org/10.5962/bhl.title.4108>
- Cross EA (1958) A revision of the bees of the subgenus *Epinomia* in the New World (Hymenoptera-Halictidae). *The University of Kansas Science Bulletin* 38 (16): 1261-1301. <https://doi.org/10.5962/bhl.part.10975>
- Crow CT (1974) Arkansas Natural Area Plan. Arkansas Department of Planning. University of Arkansas Press, Little Rock, AR.
- Daly HV (1973) Bees of the Genus *Ceratina* in America North of Mexico. 74. University of California Publications, 114 pp.
- Droege S, Tepdino V, Lebuhn G, Link W, Minckley R, Chen Q, Conrad C (2010) Spatial patterns of bee captures in North American bowl trapping surveys. *Insect Conservation and Diversity* 3 (1): 15-23. <https://doi.org/10.1111/j.1752-4598.2009.00074.x>
- Estes J, Thorp R (1974) Pollination in *Ludwigia peploides* ssp. *glabrescens* (Onagraceae). *Bulletin of the Torrey Botanical Club* 101 (5): 272-276. <https://doi.org/10.2307/2484872>
- Fiedler A, Landis D, Arduser M (2011) Rapid shift in pollinator communities following invasive species removal. *Restoration Ecology* 20 (5): 593-602. <https://doi.org/10.1111/j.1526-100x.2011.00820.x>
- Fredrickson LH (1991) Strategies for water level manipulations in moist-soil systems. 13.4.6. U.S. Fish and Wildlife Service, Fort Collins, CO. URL: [http://digitalcommons.unl.edu/icwdmwfm/26?utm\\_source=digitalcommons.unl.edu%2Ficwdmwfm%2F26&utm\\_medium=PDF&utm\\_campaign=PDF](http://digitalcommons.unl.edu/icwdmwfm/26?utm_source=digitalcommons.unl.edu%2Ficwdmwfm%2F26&utm_medium=PDF&utm_campaign=PDF)

- Gibbs J (2010) Revision of the metallic species of *Lasioglossum* (*Dialictus*) in Canada (Hymenoptera, Halictidae, Halictini). *Zootaxa* 2591: 1-382.
- Gibbs J (2011) Revision of the metallic *Lasioglossum* (*Dialictus*) of eastern North America (Hymenoptera: Halictidae: Halictini). *Zootaxa* 3073: 1-216.
- Gibbs J, Packer L, Dumes S, Danforth B (2013) Revision and reclassification of *Lasioglossum* (*Evyllaesus*), *L. (Hemihalictus)* and *L. (Sphecodogastra)* in eastern North America (Hymenoptera: Apoidea: Halictidae). *Zootaxa* 3672 (1): 1-117. <https://doi.org/10.11646/zootaxa.3672.1.1>
- Hall HG, Ascher J (2010) Surveys of bees (Hymenoptera: Apoidea: Anthophila) in natural areas of Alachua County in North-Central Florida. *Florida Entomologist* 93 (4): 609-629. <https://doi.org/10.1653/024.093.0419>
- Harder LD, Barrett SCH (1992) The energy cost of bee pollination for *Pontederia cordata* (Pontederiaceae). *Functional Ecology* 6 (2): 226-233. <https://doi.org/10.2307/2389759>
- Hurd P, Linsley EG (1967) Squash and gourd bees of the genus *Xenoglossa* (Hymenoptera: Apoidea). *Annals of the Entomological Society of America* 60 (5): 988-1007. <https://doi.org/10.1093/aesa/60.5.988>
- Hurd PD (1979) Catalog of Hymenoptera in America North of Mexico. In: Krombein KV, Hurd PD, Smith DR, Burks BD (Eds) Superfamily Apoidea. 2. Smithsonian Institution Press, Washington D.C.
- Hurd PD, LeBerge W, Linsley EG (1980) Principal sunflower bees of North America with emphasis on the Southwestern United States (Hymenoptera, Apoidea). *Smithsonian Contributions to Zoology* 310: 15. <https://doi.org/10.5479/si.00810282.310>
- Kimoto C, DeBano S, Thorp R, Rao S, Stephen W (2012) Investigating temporal patterns of a native bee community in a remnant North American bunchgrass prairie using blue vane traps. *Journal of Insect Science* 12 (108): 1-23. <https://doi.org/10.1673/031.012.10801>
- King S, Twedt D, Wilson RR (2006) The role of the wetland reserve program in conservation efforts in the Mississippi River Alluvial Valley. *Wildlife Society Bulletin* 34 (4): 914-920. [https://doi.org/10.2193/0091-7648\(2006\)34\[914:trotwr\]2.0.co;2](https://doi.org/10.2193/0091-7648(2006)34[914:trotwr]2.0.co;2)
- Kingsbury BA, Gibson J (Eds) (2012) Habitat management guidelines for amphibians and reptiles of the Midwestern United States. Technical Publication HMG-1. 2nd Edition. Partners in Amphibian and Reptile Conservation, 155 pp.
- Kirk W (1984) Ecologically selective coloured traps. *Ecological Entomology* 9 (1): 35-41. <https://doi.org/10.1111/j.1365-2311.1984.tb00696.x>
- Klips R (1999) Pollen Competition as a reproductive isolating mechanism between two sympatric *Hibiscus* species (Malvaceae). *American Journal of Botany* 86 (2): 269-272. <https://doi.org/10.2307/2656942>
- Koh I, Lonsdorf E, Williams N, Brittain C, Isaacs R, Gibbs J, Ricketts T (2015) Modeling the status, trends, and impacts of wild bee abundance in the United States. *Proceedings of the National Academy of Sciences* 113 (1): 140-145. <https://doi.org/10.1073/pnas.1517685113>
- Kross J, Kaminski R, Reinecke K, Penny E, Pearse A (2008) Moist-soil seed abundance in managed wetlands in the Mississippi Alluvial Valley. *Journal of Wildlife Management* 72 (3): 707-714. <https://doi.org/10.2193/2007-100>

- LaBerge WE (1956) A revision of the bees of the genus *Melissodes* in North and Central America. Parts I. (Hymenoptera, Apidae). The University of Kansas Science Bulletin. 37 (18): 911-1194. <https://doi.org/10.5962/bhl.part.24549>
- LaBerge WE (1961) A revision of the bees of the genus *Melissodes* in North and Central America. Part III (Hymenoptera, Apidae). The University of Kansas Science Bulletin. 42 (5): 283-663. <https://doi.org/10.5962/bhl.part.9821>
- LaBerge WE, Ribble DW (1966) Biology of *Florilegus condignus* (Hymenoptera: Anthophoridae), with a description of its larva, and remarks on its importance in alfalfa pollination. Annals of the Entomological Society of America 59 (5): 944-950. <https://doi.org/10.1093/aesa/59.5.944>
- LaBerge WE (1967) A revision of the bees of the genus *Andrena* of the western hemisphere, Part I. *Callandrena*. Bulletin of the University of Nebraska State Museum 7: 1-316.
- LaBerge WE (1987) A revision of the bees of the genus *Andrena* of the western hemisphere, Part XII. Subgenera *Leucandrena*, *Ptilandrena*, *Scoliandrena*, and *Melandrena*. Transactions of the American Entomological Society 112: 191-248.
- LaBerge WE (1989) A revision of the bees of the genus *Andrena* of the western hemisphere, Part XIII. Subgenera *Simandrena* and *Taeniandrena*. Transactions of the American Entomological Society 115: 1-56.
- Leong JM, Thorp R (1999) Colour-coded sampling: the pan trap colour preferences of oligolectic and nonoligolectic bees associated with a vernal pool plant. Ecological Entomology 24 (3): 329-335. <https://doi.org/10.1046/j.1365-2311.1999.00196.x>
- Lichvar R, Banks DC, Kirchner WN, Melvin NC (2016) The National Wetland Plant List: 2016 wetland rating. Phytoneuron 2016: 1-17. <https://doi.org/10.21236/ada570149>
- Linsley EG, MacSwain JW (1958) The significance of floral constancy among bees of the genus *Diadasia* (Hymenoptera, Anthophoridae). Evolution 12 (2): 219-223. <https://doi.org/10.2307/2406032>
- Lippok B, Gardine A, Williamson P, Renner SS (2000) Pollination by Flies, Bees, and Beetles of *Nuphar ozarkana* and *N. advena* (Nymphaeaceae). American Journal of Botany 87 (6): 898-902. <https://doi.org/10.2307/2656897>
- Little CL (2013) Bee communities in the Arkansas River Valley. Master's thesis. University of Central, Arkansas, Conway, AR.
- Loose JL, Drummond FA, Stubbs C, Woods S, Hoffman S (2005) Conservation and management of native bees in cranberry. Maine Agricultural and Forest Experiment Station Technical Bulletin 191: 27 p.
- Michener CD (2007) The Bees of The World. 2nd edition. The John Hopkins University Press, Baltimore, MD, 953 pp.
- Minckley R, Wcislo W, Yanega D, Buchmann S (1994) Behavior and phenology of a specialist bee (*Dieunomia*) and sunflower (*Helianthus*) pollen Availability. Ecology 75 (5): 1406-1419. <https://doi.org/10.2307/1937464>
- Mitchell TB (1935) A revision of the genus *Megachile* in the nearctic region. Part II. Transactions of the American Entomological Society 61: 1-44.
- Mitchell TB (1937a) A revision of the genus *Megachile* in the nearctic region. Part VIII. Transactions of the American Entomological Society 63: 381-426.
- Mitchell TB (1937b) A revision of the genus *Megachile* in the nearctic region. Part VI. Transactions of the American Entomological Society 63: 45-83.

- Mitchell TB (1960) Bees of the eastern United States, Volume I. North Carolina Agricultural Experiment Station Technical Bulletin 141: 1-538.
- Mitsch WJ, Gosselink JG (2015) Wetlands. 5th. Wiley Press. , 747 pp.
- Moure JS, Hurd PD (1987) An annotated catalog of the halictid bees of the Western Hemisphere (Hymenoptera: Halictidae). Smithsonian Institution Press, Washington, D.C., 405 pp.
- Niering WA (1988) Endangered, threatened, and rare wetland plants and animals of the continental United States. The Ecology and Management of Wetlands. 1. Springer, Boston, MA, 592 pp.
- Norden BB, Krombein KV, Deyrup MA, Edirisinghe JP (2003) Biology and behavior of a seasonally aquatic bee, *Perdita (Alloperdita) floridensis* Timberlake (Hymenoptera: Andrenidae: Panurginae). Journal of the Kansas Entomological Society 76 (2): 236 -2249.
- Park C, Overall L, Smith L, Lagrange T, Mcmurry S (2017) Melittofauna and other potential pollinators in wetland and uplands in South Central Nebraska (Insecta: Apoidea). Zootaxa 4242 (2): 255-280. <https://doi.org/10.11646/zootaxa.4242.2.3>
- Pascarella JB, Waddington KD, Neal P (2000) The bee fauna (Hymenoptera: Apoidea) of Everglades National Park, Florida and adjacent areas: distribution, phenology, and biogeography. Journal of the Kansas Entomological Society 72 (1): 32-45.
- Rau P (1930) The nesting habits of *Emphor bombiformis* Cresson. Bulletin of the Brooklyn Entomological Society 25 (1): 28-35.
- Reader RJ (1975) Competitive relationships of some bog ericads for major insect pollinators. Canadian Journal of Botany 53 (13): 1300-1305. <https://doi.org/10.1139/b75-156>
- Rehan S, Sheffield C (2011) Morphological and molecular delineation of a new species in the *Ceratina dupla* species-group (Hymenoptera: Apidae: Xylocopinae) of eastern North America. Zootaxa 2873: 35-50.
- Rightmyer MG (2008) A review of the cleptoparasitic bee genus *Triepeolus* (Hymenoptera: Apidae). Part 1. Zootaxa 1710: 1-170.
- Roberts RB (1972) Revision of the bee genus *Agapostemon* (Hymenoptera: Halictidae). University of Kansas Science Bulletin 49: 437-590.
- Roulston T, Smith S, Brewster A (2007) A comparison of pan trap and intensive net sampling techniques for documenting a bee (Hymenoptera: Apiformes) fauna. Journal of the Kansas Entomological Society 80 (2): 179-181. [https://doi.org/10.2317/0022-8567\(2007\)80\[179:acopta\]2.0.co;2](https://doi.org/10.2317/0022-8567(2007)80[179:acopta]2.0.co;2)
- Rust RW (1980) The biology of *Ptilothrix bombiformis* (Hymenoptera: Anthophoridae). Journal of the Kansas Entomological Society 53 (2): 427-436.
- Scott HD, Ferguson JA, Hanson L, Fugitt T, Smith E (1998) Agricultural water management in the Mississippi Delta Region of Arkansas. Research Bulletin 959. <http://arkansasagnews.uark.edu/959.pdf>. Accessed on: 2017-4-08.
- Snelling RR (1970) Studies on North American bees of the genus *Hylaeus*, 5. The subgenera *Hylaeus* s. str. and *Paraprosopis*. Contributions in Science, Los Angeles County Museum of Natural History 180: 1-59.
- Snelling RR (1990) A review of the native North American bees of the genus *Chalicodoma* (Hymenoptera: Megachilidae). Contributions in Science, Los Angeles County Museum of Natural History 421: 1-3.

- Sohmer SH, Sefton DF (1978) The reproductive ecology of *Nelumbo pentapetala* (Nelumbonaceae) on the Upper Mississippi River. II. The insects associated with the transfer of pollen. *Brittonia* 30 (3): 355-364. <https://doi.org/10.2307/2806274>
- Stephenson PL (2017) Bee Communities on Managed Emergent Wetlands in the lower Mississippi Alluvial Valley of Arkansas. University of Arkansas, Fayetteville, AR. URL: <http://scholarworks.uark.edu/cgi/viewcontent.cgi?article=3966&context=etd>
- Stephen W, Rao S (2005) Unscented color traps for non-*Apis* bees (Hymenoptera: Apiformes). *Journal of the Kansas Entomological Society* 78 (4): 373-380. <https://doi.org/10.2317/0410.03.1>
- Stephen W, Rao S (2007) Sampling native bees in proximity to a highly competitive food resource (Hymenoptera: Apiformes). *Journal of the Kansas Entomological Society* 80 (4): 369-376. [https://doi.org/10.2317/0022-8567\(2007\)80\[369:snbipj\]2.0.co;2](https://doi.org/10.2317/0022-8567(2007)80[369:snbipj]2.0.co;2)
- Stephen WP (1954) A revision of the bee genus *Colletes* in America north of Mexico. *University of Kansas Science Bulletin* 36: 149-527.
- Timberlake PH (1969) A contribution to the systematics of North American species of *Synhalonia* (Hymenoptera: Apoidea). *University of California Publications in Entomology* 57: 76.
- Timberlake PH (1980) Review of North American *Exomalopsis* (Hymenoptera, Anthophoridae). *University of California Publications in Entomology* 86: 158.
- USDA National Agricultural Statistics Service (NASS) (2016) Cropland Data Layer (CDL)(2016) Published crop-specific data layer. <http://nassgeodata.gmu.edu/CropScape/>. Accessed on: 2017-4-21.
- USDA, NRCS (2017) The PLANTS National Plant Data Team, Greensboro, NC 27401-4901 USA. <http://plants.usda.gov>. Accessed on: 2017-8-11.



**Figure 1.**  
Distribution of managed palustrine emergent wetlands surveyed for bees in the lower Mississippi Alluvial Valley of Eastern Arkansas, USA in 2015 and 2016. See Table 1 for site names and coordinates.

Table 1.

Site number, site name, ownership, latitude, longitude, county, hectares and year surveyed during 2015 and 2016 in the lower Mississippi Alluvial Valley of Eastern Arkansas, USA.